



PRODUCT CATALOG

Pharmaceutical Products

in partnership with

World Medicine Pharmaceutical Company





ROFLOXAN

Ofloxacin is an antibacterial drug of the fluoroquinolone group (a quinolone-carboxylic acid derivative) that is active against a wide range of Gram-negative and Gram-positive microorganisms.

THERAPEUTIC INDICATIONS

Rofloxacin is indicated in adults for the treatment of the following bacterial infections caused by ofloxacin-susceptible microorganisms: Genitourinary infections. Pelvic inflammatory diseases (in combination with other antibacterials). Respiratory infections. Exacerbation of chronic obstructive pulmonary disease. Complicated skin and soft tissue infections.



SERAMOL

Omeprazole, a racemic mixture of two enantiomers, reduces gastric acid secretion through a highly targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. It is rapidly acting and provides control through reversible inhibition of gastric acid secretion with once-daily dosing.

Omeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the intracellular canaliculi within the parietal cell, where it inhibits the enzyme $H^+ K^+-ATPase$ - the acid pump. This effect on the final step of the gastric acid formation process is dose-dependent and provides for highly effective inhibition of both basal acid secretion and stimulated acid secretion, irrespective of stimulus.



SYRPIL

Paracetamol provides onset of pain relief within 5 to 10 minutes after the start of infusion. The peak analgesic effect is obtained in 1 hour and the duration of this effect is usually 4 to 6 hours.

Paracetamol reduces fever within 30 minutes after the start of infusion with a duration of the antipyretic effect of at least 6 hours.


THERAPEUTIC INDICATIONS

The drug is indicated for use in adults, adolescent and children weighing over 33 kg for:

- short-term treatment of moderate pain syndrome, especially following surgery;
- short-term treatment of pyrexia, when the intravenous route of administration is clinically justified by an urgent need to treat pain or hyperthermic syndrome and/or when other routes of administration are not possible.

METHOD OF ADMINISTRATION AND POSOLOGY

For intravenous infusion. Dosing based on patient weight.



ZIROMIN

Azithromycin is a semisynthetic azalide derivative with a 15-membered lactone ring; azalides belong to the group of macrolide antibiotics. The mechanism of action of azithromycin is based on inhibition of bacterial protein biosynthesis by binding to the 50S ribosomal subunit; the drug generally has a bacteriostatic effect.

THERAPEUTIC INDICATIONS

Ziromin is indicated for the treatment of infections caused by pathogens susceptible to azithromycin: Upper respiratory tract infections. Lower respiratory tract infections. Acute otitis media. Skin and soft tissue infections. Uncomplicated sexually transmitted infections caused by Chlamydia trachomatis or Neisseria gonorrhoeae (non-multiresistant strains).



LOXIDOL

Meloxicam is a nonsteroidal anti-inflammatory drug (NSAID) belonging to the oxycam family, which has anti-inflammatory, analgesic and antipyretic properties. The anti-inflammatory activity of meloxicam has been proven in classical models of inflammation. As with other NSAIDs, the exact mechanism of action remains unknown. However, there is at least one common mechanism of action typical for all NSAIDs (including meloxicam). This mechanism is to inhibit the synthesis of prostaglandins that are inflammatory mediators.

THERAPEUTIC INDICATIONS

Short-term symptomatic therapy of active arthrosis. Long-term symptomatic therapy of rheumatoid arthritis (chronic polyarthritis) or ankylosing spondylitis (ankylosing spondylitis).



MEDOLAPRAM

Escitalopram is a selective serotonin reuptake inhibitor (5-HT) with a high affinity for the primary binding site. It also binds to an allosteric site on the serotonin transporter, with a 1000-fold lower affinity. Escitalopram has no or low affinity for a number of receptors including 5-HT_{1A}- и 5-HT₂-receptors, dopamine D₁- and D₂-receptors, α₁-, α₂- и β-adrenoceptors, histamine H₁-receptors, m-cholinergic receptors, benzodiazepine and opioid receptors.

THERAPEUTIC INDICATIONS

Major depressive episodes. Panic disorder with or without agoraphobia. Social anxiety disorder (social phobia). Generalised anxiety disorder. Obsessive-compulsive disorder.



ONCECEF

Ceftriaxone (third-generation cephalosporins) inhibits bacterial cell wall synthesis following attachment to penicillin-binding proteins. This results in the interruption of cell wall (peptidoglycan) biosynthesis, which leads to bacterial cell lysis and death.

THERAPEUTIC INDICATIONS

Cefamed is indicated for the treatment of the following infections in adults and children including term neonates (from birth): Bacterial meningitis. Community-acquired pneumonia. Hospital-acquired pneumonia. Acute otitis media. Intra-abdominal infections. Complicated urinary tract infections. Bone and joint infections. Complicated skin and soft tissue infections. Gonorrhea. Syphilis. Bacterial endocarditis.



POMPEZO

Esomeprazole is a proton pump inhibitor (PPI); it suppresses stomach acid secretion by specific inhibition of the H⁺/K⁺-ATPase system in gastric parietal cells. S- and R-isomers of omeprazole are protonated and converted in the acidic compartment of parietal cells, forming an active inhibitor, the achiral sulfenamide. By acting specifically on the proton pump, esomeprazole blocks the final step in acid production, thus reducing gastric acidity. This effect (decreased hydrochloric acid secretion in the stomach) is dose dependent, ranging from 20 mg to 40 mg per day.



BRAVEX

Ibuprofen is a nonsteroidal anti-inflammatory agent, a propionic acid derivative. Ibuprofen inhibits prostaglandin synthesis, reducing inflammatory pain, swelling and fever. Furthermore, ibuprofen reversibly inhibits platelet aggregation.

THERAPEUTIC INDICATIONS

By doctor's prescription only: for symptomatic therapy of juvenile rheumatoid arthritis.

By or without doctor's prescription: for rapid and effective reduction of fever, including post-immunization hyperthermia, and for rapid and effective relief of symptoms of cold and influenza, mild to moderate pain, such as sore throat, teething pain, toothache, earache, headache, mild pain syndromes and pain from ligament sprains.



BRONOLAC

Ibuprofen is a nonsteroidal anti-inflammatory agent, a propionic acid derivative. Ibuprofen inhibits prostaglandin synthesis, reducing inflammatory pain, swelling and fever. Furthermore, ibuprofen reversibly inhibits platelet aggregation.

THERAPEUTIC INDICATIONS

By doctor's prescription only: for symptomatic therapy of juvenile rheumatoid arthritis.

By or without doctor's prescription: for rapid and effective reduction of fever, including post-immunization hyperthermia, and for rapid and effective relief of symptoms of cold and influenza, mild to moderate pain, such as sore throat, teething pain, toothache, earache, headache, mild pain syndromes and pain from ligament sprains.



CEFIDANT

Cefixime is a third-generation oral cephalosporin that has a pronounced in vitro bactericidal activity against a wide variety of gram-positive and gram-negative organisms. Clinical efficacy has been demonstrated in infections caused by commonly occurring pathogens.

THERAPEUTIC INDICATIONS

Cefidant is an orally active cephalosporin antibiotic which has a pronounced in vitro bactericidal activity against a wide variety of Gram-positive and Gram-negative organisms. It is indicated for the treatment of the following acute infections caused by susceptible microorganisms: Upper Respiratory Tract Infections. Lower Respiratory Tract Infection. Urinary Tract Infections.



COLEDAN

The biologically active form of vitamin D stimulates the absorption of calcium ions from the small intestine, osteoclastic resorption and the release of calcium from bone tissue. In the small intestine it promotes rapid and delayed calcium uptake. Vitamin D also stimulates the passive and active transport of phosphate. In the kidney, it inhibits the excretion of calcium and phosphate by promoting tubular resorption. The production of parathyroid hormone in the parathyroid gland is directly inhibited by the biologically active form of vitamin D3. PTH secretion is inhibited additionally by the increased calcium uptake in the small intestine under the influence of biologically active vitamin D3.

THERAPEUTIC INDICATIONS

The treatment and prevention of vitamin D deficiency in adolescents, adults and old patients.

As an adjunct to the specific therapy of osteoporosis in patients with vitamin D deficiency.

OSOLOGY AND METHOD OF ADMINISTRATION

Children

Prevention of vitamin D deficiency in children aged 12-18 years: 1 capsule (20000 IU) in every 6 weeks.

Treatment of vitamin D deficiency in children aged 12-18 years: 1 capsule (20000 IU) in every 2 weeks for 6 weeks.

Adults

Prevention of vitamin D deficiency: 1 capsule (20000 IU) once a month. Higher doses may be required in certain situations.

Treatment of vitamin D deficiency: 2 capsules (40000 IU) per week for 7 weeks followed by the maintenance therapy, for example 2-3 capsules (1400-2000 IU per day) per month. Subsequent measurements of 25-(OH) D should be taken approximately 3-4 months after the initiation of the maintenance therapy to confirm that the target level has been achieved.



EKSDELAN

Dexlansoprazole is the R-enantiomer of lansoprazole. It is a proton pump inhibitor that suppresses the final stage of gastric acid secretion by inhibiting the activity of the H⁺/K⁺-ATPase enzyme in the stomach parietal cells. The enzyme inhibition is dose-dependent and reversible and the effect extends to both basal and stimulated secretion of gastric acid. Dexlansoprazole is concentrated in parietal cells where it is converted to its active form in the acidic environment and inhibits the activity of H⁺/K⁺-ATPase by interacting with its sulfhydryl groups.



FARIKTAN

Fariktan is a combination drug. The efficacy of the combination is higher than the efficacy of its components alone.

- Ibuprofen is a propionic acid derivative, NSAID that has demonstrated its efficacy by inhibition of prostaglandin synthesis. Ibuprofen reduces inflammatory pain, swellings and fever.
- Paracetamol has analgesic and antipyretic effects similar to those of acetylsalicylic acid.

THERAPEUTIC INDICATIONS

Therapy of mild to moderate pain and inflammatory syndromes of various origins. Headache (including migraine). Toothache. Fever in infectious and inflammatory diseases.



FLAPROX

A broad-action antimicrobial fluoroquinolone drug. The bactericidal action results from inhibition of both topoisomerase II (DNA-gyrase) and topoisomerase IV required for bacterial DNA replication, transcription, repair, and recombination.

THERAPEUTIC INDICATIONS

The drug is indicated for the treatment of infections caused by ciprofloxacin-susceptible microorganisms. Lower respiratory tract infections. Chronic suppurative otitis media. Urinary tract infections. Genital tract infections. Gastrointestinal tract infections. Intra-abdominal infections. Skin and soft tissue infections caused by gram-negative bacteria. Malignant external otitis. Bone and joint infections. Prophylaxis of invasive infections caused by *Neisseria meningitidis*. Inhalation anthrax (post-exposure prophylaxis and curative treatment).



LEKOXA

Celecoxib, the active substance of Lekoxa, is an oral selective COX-2 inhibitor in the clinical dose range (200-400 mg daily). There have been several clinical studies confirming efficacy and safety in osteoarthritis, rheumatoid arthritis and ankylosing spondylitis.

THERAPEUTIC INDICATIONS

Lekoxa is indicated for symptomatic relief in the treatment of osteoarthritis, rheumatoid arthritis and ankylosing spondylitis in adults. The decision to prescribe a selective cyclooxygenase-2 (COX-2) inhibitor should be based on an assessment of the individual patient's overall risks.



LEVOXIMED

Levofloxacin is a synthetic antibacterial agent of the fluoroquinolone class and is the S (-) enantiomer of racemic ofloxacin. Mechanism of action: As a fluoroquinolone antibacterial agent, levofloxacin acts on the DNA-DNA-gyrase and topoisomerase IV complex.

THERAPEUTIC INDICATIONS

The drug is indicated in adults for the treatment of the following infections:

Acute pyelonephritis and complicated urinary tract infections. Chronic bacterial prostatitis. Inhalation anthrax: post-exposure prophylaxis and treatment.

