

NP-Completeness

Def. A problem X is called **NP-complete** if

- ① $X \in \text{NP}$, and
- ② $Y \leq_P X$ for every $Y \in \text{NP}$.

Theorem If X is NP-complete and $X \in \text{P}$, then $\text{P} = \text{NP}$.

- NP-complete problems are the hardest problems in NP
- NP-hard problems are at least as hard as NP-complete problems (a NP-hard problem is not required to be in NP)

Outline

- 1 Some Hard Problems
- 2 P, NP and Co-NP
- 3 Polynomial Time Reductions and NP-Completeness
- 4 NP-Complete Problems**
- 5 Dealing with NP-Hard Problems
- 6 Summary
- 7 Summary of Studies 2024 Spring

Def. A problem X is called **NP-complete** if

- 1 $X \in \text{NP}$, and
- 2 $Y \leq_P X$ for every $Y \in \text{NP}$.

Def. A problem X is called **NP-complete** if

- ① $X \in \text{NP}$, and
- ② $Y \leq_P X$ for every $Y \in \text{NP}$.

- How can we find a problem $X \in \text{NP}$ such that every problem $Y \in \text{NP}$ is polynomial time reducible to X ? Are we asking for too much?

Def. A problem X is called **NP-complete** if

- ① $X \in \text{NP}$, and
- ② $Y \leq_P X$ for every $Y \in \text{NP}$.


- How can we find a problem $X \in \text{NP}$ such that every problem $Y \in \text{NP}$ is polynomial time reducible to X ? Are we asking for too much?
- No! There is indeed a large family of natural NP-complete problems

The First NP-Complete Problem: Circuit-Sat

Circuit Satisfiability (Circuit-Sat)

Input: a circuit


Output: whether the circuit is satisfiable



Circuit-Sat is NP-Complete

- key fact: algorithms can be converted to circuits


Fact Any algorithm that takes n bits as input and outputs 0/1 with running time $T(n)$ can be converted into a circuit of size $p(T(n))$ for some polynomial function $p(\cdot)$.



Circuit-Sat is NP-Complete

- key fact: algorithms can be converted to circuits

Fact Any algorithm that takes n bits as input and outputs 0/1 with running time $T(n)$ can be converted into a circuit of size $p(T(n))$ for some polynomial function $p(\cdot)$.



- Then, we can show that any problem $Y \in \text{NP}$ can be reduced to Circuit-Sat.
- We prove $\text{HC} \leq_P \text{Circuit-Sat}$ as an example.


$\text{check-HC}(G, S)$

- Let $\text{check-HC}(G, S)$ be the certifier for the Hamiltonian cycle problem: $\text{check-HC}(G, S)$ returns 1 if S is a Hamiltonian cycle in G and 0 otherwise.

$\text{check-HC}(G, S)$


- Let $\text{check-HC}(G, S)$ be the certifier for the Hamiltonian cycle problem: $\text{check-HC}(G, S)$ returns 1 if S is a Hamiltonian cycle in G and 0 otherwise.
- G is a yes-instance if and only if there is an S such that $\text{check-HC}(G, S)$ returns 1

$HC \leq_P \text{Circuit-Sat}$




- Let $\text{check-HC}(G, S)$ be the certifier for the Hamiltonian cycle problem: $\text{check-HC}(G, S)$ returns 1 if S is a Hamiltonian cycle in G and 0 otherwise.
- G is a yes-instance if and only if there is an S such that $\text{check-HC}(G, S)$ returns 1
- Construct a circuit C' for the algorithm check-HC

$HC \leq_P \text{Circuit-Sat}$



- Let $\text{check-HC}(G, S)$ be the certifier for the Hamiltonian cycle problem: $\text{check-HC}(G, S)$ returns 1 if S is a Hamiltonian cycle in G and 0 otherwise.
- G is a yes-instance if and only if there is an S such that $\text{check-HC}(G, S)$ returns 1
- Construct a circuit C' for the algorithm check-HC
- hard-wire the instance G to the circuit C' to obtain the circuit C

$HC \leq_P \text{Circuit-Sat}$



- Let $\text{check-HC}(G, S)$ be the certifier for the Hamiltonian cycle problem: $\text{check-HC}(G, S)$ returns 1 if S is a Hamiltonian cycle in G and 0 otherwise.
- G is a yes-instance if and only if there is an S such that $\text{check-HC}(G, S)$ returns 1
- Construct a circuit C' for the algorithm check-HC
- hard-wire the instance G to the circuit C' to obtain the circuit C
- G is a yes-instance if and only if C is satisfiable

$Y \leq_P \text{Circuit-Sat}$, For Every $Y \in \text{NP}$


- Let $\text{check-}Y(s, t)$ be the certifier for problem Y : $\text{check-}Y(s, t)$ returns 1 if t is a valid certificate for s .
- s is a yes-instance if and only if there is a t such that $\text{check-}Y(s, t)$ returns 1
- Construct a circuit C' for the algorithm $\text{check-}Y$
- hard-wire the instance s to the circuit C' to obtain the circuit C
- s is a yes-instance if and only if C is satisfiable □

$Y \leq_P \text{Circuit-Sat}$, For Every $Y \in \text{NP}$

- Let $\text{check-}Y(s, t)$ be the certifier for problem Y : $\text{check-}Y(s, t)$ returns 1 if t is a valid certificate for s .
- s is a yes-instance if and only if there is a t such that $\text{check-}Y(s, t)$ returns 1
- Construct a circuit C' for the algorithm $\text{check-}Y$
- hard-wire the instance s to the circuit C' to obtain the circuit C
- s is a yes-instance if and only if C is satisfiable □

Theorem Circuit-Sat is NP-complete.

Reductions of NP-Complete Problems



3-CNF (conjunctive normal form) is a special case of formula:

3-CNF (conjunctive normal form) is a special case of formula:

- Boolean variables: x_1, x_2, \dots, x_n

3-CNF (conjunctive normal form) is a special case of formula:

- Boolean variables: x_1, x_2, \dots, x_n
- Literals: x_i or $\neg x_i$

3-CNF (conjunctive normal form) is a special case of formula:

- Boolean variables: x_1, x_2, \dots, x_n
- Literals: x_i or $\neg x_i$
- Clause: disjunction (“or”) of at most 3 literals: $x_3 \vee \neg x_4,$
 $x_1 \vee x_8 \vee \neg x_9, \quad \neg x_2 \vee \neg x_5 \vee x_7$

3-CNF (conjunctive normal form) is a special case of formula:

- Boolean variables: x_1, x_2, \dots, x_n
- Literals: x_i or $\neg x_i$
- Clause: disjunction (“or”) of at most 3 literals: $x_3 \vee \neg x_4, x_1 \vee x_8 \vee \neg x_9, \neg x_2 \vee \neg x_5 \vee x_7$
- 3-CNF formula: conjunction (“and”) of clauses:
 $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee \neg x_4)$

3-Sat

3-Sat

Input: a 3-CNF formula

Output: whether the 3-CNF is satisfiable

3-Sat

Input: a 3-CNF formula

Output: whether the 3-CNF is satisfiable

- To satisfy a 3-CNF, we need to satisfy all clauses

3-Sat

Input: a 3-CNF formula

Output: whether the 3-CNF is satisfiable

- To satisfy a 3-CNF, we need to satisfy all clauses
- To satisfy a clause, we need to satisfy at least 1 literal


3-Sat

Input: a 3-CNF formula


Output: whether the 3-CNF is satisfiable

- To satisfy a 3-CNF, we need to satisfy all clauses
- To satisfy a clause, we need to satisfy at least 1 literal
- Assignment $x_1 = 1, x_2 = 1, x_3 = 0, x_4 = 0$ satisfies $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee \neg x_4)$

Circuit-Sat \leq_P 3-Sat




Circuit-Sat \leq_P 3-Sat



- Associate every wire with a new variable

Circuit-Sat \leq_P 3-Sat



- Associate every wire with a new variable
- The circuit is equivalent to the following formula:

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

$$(x_1 \vee \neg x_2 \vee x_5) \quad \wedge$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

$$(x_1 \vee \neg x_2 \vee x_5) \quad \wedge$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

$$(x_1 \vee \neg x_2 \vee x_5) \quad \wedge$$

$$(\neg x_1 \vee x_2 \vee x_5) \quad \wedge$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

$$(x_1 \vee \neg x_2 \vee x_5) \quad \wedge$$

$$(\neg x_1 \vee x_2 \vee x_5) \quad \wedge$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

$$\begin{aligned} & (x_4 = \neg x_3) \wedge (x_5 = x_1 \vee x_2) \wedge (x_6 = \neg x_4) \\ & \wedge (x_7 = x_1 \wedge x_2 \wedge x_4) \wedge (x_8 = x_5 \vee x_6) \\ & \wedge (x_9 = x_6 \vee x_7) \wedge (x_{10} = x_8 \wedge x_9 \wedge x_7) \wedge x_{10} \end{aligned}$$

Convert each clause to a 3-CNF

$$x_5 = x_1 \vee x_2 \quad \Leftrightarrow$$

$$(x_1 \vee x_2 \vee \neg x_5) \quad \wedge$$

$$(x_1 \vee \neg x_2 \vee x_5) \quad \wedge$$

$$(\neg x_1 \vee x_2 \vee x_5) \quad \wedge$$

$$(\neg x_1 \vee \neg x_2 \vee \neg x_5)$$

| x_1 | x_2 | x_5 | $x_5 \leftrightarrow x_1 \vee x_2$ |
|-------|-------|-------|------------------------------------|
| 0 | 0 | 0 | 1 |
| 0 | 0 | 1 | 0 |
| 0 | 1 | 0 | 0 |
| 0 | 1 | 1 | 1 |
| 1 | 0 | 0 | 0 |
| 1 | 0 | 1 | 1 |
| 1 | 1 | 0 | 0 |
| 1 | 1 | 1 | 1 |

Circuit-Sat \leq_P 3-Sat

- Circuit \iff Formula \iff 3-CNF

Circuit-Sat \leq_P 3-Sat

- Circuit \iff Formula \iff 3-CNF
- The circuit is satisfiable if and only if the 3-CNF is satisfiable


Circuit-Sat \leq_P 3-Sat

- Circuit \iff Formula \iff 3-CNF
- The circuit is satisfiable if and only if the 3-CNF is satisfiable
- The size of the 3-CNF formula is polynomial (indeed, linear) in the size of the circuit

Circuit-Sat \leq_P 3-Sat


- Circuit \iff Formula \iff 3-CNF
- The circuit is satisfiable if and only if the 3-CNF is satisfiable
- The size of the 3-CNF formula is polynomial (indeed, linear) in the size of the circuit
- Thus, Circuit-Sat \leq_P 3-Sat

Reductions of NP-Complete Problems



Recall: Independent Set Problem

Def. An **independent set** of $G = (V, E)$ is a subset $I \subseteq V$ such that no two vertices in I are adjacent in G .



Independent Set (Ind-Set) Problem

Input: $G = (V, E), k$


Output: whether there is an independent set of size k in G

3-Sat \leq_P Ind-Set

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$


3-Sat \leq_P Ind-Set

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- A clause \Rightarrow a group of 3 vertices, one for each literal
- An edge between every pair of vertices in same group




3-Sat \leq_P Ind-Set

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- A clause \Rightarrow a group of 3 vertices, one for each literal
- An edge between every pair of vertices in same group
- An edge between every pair of contradicting literals




3-Sat \leq_P Ind-Set

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- A clause \Rightarrow a group of 3 vertices, one for each literal
- An edge between every pair of vertices in same group
- An edge between every pair of contradicting literals
- Problem: whether there is an IS of size $k = \# \text{clauses}$



3-Sat \leq_P Ind-Set


- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- A clause \Rightarrow a group of 3 vertices, one for each literal
- An edge between every pair of vertices in same group
- An edge between every pair of contradicting literals
- Problem: whether there is an IS of size $k = \# \text{clauses}$



3-Sat instance is yes-instance \Leftrightarrow Ind-Set instance is yes-instance:

3-Sat \leq_P Ind-Set

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- A clause \Rightarrow a group of 3 vertices, one for each literal
- An edge between every pair of vertices in same group
- An edge between every pair of contradicting literals
- Problem: whether there is an IS of size $k = \# \text{clauses}$




3-Sat instance is yes-instance \Leftrightarrow Ind-Set instance is yes-instance:

- satisfying assignment \Rightarrow independent set of size k
- independent set of size $k \Rightarrow$ satisfying assignment


Satisfying Assignment \Rightarrow IS of Size k

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$




Satisfying Assignment \Rightarrow IS of Size k

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every clause, at least 1 literal is satisfied




Satisfying Assignment \Rightarrow IS of Size k

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every clause, at least 1 literal is satisfied
- Pick the vertex correspondent the literal




Satisfying Assignment \Rightarrow IS of Size k

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every clause, at least 1 literal is satisfied
- Pick the vertex correspondent the literal
- So, 1 literal from each group




Satisfying Assignment \Rightarrow IS of Size k

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every clause, at least 1 literal is satisfied
- Pick the vertex correspondent the literal
- So, 1 literal from each group
- No contradictions among the selected literals




Satisfying Assignment \Rightarrow IS of Size k

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every clause, at least 1 literal is satisfied
- Pick the vertex correspondent the literal
- So, 1 literal from each group
- No contradictions among the selected literals
- An IS of size k




IS of Size $k \Rightarrow$ Satisfying Assignment

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$




IS of Size $k \Rightarrow$ Satisfying Assignment

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every group, exactly one literal is selected in IS




IS of Size $k \Rightarrow$ Satisfying Assignment

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every group, exactly one literal is selected in IS
- No contradictions among the selected literals




IS of Size $k \Rightarrow$ Satisfying Assignment

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every group, exactly one literal is selected in IS
- No contradictions among the selected literals
- If x_i is selected in IS, set $x_i = 1$



IS of Size $k \Rightarrow$ Satisfying Assignment

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every group, exactly one literal is selected in IS
- No contradictions among the selected literals
- If x_i is selected in IS, set $x_i = 1$
- If $\neg x_i$ is selected in IS, set $x_i = 0$



IS of Size $k \Rightarrow$ Satisfying Assignment

- $(x_1 \vee \neg x_2 \vee \neg x_3) \wedge (x_2 \vee x_3 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee x_4)$
- For every group, exactly one literal is selected in IS
- No contradictions among the selected literals
- If x_i is selected in IS, set $x_i = 1$
- If $\neg x_i$ is selected in IS, set $x_i = 0$
- Otherwise, set x_i arbitrarily

