

Dementia and its treatments

UoM April 2026

Dr. Ross A. Dunne
GMDRC

2026-04-26

Dementia -  treatments

Slido - navigate to the page on your



⚠ Q & A is open now - submit questions for me or for our guest



Slido Link

Outline

- 20 minutes on “Dementia”, to give a modern framework for understanding
- 20 minutes on the development of symptomatic treatments
- 5 minute break
- 20 minutes on “Disease modifying” treatments.
- 5 minutes for questions for me on the material presented
- 10-20 minutes Q and A trial participant with me
- 10 minutes Qs from students

Part 1: Dementia

A modern framework

Alzheimer's disease is not synonymous with dementia

Dementia is an umbrella term used to describe a state where an individual has lost the ability to carry out daily life activities independently, because of cognitive impairment.

There are many types and causes of dementia.


The most common ones are:

1. Alzheimer's disease (AD) **pathological definition**
2. Vascular dementia **aetiological definition**
3. Dementia with Lewy bodies **molecular definition**
4. Frontotemporal dementia **anatomical definition**

Nature Neuroscience overview

Question 1

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode

Screen size



Alzheimer's disease is a disease like any other

- AD refers to the abnormal presence of A β and tau proteins which define AD among many other neurodegenerative diseases
- Some individuals may have dementia without A β or tau pathology, while others (rarely) may have no clinical symptoms, even in the presence of A β pathology.

INTERACTIVE TIMELINE

A History of *Alzheimer's Disease*

From plaques and tangles to disease-modifying therapy · 1906 — 2023

0 / 13

← back

forward →

reveal all

reset

→ OR SPACE REVEAL NEXT · ← STEP BACK · A REVEAL ALL · R RESET · CLICK ANY YEAR TO JUMP



ALOIS ALZHEIMER · AUGUSTE DETER · AD RESEARCH MILESTONES

NUMERIC SUPERSCRIPTS DENOTE PRIMARY REFERENCES IN THE SOURCE FIGURE

Biological vs clinical definition

The old way pre-2010

- Clinical syndrome → “probable AD”
- Pathology only confirmed at post-mortem
- Heterogeneous cohorts in trials
- “*Dementia of the Alzheimer type*”

The modern way 2018 NIA-AA, 2024 revised criteria


- Biology defines the disease
 $A\beta$ + tau
- Clinical stage is described separately
- Biomarkers in life: CSF, plasma, PET
- Allows pre-symptomatic diagnosis

INTERACTIVE MODEL

The Biomarker Cascade of Alzheimer's Disease

After Jack et al., Lancet Neurology 2010 and 2013. Drag the cursor across the plot to explore the disease continuum.

DRAG OR CLICK THE CHART · ← / → FINE ADJUST · 1 - 6 JUMP TO STAGE · CLICK A BIOMARKER ROW TO TOGGLE ITS CURVE



POSITION ON CONTINUUM

Preclinical AD

A+ T- N-

x = 0.300

BIOMARKERS click to toggle

	Amyloid-β CSF Aβ42/40 · AMYLOID-PET	75% +
	p-tau CSF P-TAU · TAU-PET	38% -
	Neurodegeneration MRI ATROPHY · FDG-PET	14% -
	Cognition NEUROPSYCHOLOGICAL TESTING	5% -
	Function ACTIVITIES OF DAILY LIVING	2% -

CLINICAL INTERPRETATION Biomarker evidence of AD pathology without cognitive symptoms. Aβ deposition precedes detectable cognitive change by approximately 15 to 20 years.


STAGE: 1 Normal 2 Preclinical 3 SCD 4 MCI 5 Mild AD 6 Moderate

JACK ET AL. 2010, 2013 · NIA-AA RESEARCH FRAMEWORK, 2018

Dementia - 🎨 treatments

Question 2, Higher or Lower?

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode

Screen size



100%



The AT(N)(X) framework

A: Amyloid T: Tau N: Neurodegeneration X: Other

What it measures: cerebral A β pathology.


- CSF A β 42/40 ratio **low = abnormal**
- Plasma A β 42/40 **replaces some lumbar punctures**
- Amyloid-PET **florbetapir, flutemetamol**



click to reveal: lumbar puncture (LP is painless)

Question 3

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode


Screen size



The global prevalence of dementia is increasing

- There are many causes of dementia, but AD is the most common cause in individuals older than 65 years of age.
- In 2020, it was estimated that over 55 million people worldwide were living with neurodegenerative disease at the *dementia* stage.
- The number of people affected is set to rise to **152 million** by 2050, with the most significant increases in lower-middle-income (LMIC) countries.
- Dementia (including dementia due to AD) incidence and prevalence are increasing globally but are likely to be underestimated because of underdiagnosis and misdiagnosis
- As global lifespan continues to increase, the number of people affected by dementia is expected to rise. Up to three-quarters of those with dementia worldwide have not received a diagnosis.


57.4M to 153M by 2050



Global change in dementia prevalence from: Figure 4 Decomposition of percentage change in the number of individuals with dementia between 2019 and 2050 globally and by world region

GBD=Global Burden of Diseases, Injuries, and Risk Factors Study; *Lancet PH* Volume 7, Issue 2e105-e125 February 2022

Regions of greatest change



Estimated change per region *Lancet PH* Volume 7, Issue 2e105-e125 February 2022

Dementia -  treatments

Historical development of Alzheimer's disease diagnostic criteria

- The 2023 revised criteria for diagnosis and staging of AD emphasizes that AD is a disease defined by its biology
- However, to avoid misdiagnosis, only biomarkers that have been proven to be accurate with respect to an accepted reference standard should be used for diagnosis
- It is important to note that biomarker tests should always be interpreted with clinical judgment

The clinician's expertise is crucial in the following situations:

1. When the biomarker test result contradicts the clinical presentation
2. When evaluating the probable contribution of AD versus other pathologies to clinical symptoms
3. When assessing how other medical conditions could possibly impact the biomarker results

Alzheimer's disease diagnostic biomarkers


Core 1 currently accepted for diagnosis

- CSF A β 42/40
- CSF p-tau181/A β 42
- CSF t-tau/A β 42
- Accurate plasma assays
- Amyloid-PET

Core 2 emerging / research-grade

- Plasma p-tau217 near-CSF accuracy
- Plasma MTBR-tau243
- GFAP astrocyte activation
- NfL non-specific neurodegeneration
- Tau-PET Braak staging in life

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode

Screen size

Differential diagnosis

AD

DLB

FTD

VaD


- **Onset**: insidious, episodic memory first
- **Signature**: temporo-parietal cognition; language; getting lost
- **Imaging**: medial temporal and parietal atrophy
- **Treatment cue**: AChEi and memantine; now anti-amyloid eligible

Reversible mimics

Treatable contributors to cognitive impairment

- B12 / folate deficiency
- Hypothyroidism
- Depression *pseudo-dementia*
- Normal pressure hydrocephalus *gait, urinary, cognitive*
- Drug effects *anticholinergics, benzodiazepines, opioids*
- Delirium *often superimposed*
- Obstructive sleep apnoea
- Subdural haematoma
- Alcohol misuse

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode

Screen size ☐ 100% ☐

Assessment

Clinical History

Cognitive examination

Examination

Neuroimaging


Biomarkers

A medical history from the patient and collateral history from an appropriate informant.

- Educational attainment **quantitative and qualitative**
- Occupational attainment **manual/ skilled/ professional**
- Head Injury / Contact sports **rugby and boxing**
- Lifetime Alcohol + substances **peak, sustained**
- Smoking history
- Rapid forgetfulness
- Repetitive questioning
- Loss of recent memory **with relative preservation of remote**
- Functional decline **IADLs first, then ADLs**
- BPSD **agitation, depression, sleep**

Part 2: Symptomatic treatments

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode


Screen size



The cholinergic hypothesis (1976)

- Bowen, Davies, Perry independently observed loss of choline acetyltransferase in post-mortem AD brain
- Source: degenerating cholinergic projections from the **nucleus basalis of Meynert**
- Implication: boost acetylcholine, improve cognition

AChE inhibitor mechanism



Cholinergic synapse

- AD brains have loss of cortical and hippocampal ACh
- AChEi inhibit synaptic breakdown of ACh, increasing transmitter availability per release
- Effect is presynaptic compensation; does not modify disease
- Side-effects from cholinergic excess: ↑ vagal tone, GI cramping, bradycardia, vivid dreams

AChE inhibitors: clinical use


Donepezil

Rivastigmine

Galantamine

- Once daily, oral; 5 mg increased to 10 mg after 6 weeks **most common UK first-line**
- Long half-life
- Side effects: GI upset, vivid dreams, bradycardia
- Cardiac history → ECG before starting

Memantine: NMDA receptor antagonism



- Glutamatergic excitotoxicity → tonic NMDA-R activation → calcium overload
- Memantine sits in the channel only when it's pathologically open **voltage-dependent**
- Strong physiological NMDA bursts displace it, preserving physiological learning
- **Indication:** moderate to severe AD, or AChEi-intolerant
- Often combined with donepezil


Memantine mechanism of action

Effect size of symptomatic treatment

MMSE trajectory by treatment and dementia type

Illustrative curves parameterised from Knight 2018, Xu 2021, Xu 2024, Truzzi 2022. Baseline MMSE 22 for AD, 24 for DLB.

— Untreated AD - - - AChEI in AD AChEI + memantine, moderate AD — AChEI in DLB - - - Untreated DLB



Curves on AChEI run roughly parallel to the untreated trajectory in AD, offset by ~1 MMSE point. The DLB curve is meaningfully larger and more durable.

Differential responsiveness across subtypes

Helpful in

- **AD** modest, all severities
- **DLB** often substantial; rivastigmine first choice
- **PDD** cholinergic deficit prominent
- **VaD** mixed disease often benefits

Limited or harmful in

- **bvFTD** no benefit, may worsen agitation
- **Pure VaD** inconsistent
- **Severe AD** memantine takes over

BPSD

- Behavioural and Psychological Symptoms of Dementia
- Up to 90% of patients experience at least one over the course of illness
- Drives caregiver burden, institutionalisation, mortality
- Five clusters:
 1. Agitation / aggression
 2. Psychosis delusions, hallucinations
 3. Depression / anxiety
 4. Apathy
 5. Sleep disturbance

Non-pharmacological management of BPSD

- **Always rule out a precipitant:** pain, infection (UTI, chest), constipation, drug effect, sensory deprivation
- **Cognitive Stimulation Therapy** NICE-recommended, group-based
- **Exercise** aerobic + resistance
- **Music therapy:** strong evidence in agitation
- **Environmental modification:** light, routine, signage, familiar objects
- **Caregiver education and support:** often high-yield

Pharmacological management of BPSD

! Antipsychotics are last-resort


- ↑ stroke and all-cause mortality CATIE-AD, FDA black box
 - **Severe sensitivity in DLB**: can cause irreversible parkinsonism / NMS
 - Use only when behaviour endangers patient or others, lowest dose, short duration, regular review
-
- **SSRIs** for depression (citalopram has agitation evidence, but QTc concern)
 - **Brexpiprazole**: first FDA approval for agitation in AD, 2023 not yet NICE-approved on NHS for this indication
 - **Trazodone**: sleep, agitation; mostly off-label
 - **Avoid benzodiazepines**: falls, paradoxical agitation, dependence

Limits of symptomatic treatment

- They do not slow neuronal loss
- They do not clear A β or tau
- They do not change underlying trajectory
- A patient on AChEi 5 years from diagnosis has the same brain pathology as one untreated

What is important in clinical trials?

Join at
slido.com
#2777 658



Copy joining link

Switch to Host mode

Screen size



5 minute break  

Part 3: Disease-modifying treatments

The amyloid hypothesis (Hardy & Higgins, 1992)

- **A β accumulation** is the upstream driver of AD
- Genetic anchor: every gene mutation that causes autosomal-dominant AD **APP**, **PSEN1**, **PSEN2** alters A β production or clearance
- Down syndrome (trisomy 21 $\rightarrow$ 3 copies of **APP**) $\rightarrow$ near-universal AD pathology by age 50
- Tau pathology, inflammation, neurodegeneration: downstream
- Predicted: lower A β $\rightarrow$ less downstream damage $\rightarrow$ better outcome

Failed anti-amyloid trials

- **Solanezumab**: Eli Lilly; failed Phase 3 in mild AD (2014, 2016)
- **Bapineuzumab**: Pfizer/J&J; failed (2012); high rate of vasogenic edema
- **Crenezumab**: Roche; failed (2019)
- **Gantenerumab**: Roche; first GRADUATE trials negative (2022); reformulated for prevention
- Lessons learned:
 - Treating too late **pathology has spread for 15+ years**
 - Targeting the wrong species **monomer vs oligomer vs plaque**
 - Insufficient brain penetration / dosing
 - Endpoint too noisy in mild populations

Aducanumab (2021)

- Two Phase 3 trials (EMERGE, ENGAGE): discordant; pre-specified analysis was negative
- Post-hoc reanalysis at high dose suggested cognitive benefit on EMERGE
- **FDA accelerated approval** June 2021, over near-unanimous advisory panel objection
- EMA **refused** authorisation
- NHS / NICE: **not commissioned**
- Eli Lilly withdrew it from market 2024

Lecanemab: CLARITY-AD (2023)

Phase 3 · n = 1,795 · 18 months · early symptomatic AD, biomarker-confirmed · target: soluble Aβ protofibrils

EFFICACY · CDR-SB CHANGE FROM BASELINE

The graph shows the change in CDR-SB score from baseline (0) to 18 months. The placebo group (red line) shows a decline to 1.66, while the lecanemab group (blue line) shows a decline to 1.21. The difference is 0.45 points, representing a 27% slowing of decline.

Group	CDR-SB Change from Baseline (18 mo)
placebo	1.66
lecanemab	1.21

27% slowing
 -0.45 CDR-SB difference · p < 0.001 primary & all secondaries

SAFETY · ARIA-E RATE BY APOE GENOTYPE

The bar chart shows the ARIA-E rate by APOE genotype. The rates are approximately 5% for noncarriers, 11% for heterozygotes, and 33% for homozygotes.

Genotype	ARIA-E Rate
noncarrier	~5%
heterozygote	~11%
homozygote	~33%

12.6% / 17.3%
 overall ARIA-E / ARIA-H · 9.2% symptomatic ARIA-E in homozygotes

REGULATORY

FDA · Jul 2023
 full (traditional) approval

EMA · 2025
 approved on appeal, excluding APOE4 homozygotes


NICE · rejected
 NHS use, Aug 2024 — not cost-effective at list price

van Dyck CH et al. *N Engl J Med.* 2023;388(1):9–21

Donanemab: TRAILBLAZER-ALZ 2 (2024)

Phase 3 · n = 1,736 · 76 weeks · MCI / mild AD · target: pyroglutamate-modified Aβ in plaques · stop-when-cleared design


EFFICACY · IADRS DECLINE,
LOW/INTERMEDIATE-TAU (N = 1,182)



35% slowing

iADRS · CDR-SB slowing 36% · 47% vs 29% no decline at 1 y

SAFETY · ARIA IN DONANEMAB ARM VS
PLACEBO



5.8% / 1.5%

symptomatic / serious ARIA-E · 3 deaths attributed to ARIA

REGULATORY · APOE

FDA · Jul 2024

full approval

NICE · rejected


NHS use, Oct 2024

OR 4.57

ARIA-E in APOE4 homozygotes vs noncarriers

Sims JR et al. JAMA. 2023;330(6):512–527

Antibody specificity



A β aggregation pathway 10.1002/chem.202400277

Dementia -  treatments

Patient selection

- **Confirmed A β pathology**: amyloid-PET or CSF A β 42/40
- **Disease stage**: MCI or mild AD dementia (not moderate or severe)
- **MRI suitable**: no severe cerebral amyloid angiopathy, ≤ 4 microbleeds, no superficial siderosis
- **Anticoagulation**: caution or contraindication
- **APOE4 genotyping**: needed for ARIA risk counselling
- **Capable of attending infusions** every 2–4 weeks for 18+ months
- **Capacity** to consent

APOE RISK TOOL APOE Genotype and Alzheimer's Disease Risk

Verify against source PDFs

SOURCE Farrer 1997 Genin 2011 Reiman 2020 Fortea 2024 SEX All F M AGE 65 70 75 80 85 TONE Clinical Patient

Farrer LA et al., JAMA 1997;278:1349–56. Meta-analysis, Caucasian clinic/autopsy cohorts. ORs vs ε3/ε3, age- and study-adjusted.

1 Build a genotype

One allele from each biological parent. Drag (or click) one allele from each row.

P1

ε2ε3ε4

→

P2

ε2ε3ε4

→

YOUR GENOTYPE

— pick one from each parent —

Clear

Canonical order

2 How it is inherited

Set each parent's genotype. The four cells show the possible genotypes of any one child.

PARENT 1

ε3ε4

tap to change

PARENT 2

ε3ε3


tap to change

	ε3	ε3
ε3	ε3 / ε3 25%	ε3 / ε3 25%
ε4	ε3 / ε4 25%	ε3 / ε4 25%


Each child: 50% ε3/ε3, 50% ε3/ε4.

3 What this means for risk

Population frequency



Relative risk vs ε3/ε3



Absolute lifetime risk

Build a genotype to see absolute lifetime risk.

Context. Figures derive from cohorts of predominantly European ancestry. Risk is age-dependent. APOE is one of many genetic and environmental contributors; modifiable factors (hypertension, hearing loss, smoking, diabetes, social isolation) also shape lifetime risk. Genotype is probabilistic, not diagnostic.

ARIA

ARIA-E oedema, effusion

- Vasogenic oedema, leptomeningeal effusion
- Best seen on **FLAIR**
- Most asymptomatic; caught on surveillance MRI
- Can present with headache, confusion, focal symptoms, seizures
- Severity is mostly mild; ~3% serious; rare deaths reported
- Surveillance MRI before infusions 5, 7, 14 (lecanemab) or before 2, 3, 4, 7 (donanemab)

ARIA-H haemorrhage

- Microbleeds, superficial siderosis
- Best seen on **SWI / GRE**
- Usually asymptomatic
- Persists indefinitely after resolution

APOE4 and ARIA risk

APOE genotype	Lecanemab ARIA-E	Donanemab ARIA-E
Non-carrier	~5%	~16%
Heterozygote	~11%	~23%
Homozygote	~33%	~41%

- ϵ 4 homozygotes are ~2% of the population but bear most of the risk
- **Counselling**: genotype results have implications beyond the individual **family members, life insurance**
- Donanemab US label restricts use in homozygotes; EMA mandates discussion of relative risk
- Patients can decline genotyping but must be counselled to the higher-risk assumption

UK implementation

- **Drug cost:** £20–25k per patient per year (lecanemab list)
- **Infrastructure:** infusion suite every 2 (lecanemab) or 4 (donanemab) weeks
- **Surveillance MRI:** 4–7 scans per patient course
- **Workforce:** trained nurses, dementia specialists, neuroradiologists, raters, pharmacists
- **NICE August 2024:** rejected lecanemab for NHS commissioning; benefit too small at price
- Currently available only privately in the UK; major equity concern
- Open question whether amyloid lowering alone is sufficient or whether combinations will be needed

Other approaches

- **Anti-tau antibodies:** E2814, JNJ-2056 (zaganemab discontinued 2023) Phase 2
- **Tau ASOs** intrathecal antisense oligonucleotides, BIIB080
- **Inflammation:** TREM2 agonists in early trials
- **Vaccines:** UB-311, ABvac40 (Phase 2)
- **Prevention trials** in pre-symptomatic carriers:
 - **DIAN-TU:** autosomal-dominant AD families
 - **AHEAD 3-45:** sporadic, plasma p-tau217 stratified
 - *A4 closed early; solanezumab failed*

Current state

- The first disease-modifying drugs exist. Their effect size is modest. ARIA limits use. Access in the UK is restricted.
- **Diagnosis**: plasma p-tau217 may make primary-care testing feasible.
- **Prevention** may matter more at a population level than any antibody:
 - *Lancet Commission 2024: 14 modifiable risk factors account for ~45% of dementia risk*
 - Hearing, education, BP, smoking, obesity, depression, physical inactivity, diabetes, alcohol, social isolation, air pollution, head injury, vision, cholesterol

What is a Brain Health Centre?


Same patient. Same disease. Different future.

A Brain Health Centre is not a relabelled memory clinic. It redesigns the pathway so that biomarkers, vascular workup, and behavioural support arrive together, early, before too much brain has been lost.

Mary, 64. "My memory's not what it was. My dad had dementia. Should I be worried?"

MEMORY ASSESSMENT SERVICE

cognition-led, no biomarkers, sequential appointments




"probable Alzheimer's disease"

- × no biomarkers, no diagnostic certainty
- × ineligible for disease-modifying therapy
- × no structured risk-reduction plan

BRAIN HEALTH CENTRE

biomarker-led, integrated, with active follow-up



biomarker-confirmed diagnosis

- ✓ blood pressure normalised, statin started
- ✓ hearing aid referral, social plan
- ✓ DMT-ready, in research, year-on review booked

1 Confirm with biology, not exclusion. Plasma pTau217, MRI and CSF give a positive diagnosis of Alzheimer's pathology, instead of a diagnosis of last resort.

2 Treat the brain like the heart. Forty-five percent of dementia worldwide is linked to fourteen modifiable risk factors. Most are vascular and behavioural.

3 Be ready for therapy. Lecanemab and donanemab need confirmed amyloid. Without biomarkers, eligibility is impossible to establish.

Today only **1 in 800** UK patients with memory symptoms gets a biomarker-confirmed diagnosis. The other 799 are Mary on the upper track.



Manchester Brain Health Centre

GMMH

mental health · old-age psychiatry · clinical lead


MFT

imaging · cardiology · labs · biobank host

academic partner: The University of Manchester

Alzheimer's Society

funding · Brain Health Advisors · public voice



49_d

simulated median pathway,
referral to disclosure

vs 187 days currently

73%

reduction in time to diagnosis

Phase 2 vs Phase 1b

~80%

of patients spared a lumbar
puncture

blood-first triage

0

serious LP adverse events to
date

~92% acceptance, atraumatic technique



What's next, and why it matters

45% of dementia worldwide is linked to **14 modifiable risk factors** (Lancet Commission, 2024)

EARLY LIFE	MIDLIFE	LATE LIFE
Less education 5%	Hearing loss 7% High LDL cholesterol 7% Depression 3% Traumatic brain injury 3% Physical inactivity 2% Smoking 2% Diabetes 2% Hypertension 2% Obesity 1% Excessive alcohol 1%	Social isolation 5% Air pollution 3% Vision loss 2%

What's next at MBHC

- One-stop blood-first pathway live in clinic from **Q3 2026**
- First UK site to deploy **pTau217 Elecsys** for routine NHS care
- Brain Health Advisors trained in **motivational interviewing**, embedded in pathway
- Five-year cohort study with **GSK**, ASL imaging endpoints for future trials
- Hub-and-spoke model across **Greater Manchester**

Why it matters for psychology

- Pathology starts **20+ years** before symptoms. Behaviour change has the most leverage when started early.
- Disease-modifying drugs (lecanemab, donanemab) are MHRA-licensed but **not yet on the NHS**. Eligibility depends on biomarkers.
- Most modifiable risk factors are **psychological and behavioural**: hearing aid use, exercise, sleep, social engagement, mood.
- This is where psychologists work, with clinicians, on the things that actually shift the curve.



Questions on the material (5 min)



Q&A with trial participant (10–20 min)

Welcome!



Student questions (10 min)

over to you

Summary

1. **Dementia is a syndrome; AD is defined biologically.** $A\beta$ and tau can be detected in life via CSF, plasma p-tau217, or PET.
2. **Symptomatic treatments help modestly and do not change disease course.** AChEi and memantine offer about a 6-month delay. Non-pharmacological BPSD interventions are first-line. Antipsychotics carry mortality risk.
3. **Anti-amyloid antibodies are the first disease-modifying drugs.** Effect size is modest. ARIA limits use. NICE has rejected them for the NHS. Prevention may matter more.

References

- Bowen, David M., Charles B. Smith, Peter White, and Alan N. Davison. 1976. 'Neurotransmitter-Related Enzymes and Indices of Hypoxia in Senile Dementia and Other Abiotrophies'. *Brain* 99 (3): 459–96.
- Dyck, Christopher H. van, Chad J. Swanson, Paul Aisen, Randall J. Bateman, Christopher Chen, Michelle Gee, Michio Kanekiyo, et al. 2023. 'Lecanemab in Early Alzheimer's Disease'. *New England Journal of Medicine* 388 (1): 9–21.
- Hardy, John A., and Gerald A. Higgins. 1992. 'Alzheimer's Disease: The Amyloid Cascade Hypothesis'. *Science* 256 (5054): 184–85.
- Jack, Clifford R., David A. Bennett, Kaj Blennow, Maria C. Carrillo, Billy Dunn, Samantha Budd Haeberlein, David M. Holtzman, et al. 2018. 'NIA-AA Research Framework: Toward a Biological Definition of Alzheimer's Disease'. *Alzheimer's & Dementia* 14 (4): 535–62.
- Livingston, Gill, Jonathan Huntley, Kathy Y. Liu, Sergi G. Costafreda, Geir Selbæk, Suvarna Alladi, David Ames, et al. 2024. 'Dementia Prevention, Intervention, and Care: 2024 Report of the Lancet Standing Commission'. *The Lancet* 404 (10452): 572–628.
- Niu, Zheng, Xinrui Gui, Shuang Feng, and Bernd Reif. 2024. 'Aggregation Mechanisms and Molecular Structures of Amyloid- β in Alzheimer's Disease'. *Chemistry - A European Journal* 30 (August). <https://doi.org/10.1002/chem.202400277>.
- Palmqvist, Sebastian, Pontus Tideman, Niklas Mattsson-Carlgrén, Suzanne E. Schindler, Ruben Smith, Rik Ossenkoppele, Shorena Janelidze, Erik Stomrud, Oskar Hansson, et al. 2024. 'Blood Biomarkers to Detect Alzheimer Disease in Primary Care and Secondary Care'. *JAMA* 332 (15):