

Machine Learning

Dr. Mehran Safayani safayani@iut.ac.ir safayani.iut.ac.ir



https://www.aparat.com/mehran.safayani



https://github.com/safayani/machine_learning_course



Department of Electrical and computer engineering, Isfahan university of technology, Isfahan, Iran

Cost Function

$$h_{\theta}(x_i) = \theta_0 + \theta_1 x_i$$

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x_i) - y_i)^2$$

Minimize
$$J(heta_0$$
 , $heta_1$) $heta_0$, $heta_1$

If
$$J(\theta_1) = (\theta_1 - 2)^2$$

$$\frac{dJ(\theta_1)}{d\theta_1} = 0$$

$$\frac{dJ(\theta_1)}{d\theta_1} = 2 (\theta_1 - 2) = 0 \longrightarrow \theta_1 = 2$$

Minimize
$$J(heta_0$$
 , $heta_1$) $heta_0$, $heta_1$

Minimize
$$J(\theta_0$$
 , θ_1 , ... , θ_n) θ_0 , θ_1 , ... , θ_n

Repeat until convergence: {

For j=0,...,n

$$\theta_j = \theta_j - \alpha \frac{dJ(\theta_0, \theta_1, ..., \theta_n)}{d\theta_j}$$

 α is learning rate

Updating all $heta_{i}$ Simultaneously

Convergence condition:

$$\|\theta^{t+1} - \theta^t\|_2 \le \varepsilon$$

Correct form

temp0 =
$$\theta_0 - \alpha \frac{dJ(\theta_0, \theta_1)}{d\theta_0}$$

temp1 =
$$\theta_1 - \alpha \frac{dJ(\theta_0, \theta_1)}{d\theta_1}$$

 θ_0 = temp0

 θ_1 = temp1




Incorrect form

$$\theta_0 = \theta_0 - \alpha \frac{dJ(\theta_0, \theta_1)}{d\theta_0}$$

$$\theta_1 = \theta_1 - \alpha \frac{dJ(\theta_0, \theta_1)}{d\theta_1}$$






خطوط مماس نشان داده شده دارای شیب یا مشتق مثبت هستند. در نتیجه:

$$\frac{dJ(\theta^{1})}{d\theta^{1}} > 0, \ \alpha > 0 \implies \alpha \frac{dJ(\theta^{1})}{d\theta^{1}} > 0$$

$$\implies \theta^{2} = \theta^{1} - \alpha d\theta^{1}$$

 θ کوچکتر میشود و به سمت چپ حرکت میکنیم.




خطوط مماس نشان داده شده دارای شیب یا مشتق مثبت هستند. در نتیجه:

$$\frac{dJ(\theta^1)}{d\theta^1} < 0, \ \alpha > 0 \implies \alpha \frac{dJ(\theta^1)}{d\theta^1} < 0$$


$$\implies \theta^2 = \theta^1 - \alpha d\theta^1$$

بزرگتر میشود و به سمت راست حرکت میکنیم. θ

Choosing Learning Rate



Gradient Descent Weakness



Linear regression model

$$h_{\theta}(x_i) = \theta_0 + \theta_1 x_i$$

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^{m} (h_{\theta}(x_i) - y_i)^2$$

$$\frac{dJ(\theta_0, \theta_1)}{d\theta_0} = \frac{1}{m} \sum_{i=1}^{m} (h_{\theta}(x_i) - y_i)$$

$$\frac{dJ(\theta_{0}, \theta_{1})}{d\theta_{1}} = \frac{1}{m} \sum_{i=1}^{m} (h_{\theta}(x_{i}) - y_{i}) x_{i}$$

Linear regression model

Repeat until convergence:

$$\boldsymbol{\theta_0} = \boldsymbol{\theta_0} - \alpha \frac{1}{m} \sum_{i=1}^{m} (h_{\theta}(x_i) - y_i)$$

$$\boldsymbol{\theta_1} = \boldsymbol{\theta_1} - \alpha \frac{1}{m} \sum_{i=1}^{m} (h_{\theta}(x_i) - y_i) x_i$$

روز رسانی همزمان

$$\theta^t = \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix} \quad , \qquad \theta^{t+1} = \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix} \quad , \qquad d\theta = \begin{bmatrix} d\theta_0 \\ d\theta_1 \end{bmatrix}$$

Convergence condition:

$$\|\theta^{t+1} - \theta^t\|_2 = \sqrt[2]{(\theta_0^{t+1} - \theta_0^t)^2 + (\theta_1^{t+1} - \theta_1^t)^2} < \varepsilon$$

 $\|d\theta\|_2 < \varepsilon$

Batch Gradient Descent

$$\frac{dJ(\theta)}{d\theta} = \frac{1}{m} \sum_{i=1}^{m} (h_{\theta}(x_i) - y_i)$$

Batch Gradient Descent

```
\theta_0 \longleftarrow random, \theta_1 \longleftarrow random
Repeat until convergence:
J— 0, d\theta_1 — 0, d\theta_0 — 0
For i = 1 to m:
              h_{\theta}(x_i) = \theta_0 + \theta_1 x_i
              j += (h_{\theta}(x_i) - y_i)^2
              d\theta_1 += 2(h_\theta(x_i) - y_i) x_i
               d\theta_0 += 2 (h_\theta(x_i) - y_i)
J/=2m
d\theta_1 \neq 2m
d\theta_0 /= 2m
\theta_1 = \theta_1 - \alpha d\theta_1
\theta_0 = \theta_0 - \alpha d\theta_0
```