Calc in 3d Notes

$saffron_{\scriptscriptstyle{-}}$

Contents

Chapter 2: Vectors in Space	2
Appendix	4

Chapter 2: Vectors in Space

Graphing

TEMPLATE TEMPLATE

The Vector

ex image here lacktriangle a quantity with a magnitude and direction

• unit vector $\hat{u} = \frac{\vec{v}}{\|\vec{v}\|}$

Vector Operations

• Addition

$$- \vec{a} = \langle 1, 2 \rangle, \vec{b} = \langle 3, 4 \rangle$$
$$- \vec{a} + \vec{b} = \langle 4, 6 \rangle$$

• Scalar Multiplication

$$- \vec{v} = \langle 1, 3 \rangle, c = 2$$
$$- c\vec{v} = \langle 2, 6 \rangle$$

simple inverses for subtraction and scalar division exist.

• Dot Product (also 2d!)

$$\begin{aligned}
& \text{img} \\
& \vec{u} \cdot \vec{v} = \|\vec{u}\| \, \|\vec{v}\| \cos \theta
\end{aligned}$$

$$\vec{u} = \langle u_1, u_2, u_3 \rangle, \ \vec{v} = \langle v_1, v_2, v_3 \rangle$$

 $\vec{u} \cdot \vec{v} = \sum u_i v_i$

$$- \ \vec{v} \cdot \vec{v} = \|\vec{v}\|^2$$

– two vectors are orthogonal aka \perp iff their dot product is 0

$$-\cos\theta = \frac{\vec{u}\cdot\vec{v}}{\|u\|\,\|v\|} \text{ wtf is equation } 2.5$$
 "unique over this range" on abt

$$- \text{ work} = \vec{F} \cdot \vec{D}$$

- comp (scalar projection)

$$\begin{aligned} \operatorname{comp}_{\vec{v}} \vec{u} &= \|\vec{v}\| \cos \theta \\ &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\|} \end{aligned}$$

proj (vector projection)

$$\begin{aligned} \operatorname{proj}_{\vec{u}} \vec{v} &= \operatorname{comp}_{\vec{u}} \vec{v} \cdot \frac{\vec{u}}{\|\vec{v}\|} \\ &= \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\|^2} \vec{u} \end{aligned}$$

• Cross Product (3d)

$$\lim_{\|\vec{u} \times \vec{v}\| = \|\vec{u}\| \|\vec{v}\| \sin \theta$$

$$- \vec{u} \times \vec{v} = \det \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

- result is \perp to both input vectors, direction by right hand rule

- triple scalar product:

$$\vec{u} \cdot (\vec{v} \times \vec{w})$$

$$= (\vec{u} \times \vec{v}) \cdot \vec{w}$$

$$= \det \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}$$

- if \vec{u} and \vec{v} are the sides of a parallelogram, then its area is $\|\vec{u} \times \vec{v}\|$

- if a parallelepiped has edges $\vec{u}, \vec{v}, \vec{w}$, its volume is the absolute value of its triple scalar product

- torque =
$$\vec{\tau} = \vec{r} \times \vec{F}$$

Appendix

matrices, 3x3 determinants

Recall that a elements of a matrix are enumerated a_{ij} where i is column and j is row, both 1-indexed.

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

Recall that the minor of the matrix element here is the 2 by 2 determinant when you take away the row and column of the element in a 3 by 3 matrix. Picking an arbitrary row (or even column), with a_{ij} being an element and M_{ij} being a minor, a 3 by 3 determinant is calculated by

$$\sum_{j=1}^{3} (-1)^{i+j} a_{ij} M_{ij}$$

In the figure media/3x3determinant.png, $(-1)^{i+j}$ is in green, a_{ij} in orange, and M_{ij} in blue.

cross/dot products

- see:
 - media/cross_prod_area_ex.png
 - media/scalar_projection_composition_ex.png
 - media/scalar_triple_prod_parallelepiped_ex.png