

GRE Physics Study Notes v2

Courtesy of Nicole Duncan, transcribed by Jeren Suzuki

Last Edited 18th September 2013

Contents

1	Hermitian Matrix	1
2	Doppler Effect	1
2.1	Relativistic	1
3	Lagrangian and Hamiltonian	1
4	The Structure of Hydrogen	2
5	Particle in a Box	2
6	Free Particle	2
7	Schrödinger's Equation	3
8	Index of Refraction	3
8.1	Cherenkov Radiation	3
8.2	Bremsstrahlung Radiation	3
9	Gauss' Laws	3
10	Damped Driven Oscillator	3
10.1	Critically Damped $\rightarrow \omega = \beta$	3
10.2	Overdamped $\rightarrow \omega < \beta$	4
10.3	Underdamped $\rightarrow \omega > \beta$	4
11	Traveling Wave Formalism	4
12	Maxwell Velocity Distribution	4
13	Mean Free Path	4
14	Cross Section	4
15	Particle Diffusion (Fick's Law)	4
16	Thermal Diffusion (Fourier's Law)	4
17	Thermodynamic Identity	4
18	Heat Capacity	5
19	Water	5
20	Decay	5
21	Beats	5
22	First Law of Thermodynamics	5
22.1	Second Law of Thermodynamics	5
22.2	Third Law of Thermodynamics	5
23	Fundamental Assumption of Statistical Mechanics	5
24	Isothermal	6
25	Adiabatic Compression	6
26	Heat	6
27	Cyclotron	6
28	Fermi	6
29	Telescope	6
30	Multiplicity/States	7
31	Rocket Motion	7
32	Collisions	7
33	Springs/Single Harmonic Oscillator	7
34	Thin Films	7
35	Conductivity/Current Density	8
36	Resistance	8
37	Boltzmann Statistics	8
38	Density of State Distribution	8
39	Band Pass Filter	8
40	Resonant Frequency	8
41	Central Force Motion	9
42	Moments of Inertia	9
43	Blackbody Radiation	9
44	Heat Engine	9
45	Refrigerator	9
46	Space-Time Diagram	9
47	Low-Pass	9

48	High Pass	10
49	Special Relativity	10
49.1	Time Dilation	10
49.2	Length Contraction	10
49.3	Invariant	10
49.4	Energy	10
50	Finite Potential Well	11
51	Fundamental Particles	11
52	Single Slit Diffraction	11
53	Double Slit	11
54	Diffraction Grating	11
55	Bragg	11
56	Aperture Limited: Airy Disk: Diffraction Limit: Angular Resolution	11
57	Electrostatics	12
57.1	Limits	12
57.2	Motion Through a Capacitor, etc.	12
57.3	Dipole	12
57.4	Current Density	12
57.5	Drift Speed	12
57.6	Conductivity	12
58	Magnetic Field	12
58.1	Tesla	13
58.2	Force	13
58.3	Cyclotron	13
58.4	Cycloid	13
58.5	Examples	13
58.5.1	Solenoid	13
58.5.2	Ring	13
58.5.3	Sheet of Current	13
58.5.4	Toroid	13
58.5.5	Dipole	13
59	Inductance	14
60	Radiation	14
61	Maxwell's Equations	14
62	Boundary Conditions	14
63	E&M Fields	14
64	Relativistic E&M	15
65	Coordinate Systems	15
66	Positronium	15
67	Free Expansion	15
68	Entropy	15
69	P/N Junctions	15
70	Wave Velocity	15
71	Polarizers	16
72	Heisenberg	16
73	Compton Effect	16
74	Photoelectric Effect	16
75	Phonon	16
76	Superconductor	16
77	Mirrors	16
77.1	Concave	17
77.2	Convex	17
78	Reflection/Refraction	17
79	Bloch's Theorem	17
80	Stern-Gerlach	17
81	Mixing Gases	17

82	Equipartition Theorem	17
83	Degrees of Freedom	18
84	Simple Pendulum	18
85	Gravity	18
86	Drag Force	18
87	Selection Rules	18
88	Stationary States	18
89	Spectroscopic Notation	19
90	Matter Waves	19
91	Pipes/Tubes/Fixed and Open Ends	19
92	Solutions to Time-Independent Schrödinger's Equation	19
93	Physical Pendulum	19
94	Intrinsic Magnetic Moment	20
95	Equations of Motions	20
96	Moments of Inertia	20
97	Hermitian Matrix	20
98	Balancing Problem	20
99	Decay Rates	20
100	Interferometer	20
101	Springs	21
102	Speed of Sound	21
103	Commutator Identities	21
104	Motion in a Circle	21
105	Specific Heat in a Solid	21
106	Doppler Shift	21
107	Fission	22
108	Wire Resistance	22
109	Spin Matrices	22
110	For What v Will a Car Stay on a Hill?	22
111	Böhr Model	22
112	Fluids	22
113	Gauss' with non-uniform	23
114	Capacitors	23
115	Diffraction Limit	23
116	Normal Modes	23
117	Radiation in Atoms	23
118	Ionization Energy	24
119	Binding Energy	24
120	Hierarchy of Forces	24
121	Pair Production	24
122	Spectral Lines	24
123	Photon Interactions with Matter	24
124	Neutron	25
125	Deuteron	25
126	Protium	25

1 Hermitian Matrix

1. Square Matrix
2. $A = A^\dagger \rightarrow$ the matrix is equal to its conjugate transpose
3. Entries on the diagonal are real
4. Sum of 2 Hermitian matrices is Hermitian
5. Product of 2 Hermitian matrices is Hermitian only if they commute
6. Eigenvalues are orthogonal
7. The determinant is real

2 Doppler Effect

$$f = f_0 \left[\frac{v + v_s}{v + v_0} \right] \quad (1)$$

$$\Delta d \downarrow = \begin{cases} +v_s \\ -v_0 \end{cases}$$

towards e/o

$$\Delta d \uparrow = \begin{cases} -v_s \\ +v_0 \end{cases}$$

away from e/o

2.1 Relativistic

$$\frac{f_0}{f} = \frac{\lambda}{\lambda_0} = \sqrt{\frac{1 + \beta}{1 - \beta}}, \beta = \frac{v}{c} \quad (2)$$

3 Lagrangian and Hamiltonian

$$L = T - U \quad (3)$$

$$\frac{\partial L}{\partial x} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}} \right) = 0 \quad (4)$$

$$H = T + U \text{ iff } U \neq U(\dot{x}), U \neq U(t) \quad (5)$$

$$\rho = \frac{\partial L}{\partial \dot{q}} \quad \dot{q} = \frac{\partial H}{\partial \rho} \quad \dot{p} = \frac{\partial H}{\partial q} \quad (6)$$

4 The Structure of Hydrogen

1. Fine: Spin/orbit + relativistic correction
Breaks l degeneracy, preserves j
why $E_{2s} < E_{2p}$
2. Hyperfine: spin/spin coupling of e^- /nucleus
Responsible for 21 cm line

$$\mu_p = \frac{ge}{2m_p} \langle \bar{s}_p \rangle \quad \mu_e = -\frac{e}{m_e} \langle \bar{s}_e \rangle \quad (7)$$

$$E_{n'f} = \frac{\mu_0 g_p e^2}{3\pi m_p m_e a^3} \langle \bar{s}_p \cdot \bar{s}_e \rangle \quad (8)$$

3. Stark Effect: Atom in external E
Not spin dependent
 $H' = eE_z$ if $E = \hat{E}_z$
Hydrogen:

$$E'_1 = \langle H' \rangle = eE \int_0^\infty d^3r z |\Psi_{100}|^2 = 0 \quad (9)$$

4. Zeeman Effect: Atom in external B
Spin/orbital angular momentum(l) + B coupling

$$H'_z = (\bar{\mu}_e + \bar{\mu}_s) \cdot B_{\text{ext}} \quad (10)$$

Weak: $B_{\text{ext}} \ll B_{\text{int}}$
 $E' \propto m_j \rightarrow$ breaks into $2j + 1$ levels
 Strong: $B_{\text{ext}} \gg B_{\text{int}}$
 $E' = \mu_B B_{\text{ext}} (m_e + 2m_s)$

5 Particle in a Box

$$E_n = n^2 E_0 \quad (11)$$

$$E_0 = \frac{\hbar^2 k^2}{2m} \quad (12)$$

$$k = \frac{n\pi}{a} \quad (13)$$

$$\Psi = \sqrt{\frac{2}{a}} \sin(kx), p = \hbar k \quad (14)$$

$$\text{3D: } E = \frac{\hbar^2}{2m} [k_x^2 + k_y^2 + k_z^2] \quad (15)$$

6 Free Particle

$$\Psi = A e^{i(kx - \omega t)} \quad (16)$$

$$\Delta p \Delta x = \frac{\hbar}{2} \quad (17)$$

$$\Delta x \Delta k \sim 1 \quad (18)$$

Packet moves with group velocity... $v_g = \frac{\partial \omega}{\partial k}$

$$\Psi = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Phi(k) e^{ikx} dk \quad \Phi = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Psi(x) e^{-ikx} dx \quad (19)$$

7 Schrödinger's Equation

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + V \right) \Psi = i\hbar \frac{\partial \Psi}{\partial t} \quad (20)$$

Separable $\Psi(x, t) = \Psi(x)\phi(t)$

$$\phi = e^{-iE_n t/\hbar} \quad (21)$$

8 Index of Refraction

$$n = \frac{c}{v} = \sqrt{\frac{\epsilon \mu}{\epsilon_0 \mu_0}} \quad v = v_\phi = \sqrt{\frac{1}{\epsilon \mu}} \quad (22)$$

$\lambda = \frac{\lambda_0}{n}$ inside a medium

8.1 Cherenkov Radiation

A charged particle passing through a medium which travels faster than the speed of light in that medium will emit light

$$n = \frac{c}{v} \longrightarrow v_{\min} = \frac{c}{n} \quad (23)$$

8.2 Bremsstrahlung Radiation

Continuous spectrum of radiation emitted when a charged particle is decelerated in a metal target

9 Gauss' Laws

$$\int E \cdot da = \frac{Q_{\text{in}}}{\epsilon_0} \rightarrow \nabla \cdot E = \frac{\rho_{\text{in}}}{\epsilon_0} \quad (24)$$

$$\int B \cdot da = 0 \rightarrow \nabla \cdot B = 0 \quad (25)$$

$$\int g \cdot da = 4\pi MG \rightarrow \nabla \cdot g = -4\pi G \rho_m \quad (26)$$

10 Damped Driven Oscillator

$$F = -\underbrace{kx}_{\text{spring}} - \underbrace{b\dot{x}}_{\text{damp}} + \underbrace{A \cos \theta}_{\text{driving}}, \omega = \sqrt{\frac{k}{m}}, \beta = \frac{b}{2m} \quad (27)$$

10.1 Critically Damped $\rightarrow \omega = \beta$

$$X_e = Ae^{-\omega_0 t} + A_2 t e^{-\omega_0 t} \quad (28)$$

10.2 Overdamped $\rightarrow \omega < \beta$

$$X_o = Ae^{-\beta t} e^{-\omega'' t}, \omega'' = \sqrt{\beta^2 - \omega_0^2} \quad (29)$$

10.3 Underdamped $\rightarrow \omega > \beta$

$$X_u = Ae^{-\beta t} \cos(\omega' t + \phi), \omega' = \sqrt{\omega_0^2 - \beta^2} \quad (30)$$

11 Traveling Wave Formalism

$$v_\phi = \frac{\omega}{k} \quad \psi = A \cos k(vt - x) = A \cos(\omega t - kx) \quad (31)$$

In one period, $x - vt = 2\pi$

12 Maxwell Velocity Distribution

$$D(v) \propto v^2 e^{-E/kT} \quad (32)$$

13 Mean Free Path

$$l = \frac{1}{n\sigma} \quad (33)$$

14 Cross Section

$$N_s = N_i \frac{N_t}{A} \sigma \quad (34)$$

15 Particle Diffusion (Fick's Law)

$$J = -D \nabla n \quad (35)$$

16 Thermal Diffusion (Fourier's Law)

$$\phi = -\sigma \nabla T \quad (36)$$

Interaction	Quantity	Variable	Formula
Mech	vol	P	$P = -\frac{\partial U}{\partial V} = T \left(\frac{\partial S}{\partial V} \right)$
Thermal	temp/energy	T	$T = \frac{\partial S}{\partial S}$
Diffuse	particles	μ	$\mu = -\frac{\partial U}{\partial N} = T \frac{\partial S}{\partial N}$

17 Thermodynamic Identity

$$TdS - PdV + \mu dN = dU \quad (37)$$

18 Heat Capacity

$$C \equiv \frac{dQ}{dT} \quad (38)$$

$$C_p = \frac{dQ}{dT} = T \frac{dS}{dT} \quad C_v = \frac{dQ}{dT} = \frac{dU}{dT} \quad (39)$$

At constant P , E lost to work $\Rightarrow T_p < T_V \Rightarrow C_p > C_v$

19 Water

$\rho = 1 \text{ g/cm}^3$, $1 \text{ L} = 1 \text{ kg}$

20 Decay

$$\begin{array}{ccc} {}^0_{-1}\beta + \bar{\nu} & {}^0_1\beta + \nu & {}^2_1D \\ {}^4_2\alpha & {}^0_0\gamma & {}^A_ZX \end{array}$$

21 Beats

$$f_0 = f_1 - f_2, \quad T_b = \frac{1}{f_1 - f_2}, \quad \text{occur when } f_1 + f_2 \text{ are close} \quad (40)$$

The tuned frequency:

$$f = \underbrace{n}_{\text{harmonic}} \underbrace{f_f}_{\text{fundamental}} \quad (41)$$

22 First Law of Thermodynamics

$$U = Q + W \quad (42)$$

22.1 Second Law of Thermodynamics

S increases or stays the same for any cyclic process.

22.2 Third Law of Thermodynamics

$$S(T=0) = 1 \quad C_v \rightarrow 0 \text{ as } T \rightarrow 0 \quad (43)$$

23 Fundamental Assumption of Statistical Mechanics

All accessible microscopic states are equally likely

24 Isothermal

Slow so T can equalize.

$$P_1 V_1 = P_2 V_2 \quad W = Nk \ln \left(\frac{V_i}{V_f} \right) \quad , W = - \int_{V_i}^{V_f} P dV \quad (44)$$

$$U = 0 \quad \text{since } \Delta T = 0 \quad , U = \frac{f}{2} Nk \Delta T \quad (45)$$

25 Adiabatic Compression

Fast, so no ΔQ lost, like opening a soda can.

$$\Delta Q = 0 \rightarrow U = W \quad (46)$$

$$\Delta U = Nk \Delta T = W \quad (47)$$

$$\gamma = \frac{f+2}{f} \quad , W = \frac{P_f V_f - P_i V_i}{1-\gamma} \quad , V_f^\gamma P_f = V_i^\gamma P_i \quad (48)$$

26 Heat

$$Q = T dS \quad (49)$$

$$Q = mc \Delta T \quad (50)$$

$$Q = Pt \quad (51)$$

$$U = Q + W \quad (52)$$

27 Cyclotron

$$\omega = \frac{qB}{m} \quad F_c = F_B \rightarrow \frac{mv^2}{r} = qvB \rightarrow v = \frac{qBr}{m} = r\omega \quad (53)$$

28 Fermi

$$T_f = \frac{E}{k_B} \quad E_f = \frac{\hbar^2 k^2}{2m} \quad p = \hbar k \quad v_f = \frac{p_f}{m} \quad k_F = \left(\frac{3\pi^2 N}{\text{vol}} \right)^{1/3} \quad (54)$$

$$p_f = \frac{2}{3} \frac{E_f}{v} \quad (55)$$

Degenerate Fermi gas: so cold that all states below E_F are occupied

29 Telescope

$$M = - \frac{f_{\text{object}}}{f_{\text{eye}}} = \frac{\theta_{\text{eye}}}{\theta_{\text{object}}} \quad (56)$$

30 Multiplicity/States

Probability (Ω_n) = $\frac{\Omega_n}{\Omega_{all}}$ where Ω is the multiplicity # of things.

1. Total # microstates: (# of states can be in)^{# of things}
2. Ways to choose n things from N

$$\Omega \binom{N}{n} = \frac{N!}{(N-n)!n!} \quad (57)$$

31 Rocket Motion

$$u \frac{dm}{dt} + M \frac{dv}{dt} = 0 \quad (58)$$

$$v_f = v_0 + u \ln \left(\frac{M_i}{M_f} \right) \quad (59)$$

32 Collisions

1. Momentum is always conserved $p_i = p_f$
Don't forget to use (+) and (-) for before and after velocity collisions
2. KE and U conserved before or after collision only
- 3.

$$\epsilon = \frac{\overbrace{|v_1| + |v_2|}^{\text{after}}}{\underbrace{|U_1| + |U_2|}_{\text{before}}} \quad (60)$$

4. Impulse $J = F\Delta t = \Delta p = \Delta L$
5. Cross section $N_s = N_I \frac{N_t}{A} \sigma$

33 Springs/Single Harmonic Oscillator

$$F = -kx \quad U = \frac{1}{2}kx^2 \quad \omega = \sqrt{\frac{k}{m}} \quad (61)$$

$$ma = -kx = m\ddot{x} \quad (62)$$

$$E_{\text{tot}} = \frac{1}{2}kA^2, \quad A = \text{max amplitude} \quad (63)$$

34 Thin Films

$$\Delta\phi = \begin{cases} 0 & n_2 < n_1 \\ \pi & n_2 > n_1 \end{cases} \quad 2d = \begin{cases} n\lambda/2 & \Delta\phi_{\text{tot}} = \pi \\ n\lambda & \Delta\phi_{\text{tot}} = 0, 2\pi \end{cases}, \quad n = \text{odd \#s only}$$

35 Conductivity/Current Density

$$\bar{J} = ne\bar{v} = \sigma E \quad (64)$$

$$\sigma = \frac{ne^2\tau}{m} \quad (65)$$

36 Resistance

$$R = \frac{\rho L}{A} \quad (66)$$

37 Boltzmann Statistics

$$Z = \sum_i g_i e^{-E_i/kT}, g_i = \text{degeneracy of state } i \quad (67)$$

$$p_s = \frac{g_s e^{-E_s/k_B T}}{Z} \quad \frac{p_A}{p_B} = \frac{g_A}{g_B} \frac{e^{-A/kT}}{e^{-B/kT}} = \frac{g_A}{g_B} e^{(-A+B)/kT} \quad (68)$$

$$\langle \bar{X} \rangle = \frac{\sum_i e^{-E_i/kT}}{Z} \rightarrow \langle \bar{E} \rangle = \frac{1}{Z} \sum_i E_i e^{-E_i/kT} \quad (69)$$

$$U = N\bar{E} \rightarrow \text{total Energy of system} \quad (70)$$

38 Density of State Distribution

Fermions:

$$N_i = \frac{g_i}{e^{(E_i - \mu)/kT} + 1} \quad (71)$$

Bosons:

$$N_i = \frac{g_i}{e^{(E_i - \mu)/kT} - 1} \quad (72)$$

39 Band Pass Filter

$$j\omega C + \frac{1}{j\omega L} = Z = \frac{-\omega^2 CL + 1}{j\omega L} \quad \omega_o = \frac{1}{\sqrt{LC}} \quad (73)$$

40 Resonant Frequency

Inductor and capacitor in series:

$$j\omega L = \frac{1}{j\omega C} \rightarrow \omega_o^2 LC = 1 \rightarrow \omega = \frac{1}{\sqrt{LC}} \quad (74)$$

Inductor and capacitor in parallel:

$$j\omega C + \frac{1}{j\omega L} = 0 \rightarrow j\omega C = \frac{1}{j\omega L} \rightarrow \omega = \frac{1}{\sqrt{LC}} \quad (75)$$

41 Central Force Motion

$$\mu = \frac{m_1 m_2}{m_1 + m_2} \quad R_{\text{CM}} = \frac{\sum m_i r_i}{\sum m_i} \quad T = \frac{1}{2} \mu |\dot{r}|^2$$

$$r_1 = \frac{m_2}{m_1 + m_2} r \quad r_2 = \frac{m_1}{m_1 + m_2} r \quad \bar{r} = \bar{r}_1 - \bar{r}_2$$

42 Moments of Inertia

1. Hoop: MR^2
2. Disk: $\frac{1}{2}MR^2$
3. Solid Sphere: $\frac{2}{3}MR^2$
4. Hollow Sphere: $\frac{2}{5}MR^2$
5. Rod End: $\frac{1}{3}ML^2$
6. Rod Middle: $\frac{1}{12}ML^2$

43 Blackbody Radiation

$$T\lambda = 3 \text{ mmK} \quad P \propto T^4 \quad \rho \propto AT^4 \text{ (for photons)} \quad (76)$$

44 Heat Engine

$$e \leq 1 - \frac{T_c}{T_h} \quad e = \frac{\text{benefit}}{\text{cost}} = \frac{W}{Q_h} \quad W = Q_h - Q_c \quad (77)$$

45 Refrigerator

$$e \leq \frac{T_c}{T_h - T_c} \quad e = \frac{\text{benefit}}{\text{cost}} = \frac{Q_c}{W} \quad (78)$$

46 Space-Time Diagram

$$s^2 = x^2 - (ct)^2 \quad (79)$$

1. $s > 0$ Spacelike
 Δt can equal 0, simultaneous events occur
2. $s < 0$ Timelike
Events can occur at same point in space, $\Delta x = 0$, but not simultaneously $\Delta t \neq 0$
3. $\Delta s = 0$ Lightlike

47 Low-Pass

RC or LR perpendicular to each other. For RC:

$$\frac{1}{R + \frac{1}{j\omega C}} = \frac{1}{j\omega CR + 1} \quad , \omega \rightarrow 0, \rightarrow 1 \quad (80)$$

For LR:

$$\frac{R}{R + j\omega L}, \omega \rightarrow 0, \rightarrow 1 \quad (81)$$

Square signals turn into wave-like signals with crests.

$$V_{\text{out}} = V_{\text{in}} \left(\frac{Z_2}{Z_1 + Z_2} \right) \quad (82)$$

Where Z_1 and Z_2 are on either side of a perpendicular V_{out} with a V_{in} leading into Z_1 .

48 High Pass

CR or RL, turns square signals into signals where the flat tops of square turns into decaying to 0.

49 Special Relativity

v/c	γ	
.1	1.005	Motion in \hat{x} :
.25	1.033	
.5	1.151	
.75	1.55	
.9	2.3	

$$x = \gamma(x' + vt') \quad (83)$$

$$t = \gamma \left(t' + \frac{vx}{c^2} \right) \quad (84)$$

$$u'_x = \frac{u_x + v}{1 + \frac{u_x v}{c^2}} \quad (85)$$

$$u'_z = \frac{u_z}{\gamma \left(1 + \frac{u_x v}{c^2} \right)} \quad (86)$$

49.1 Time Dilation

$$t' = \gamma t_0 \quad (87)$$

49.2 Length Contraction

$$x' = \frac{x}{\gamma} \quad (88)$$

49.3 Invariant

$$\Delta s^2 = \Delta x^2 - (ct)^2 \quad (89)$$

49.4 Energy

$$E_r^2 = E_0^2 + (pc)^2 \quad E_r \neq \frac{p_r^2}{2m} \quad E_r = \gamma E_0 \quad p_r = \gamma mv = \gamma p \quad (90)$$

50 Finite Potential Well

$$E \propto n^2 \quad d \propto \frac{1}{\sqrt{V - E_n}} \rightarrow d \propto n \quad (91)$$

51 Fundamental Particles

1. Bosons:
Gauge Boson - Gluon - Strong
Photon - E&M
W,Z Bosons - a.k.a weak bosons
Higgs
Graviton
Pion
2. Fermions:
Quarks (Up, down, top, bottom, strange, charm)
Leptons (Electron, Muon, Tauon) and neutrino variants of each
3. Composite Fermions: Protons and Neutrons, etc.

52 Single Slit Diffraction

$$d \sin \theta = n\lambda \quad \theta = \text{angle between central max and first minimum} \quad (92)$$

$$\tan \theta = \frac{y}{L} \quad \text{central max, width } \Delta y_{\max} = \frac{2L\lambda}{d} \quad (93)$$

53 Double Slit

$$d \sin \theta = n\lambda \quad \Delta y = L \tan \theta \quad (94)$$

54 Diffraction Grating

$$d \sin \theta = n\lambda \quad y = L \tan \theta = L \frac{\sin \theta}{\cos \theta} = \frac{Ln\lambda}{d \cos \theta} \quad (95)$$

55 Bragg

$$2d \sin \theta = n\lambda \quad d = \frac{a}{\sqrt{l^2 + h^2 + k^2}}, \quad \text{letters are miller indices} \quad (96)$$

56 Aperture Limited: Airy Disk: Diffraction Limit: Angular Resolution

$$\sin \theta = \frac{1.22\lambda}{D}, \quad D = \text{diameter of aperture}, \quad \theta = \text{angular separation} \quad (97)$$

57 Electrostatics

$$F = \frac{kq_1q_2}{r^2} \quad \epsilon = k\epsilon_0 \quad k = 9 \times 10^9 \frac{\text{Nm}}{\text{C}^2} \quad (98)$$

1. Sphere: $\propto \frac{1}{r^2}$
2. Infinite Line: $\propto \frac{1}{r}$
3. Infinite Plane doesn't fall off: $E = \frac{\sigma}{2\epsilon_0}$
4. Ring: $\propto \frac{x}{d^3}$, $d = \sqrt{x^2 + R^2}$
5. Capacitor doesn't fall off: $E_{\text{out}} = 0$, $E_{\text{in}} = \frac{\sigma}{\epsilon_0}$

57.1 Limits

As $x \rightarrow \infty$ all objects look like point objects. Sometimes use binomial approximation to get behavior at ∞ , $(1+x)^n \sim 1 + nx$ for $x \ll 1$.

57.2 Motion Through a Capacitor, etc.

Use kinematics equations $F = ma = qE$ find V, a, t to get θ deflection.

57.3 Dipole

$$\bar{p} = q\bar{d} \quad (99)$$
$$\bar{E}_{\text{dipole}} = \begin{cases} \frac{2k\bar{p}}{r^3} & \text{axis of } \bar{d} \\ -\frac{k\bar{p}}{r^3} & \perp \text{ to } \bar{d} \end{cases}$$

57.4 Current Density

$$J = nev_d, I = JA = \frac{\text{current of cross section}}{m^2} \quad (100)$$

57.5 Drift Speed

$$J = \sigma E = \frac{ne^2\tau}{m} E \rightarrow v_d = \frac{\sigma E}{ne} = \frac{e\tau E}{m} \quad (101)$$

57.6 Conductivity

$$\sigma = \frac{ne^2\tau}{m} \quad (102)$$

58 Magnetic Field

$$B = \frac{\mu_0 I}{4\pi} \frac{d\bar{l} \times \hat{r}}{r^2}, d\bar{l} = \text{length and direction} \quad (103)$$

58.1 Tesla

$$T = \frac{N}{A \cdot m}, \text{ current } I = \int J \cdot da_{\perp} \quad (104)$$

58.2 Force

$$F = q\bar{v} \times \bar{B} = I(d\bar{l} \times \bar{B}) \quad (105)$$

58.3 Cyclotron

$$E \parallel B \rightarrow \text{Helical motion} \quad (106)$$

$$v_{\parallel} B \rightarrow \text{Helical} \quad (107)$$

$$\frac{mv^2}{r} = qvB \rightarrow F_c = F_m \quad (108)$$

58.4 Cycloid

$$E \perp B \quad (109)$$

58.5 Examples

58.5.1 Solenoid

$$B = \begin{cases} 0 & \text{outside} \\ \frac{\mu_0 IN}{L} & \text{inside} \end{cases}$$

58.5.2 Ring

$$B = \frac{\mu_0 I}{2R} \quad (110)$$

Any displacement along center of ring should reduce tho this equation as $x \rightarrow 0$.

58.5.3 Sheet of Current

$$B = \begin{cases} -\frac{\mu_0}{2} & z > 0 \\ \frac{\mu_0}{2} & z < 0 \end{cases}$$

58.5.4 Toroid

$$B = \begin{cases} 0 & \text{out} \\ \frac{\mu_0 IN}{2\pi R} & \text{in} \end{cases}$$

58.5.5 Dipole

$$B \propto \frac{\mu}{r^3}, \mu = IA = \text{dipole moment} \quad (111)$$

$$B \propto \frac{IA}{r^3}, \text{ as } x \rightarrow \infty, \text{ this is twice the field of a current loop} \quad (112)$$

59 Inductance

$$\Phi = LI \quad \epsilon = -\frac{d\Phi}{dt} \quad \Phi_B = \int B \cdot dA \quad (113)$$

$$W = \frac{1}{2}LI^2 \quad \left(\text{corollary: cap } W = \frac{t}{2}CV^2 \right) \quad (114)$$

$$I(t) = \frac{\epsilon_0}{R} \left[1 - e^{-(R/L)t} \right] \quad \tau = \frac{L}{R} \quad (115)$$

60 Radiation

Electric Dipole	Magnetic Dipole	Point Charge
$P \propto q^2 d^2 \omega^4$	$P \propto I^2 \omega^4$	$P \propto q^2 a^2$

$$P_{\max} \perp \text{ to } \hat{a}.$$

61 Maxwell's Equations

$$\begin{aligned} \nabla \cdot E &= \frac{\rho_{\text{in}}}{\epsilon_0} & \nabla \times E &= -\frac{\partial B}{\partial t} \\ \nabla \cdot B &= 0 & \nabla \times B &= \mu_0 J - \mu_0 \epsilon_0 \frac{\partial E}{\partial t} \\ \oint E \cdot dA &= \frac{Q_{\text{in}}}{\epsilon_0} & \oint E \cdot dl &= -\frac{\partial \Phi_B}{\partial t} \\ \oint B \cdot dA &= 0 & \oint B \cdot dl &= \mu_0 I + \mu_0 \epsilon_0 \frac{\partial \Phi_E}{\partial t} \end{aligned}$$

62 Boundary Conditions

$$E_{\parallel} = 0, \quad B_{\perp} = 0 \quad \text{for reflected waves} \quad E_{\text{tot}} = 0 \quad B_{\text{tot}} = 2B_{\text{wave}} \quad (116)$$

$$\epsilon_1 E_1^{\perp} - \epsilon_2 E_2^{\perp} = \sigma_F \quad E_1^{\parallel} = E_2^{\parallel} \quad (117)$$

$$B_1^{\perp} = B_2^{\perp} \quad \frac{B_1^{\parallel}}{\mu_1} - \frac{B_2^{\parallel}}{\mu_2} = k_f \times \hat{n} \quad (118)$$

63 E&M Fields

B/E are in phase and perpendicular

$$B_{+0} = \frac{k}{\omega} E_0 = \frac{1}{c} E_0 \quad (119)$$

$$\text{Energy Density } \langle U \rangle = \frac{\epsilon_0}{2} E^2 \quad \text{Intensity } \langle I \rangle = \frac{1}{2} c \epsilon E^2 \quad (120)$$

$$\text{Radiation Pressure } p = \frac{\langle s \rangle}{c} \quad \bar{s} = \frac{\bar{E} \times \bar{B}}{\mu_0}, \quad \hat{s} = \text{propagation of } \frac{E}{\mu} \quad (121)$$

64 Relativistic E&M

Processes change between frames, but outcome is same.

Example: Parallel Plate Capacitor:

$$\sigma_0 = \frac{Q}{A} \quad \sigma' = \frac{Q}{A'} \quad , \quad \text{length contracts, } A' < A \rightarrow \sigma' > \sigma_0 \quad \sigma' = \gamma\sigma_0 \quad (122)$$

$$\text{Also, } A = lw \quad a' = \frac{l}{\gamma}w = \frac{lw}{\gamma} \quad (123)$$

$$\text{so, } E_{\perp} = \gamma E_0 \quad E_{\parallel} = E_{\parallel} \quad (124)$$

65 Coordinate Systems

1. Cartesian: $dl = \hat{x}dx + \hat{y}dy + \hat{z}dz$, $dV = dxdydz$
2. Spherical: $dl = \hat{r}dr + r d\theta\hat{\theta} + r \sin\theta d\phi\hat{\phi}$, $dV = r^2 \sin\theta dr d\phi d\theta$
3. Cylindrical: $dl = \hat{s}ds + s d\phi\hat{\phi} + \hat{z}dz$, $dV = s ds d\phi dz$

66 Positronium

$$\mu = \frac{m_e}{2} \rightarrow E_p = \frac{E_H}{2} \quad E_{pos} = \frac{-13.6 \text{ eV}}{2} \quad E_p = \frac{-6.8}{n^2} \quad (125)$$

67 Free Expansion

$$W = 0 \quad , \quad Q = 0 \quad , \quad \Delta S > 0 \quad , \quad \Delta S = Nk \ln \left(\frac{V_i}{V_f} \right) \quad (126)$$

68 Entropy

$$S = k \ln(\Omega) \quad , \quad Q = TdS \quad S_{\text{tot}} = S_A + S_B \quad (127)$$

69 P/N Junctions

1. n donate e^- to CB
2. p donate holes to VB

70 Wave Velocity

$$\text{Group Velocity } v_g = \frac{\partial \omega}{\partial k} \quad \text{phase } v_{\phi} = \frac{\omega}{k} = \sqrt{\frac{1}{\epsilon\mu}} = \frac{\lambda}{T} \quad (128)$$

71 Polarizers

1. $I = I_0 \cos^2 \theta$ for plane polarized
2. $I = \frac{I_0}{2}$ for natural light

72 Heisenberg

$$\sigma_p \sigma_x \geq \frac{\hbar}{2} \quad \sigma_A \sigma_B \geq \frac{1}{2i} \langle [\hat{A}, \hat{B}] \rangle \quad (129)$$

73 Compton Effect

Elastic scattering of photons - shows particle nature of light

$$\Delta\lambda = \lambda_c(1 - \cos \theta) \quad \lambda_c = \frac{hc}{E_0} = \frac{h}{mc} \quad (130)$$

74 Photoelectric Effect

$$KE_{\max} = E_p - \Phi = \hbar\omega - \Phi = \frac{hc}{\lambda} - \Phi = hf - \Phi \quad (131)$$

$$\text{Energy of photon} = \frac{hc}{\lambda} = hf = \frac{h}{2\pi} \omega = \hbar\omega \quad (132)$$

Einstein's Equation: $eV = hf - \Phi$, $V = (-)$ value which e^- can be stopped from hitting the cathode, $I \rightarrow 0$ (133)

75 Phonon

Displacement from equilibrium values of plane spacing

$$E_{\text{phonon}} = \hbar\omega \quad (134)$$

76 Superconductor

1. Meissner: $B = 0$ inside S_c
2. $\rho \rightarrow 0$ at critical temp
3. λ_c penetration depth measures how far B penetrates before $\rightarrow 0$
 $B = B_0 e^{-x/\lambda_c}$

77 Mirrors

$$\frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f} \rightarrow d_i = \frac{d_o f}{f - d_o} \quad (135)$$

$$M = \frac{h_i}{h_o} = -\frac{d_i}{d_o} \quad f = \frac{R}{2} \quad (136)$$

Rays go through center and continues in a straight line and goes parallel then through focal point.

77.1 Concave

Converging Mirror, what most diagrams are of

77.2 Convex

Diverging Mirror. Images always smaller, virtual, and upright. They cover a wide field of view.

78 Reflection/Refraction

Snell's Law:

$$n_1 \sin \theta_1 = n_2 \sin \theta_2 \quad (137)$$

$$\theta_i = \theta_r, \theta \text{ measured relative to normal} \quad (138)$$

For total internal reflection:

$$n_1 \sin \theta_1 = n_2 \sin 90 \quad (139)$$

$$\sin \theta_1 = \frac{n_2}{n_1} \quad (140)$$

$$n = \frac{c}{v} = \sqrt{\frac{\epsilon\mu}{\epsilon_0\mu_0}} \quad \epsilon = k\epsilon_0 \quad \frac{n_1}{n_2} = \frac{v_2}{v_1} \quad (141)$$

79 Bloch's Theorem

Ψ solutions to Schrödinger's Equation are plane waves modulated by a function with the periodicity of the lattice

80 Stern-Gerlach

Expected $2s + 1$ states, saw $2s + 1 = 2$ states. Implied that $H = -\gamma \vec{S} \cdot \vec{B}$

81 Mixing Gases

$$\begin{cases} \text{if } A \neq B & \Delta s = \Delta s_A + \Delta s_B \\ \text{if } A = B & \Delta s = 0 \text{ since } \Delta \ln(\Omega) \sim 0 \text{ since } \Omega = \text{large } \# \end{cases}$$

82 Equipartition Theorem

$$U = \frac{f}{2} N k T \quad (142)$$

$f = \#$ quadratic terms in Hamiltonian (not degrees of freedom, but in general they're equal).

$$kT \sim \frac{1}{40} \text{ eV @ room temperature} \quad (143)$$

83 Degrees of Freedom

1. A quadratic term in PE or KE
2. Translational ($\frac{mv^2}{2}$)
3. Rotational ($I\omega^2$)
4. Vibrational (Counts as 2) (kx^2, mv^2)

84 Simple Pendulum

$$\omega = \sqrt{\frac{g}{l}} \quad \theta = \theta_{\max} \sin(\omega t) \quad (144)$$

85 Gravity

$$F_g = \frac{Gm_1m_2}{r^2} = mg \quad \text{Kepler: } T^2 \propto a^3 \quad (145)$$

Escape velocity:

$$F_c = F_g \rightarrow mg = \frac{mv^2}{r} \rightarrow v = \sqrt{gr} \quad (146)$$

Gauss:

$$\int g \cdot dA = -4\pi G \int dM_{\text{in}} \rightarrow \int g \cdot dA = -4\pi GM_{\text{in}} \quad (147)$$

86 Drag Force

$$F_D \propto v^n \quad \text{air: } F_D \propto v^2 \quad (148)$$

$$\text{Terminal velocity } F_D = F_g \rightarrow v = \sqrt{\frac{g}{k}} \quad (149)$$

$$F_D = mkv^2 \quad (150)$$

87 Selection Rules

$$\Delta l = \pm 1 \quad \Delta m = \pm 1, 0 \quad \Delta s \text{ no rule} \quad \Delta n \geq 1 \quad (151)$$

88 Stationary States

$$\langle \Psi | \Psi \rangle \text{ is not a function of time} \quad (152)$$

89 Spectroscopic Notation

$$^{2s+1}L_j \quad (153)$$

1. Spin isn't always $\frac{1}{2}$
2. L = orbital angular momentum
3. $j = s + L$ = total angular momentum

$$L = \begin{cases} s & 0 \\ p & 1 \\ d & 2 \\ f & 3 \end{cases}$$

90 Matter Waves

$$p = \hbar k \quad k = \frac{2\pi}{\lambda} \quad p = \frac{h}{\lambda} \rightarrow \lambda = \frac{h}{p} \quad (154)$$

91 Pipes/Tubes/Fixed and Open Ends

1. Closed on both ends:
Ends are nodes, $\sin kx$ dependence, $\Psi = A \sin kx \cos \omega t$
 $\lambda = \frac{2L}{n}$ longest λ is $2L$
2. Open on both ends:
Ends are antinodes, $\cos kx$ dependence
 $\lambda = \frac{2L}{n}$ $\Psi = A \cos kx \cos \omega t$
3. Closed/Open ends:
Node/Antinode Longest $\lambda = 4L$ (so $\frac{1}{4}\lambda$ can fit end to end)
 $\lambda = \frac{4L}{2n-1}$ (-1 if n starts at 1)
 $\Psi = A \sin kx \cos \omega t$

92 Solutions to Time-Independent Schrödinger's Equation

$$\langle H \rangle = \sum_n |C_n|^2 E_n = C_1 E_1 + C_2 E_2 + \dots, \quad \sum_n |C_n|^2 = 1 \quad (155)$$

$$\text{Prob}(a < x < b) = \int_a^b |\Psi(x)|^2 dx \rightarrow \text{area under the } |\Psi|^2 \text{ vs } x \text{ graph} \quad (156)$$

93 Physical Pendulum

$$\tau = I\alpha = I\dot{\omega} = mg \sin \theta L_{\text{cm}} = \bar{r} \times \bar{F} \quad (157)$$

$$\ddot{\theta} = \frac{mgL_{\text{CM}}}{I}\theta \rightarrow \omega = \sqrt{\frac{mgL_{\text{CM}}}{I}}, \quad L_{\text{CM}} = \text{the distance from the pivot point to the center of mass} \quad (158)$$

94 Intrinsic Magnetic Moment

$$\bar{\mu}_s = \frac{gq}{2m} \bar{s} \quad m \text{ is dominant factor} \quad (159)$$

95 Equations of Motions

Look for boundary values $I(s)$ given $x(t)$ and $y(t)$. Differentiate and see which one yields v_0

96 Moments of Inertia

The moment of an object stretched along the axis of rotation doesn't change

$$I_{\text{disk}} = I_{\text{cylinder}} \quad (160)$$

The moment of a cuboid:

$$I = \frac{M}{12}(x^2 + y^2) \quad (161)$$

“Twin Plate”, $y = 0$ so $I_z = \frac{M}{12}x^2$. Then, $I_z = \frac{1}{12}(2d)^2 = \frac{M}{3}d^2$

97 Hermitian Matrix

Real eigenvalues, square

$$A = A^T = A^{*T} \text{ the entries are equal to their conjugate transpose} \quad (162)$$

All diagonals must be real. The sum of two Hermitian matrices is also Hermitian.

$$\langle f | \hat{A} f \rangle = \langle \hat{A} f | f \rangle \Rightarrow A = A^* \quad (163)$$

98 Balancing Problem

Easiest to use center of mass.

$$\text{Center of Mass} = \frac{\sum m_i r_i}{\sum m_i} \text{ one mass is at } -r \quad (164)$$

99 Decay Rates

$$\frac{dA}{dt} = -kA \rightarrow A = A_0 e^{-kt}; \frac{A}{A_0} = \frac{1}{2} = e^{-kt} \rightarrow t_{1/2} = \frac{\ln(2)}{k} \quad (165)$$

100 Interferometer

Fringe shifts occur for changing distance or λ .

$$2d = m\lambda \quad d = \text{change in distance} \quad \lambda = \Delta\lambda \quad \lambda_{\text{gas}} = \frac{\lambda_{\text{vac}}}{n} \quad (166)$$

In a tube where gas \rightarrow vacuum:

$$2d = m(\lambda_{\text{gas}} - \lambda_{\text{vac}}) = m\lambda_{\text{vac}} \left(\frac{1}{n} - 1 \right) \quad (167)$$

101 Springs

Add like capacitors. Makes sense because in series they can stretch more so $F = kx$ must be decreased, in parallel, they stretch less so $k \uparrow$ for the same force.

$$\text{Springs in series: } \frac{1}{k_{\text{tot}}} = \frac{1}{k_1} + \frac{1}{k_2} \quad (168)$$

$$\text{Springs in parallel: } k = k_1 + k_2 \quad (169)$$

102 Speed of Sound

In an ideal gas: $v \propto \sqrt{T}$

103 Commutator Identities

$$[A, B] = -[B, A] \quad (170)$$

$$[AB, C] = A[B, C] + [A, C]B \quad (171)$$

$$[A, BC] = B[A, C] + [A, B]C \quad (172)$$

104 Motion in a Circle

Always a_{radial} component. Only a_{tan} if v_{tan} changes.

$$F = \frac{mv^2}{r} = ma_r \rightarrow a_r = \frac{v^2}{r} \quad (173)$$

$$a_r = r \times \alpha \quad v = r \times \omega \quad (174)$$

105 Specific Heat in a Solid

1. Einstein Model:
Treats atoms as harmonic oscillators, $3N$ total harmonic oscillators and they all have the same energy (frequency) using Bose-Einstein statistics
2. Debye:
Also $3N$ harmonic oscillators. Assigns a range of energies (frequencies) and treats the lattice vibrations as phonons in a box
3. Dulong-Petit:
High temperature, uses equipartition theorem with harmonic oscillators ($f = 6, c = 3Nk$). Debye and Einstein models reduce to this in the high T limit.

106 Doppler Shift

$$f = f_0 \left(\frac{1 + v_s}{1 + v_0} \right) \quad \frac{\lambda}{\lambda_0} = \frac{f_0}{f} = \sqrt{\frac{1 + \beta}{1 - \beta}}, \beta = \frac{v}{c} \quad (175)$$

The “redshift”: $z = \frac{\lambda_0 - \lambda}{\lambda} = \frac{f - f_0}{f}$

107 Fission

Conservation of energy, binding energy of nucleus is always $(-)$, like e^- binding energy.

$$-BE_i + KE_i = -BE_f + KE_f \quad (176)$$

108 Wire Resistance

$$R = \frac{\rho L}{A} \quad (177)$$

109 Spin Matrices

$$S_i \psi = \frac{\hbar}{2} \sigma_i \psi \quad (178)$$

For example, eigenstate of S_x with $-\frac{\hbar}{2}$ and $\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

$$\frac{1}{\sqrt{2}}(|\uparrow\rangle - |\downarrow\rangle) = \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 1 \\ 0 \end{pmatrix} - \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right] = \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 1 \\ -1 \end{pmatrix} \right] \quad (179)$$

$$\begin{pmatrix} 1 \\ -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix} \rightarrow S_x \psi = \frac{\hbar}{2} \begin{pmatrix} 1 \\ \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = -\frac{\hbar}{2} \begin{pmatrix} 1 \\ \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} -1 \\ 1 \end{pmatrix} = -\frac{\hbar}{2} \psi \quad (180)$$

$$|\uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad |\downarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (181)$$

110 For What v Will a Car Stay on a Hill?

$$F_c = F_g \quad \frac{mv^2}{r} = mg \quad (182)$$

111 Bohr Model

1. e^- have classical motions
2. $\Delta E = hf$
3. Angular momentum quantized $L = n\hbar$
4. $E_n = -\frac{Z^2 E_0}{n^2} \quad E_n \propto \mu$
5. $\Delta E = E_0 \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right) \rightarrow \frac{1}{\lambda} = R_y \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right), R_y = 1 \times 10^7 \text{ m}^{-1}$
6. Positronium: $\mu = \frac{m_e}{2} \rightarrow E_p = \frac{E_0}{2n^2}$

112 Fluids

Equilibrium when $F_A = F_B$, where $F = mg$.

113 Gauss' with non-uniform

Must integrate.

$$\rho = Ar^2 \quad dV = 4\pi r^2 dr \quad \int E \cdot dA = \frac{Q_{\text{in}}}{\epsilon_0} = \frac{\int \rho dV}{\epsilon_0} = \frac{\int Ar^2 4\pi r^2 dr}{\epsilon_0} = \frac{44\pi r^5}{5\epsilon_0} \quad (183)$$

114 Capacitors

In series, $Q_1 = Q_2$ while in parallel, $V_1 = V_2$

115 Diffraction Limit

$$\sin \theta = \frac{1.22\lambda}{d} \quad d = \text{diameter of lens} \quad (184)$$

116 Normal Modes

1. Highest normal mode frequency when out of phase
2. Use limits if possible, with $M \rightarrow \infty$
3. # frequency = # masses
4. If odd # masses, one ω will be ω_0 , others above and below

For two hanging masses connected by a spring,

$$\text{In phase: } \omega = \sqrt{\frac{g}{l}} \quad \text{spring's } \Delta x = 0 \quad (185)$$

$$\text{Out of phase: } \omega = \sqrt{\frac{2k}{m} + \frac{g}{l}} \quad (186)$$

For three masses connected by two springs with the mass in the middle larger than the equal masses on the sides:

$$\omega = \sqrt{\frac{k}{m}}, \text{ like attached to a wall} \quad (187)$$

Side masses are in phase, middle mass is out of phase, $\omega = \sqrt{\frac{2k}{m}}$.

For 2 masses connected by 3 springs with the side masses connected to a wall:

$$\text{In phase: } \Delta x_1 = \Delta x_2 \text{ and } k' \text{ isn't expanded, } \omega = \sqrt{\frac{k}{m}} \quad (188)$$

$$\text{Out of phase: } \Delta x_1 = -\Delta x_2 \text{ and center of mass } k' \text{ stays in place } \omega = \sqrt{\frac{k + 2k'}{m}} \quad (189)$$

117 Radiation in Atoms

$n_f = \begin{matrix} \text{K} & \text{L} & \text{M} & \text{N} \\ 1 & 2 & 3 & 4 \end{matrix}$ series specify the final states coming from infinity, $n_i = \infty$ when being bombarded.

118 Ionization Energy

E required to liberate the outermost e^- . On the periodic table, increases in the $+y, +x$ direction.

119 Binding Energy

How tightly bound nucleons are

1. Peak at Fe/N \rightarrow elements $Z < Z_{\text{Fe}}$ undergo Fusion, $Z > Z_{\text{Fe}}$ undergo fission to release energy
2. When BE/nucleon increases, in reaction, energy is released
3. The mass of a nucleus is always less than the \sum particle's masses
4. "More tightly bound" = less mass/nucleon, more BE/nucleon
5. Created by the strong force
6. Energy given off in fusion/fission is the ΔE between fuel and products

120 Hierarchy of Forces

1. Strong
2. E&M
3. Weak
4. Gravity

121 Pair Production

1. Creation of elementary particle and anti-particle from photon
2. Cannot occur in free space, usually near a nucleus or other photon
3. For e^- , the photon E must exceed twice the rest energy of the e^- , ≈ 1 MeV
4. If 2 photons, 500 keV
5. Dominates at high E

122 Spectral Lines

1. Less dense gas \rightarrow more sharp and precise lines - don't lose E due to collisions
2. Sodium doublet - created by spin/orbit coupling, more pronounced in an external B

123 Photon Interactions with Matter

1. Compton Effect: low E , elastic scattering $< 10^6$ MeV
2. Photoelectric Effect: mid E , $< 10^7$ MeV
3. Pair Production: high E , $> 10^6$ MeV

124 Neutron

A fermion with:

$${}_0^1n \tag{190}$$

$$\text{Decay: } {}_0^1n \rightarrow {}_1^1p^+ + {}_{-1}^1e + \bar{\nu} \tag{191}$$

$$\text{Capture: } {}_1^1p^+ + {}_{-1}^1e \rightarrow {}_0^1n + \bar{\nu} \tag{192}$$

125 Deuteron

“Heavy Hydrogen”, ${}_1^2H$. Also, a boson.

126 Protium

A proton, ${}_1^1H$, Hydrogen nucleus, a fermion