

GRE Physics Study Notes

Courtesy of Nicole Duncan, transcribed by Jeren Suzuki

Last Edited 28th November 2012

Contents

1	Waves	1
1.1	Doppler Effect	1
2	Optics	1
2.1	Thin Lenses	1
2.2	Ray Diagrams	1
2.3	Index of Refraction	2
2.4	Telescope and Magnification	2
2.5	Thin Films	2
2.6	Resistance	2
3	Electricity and Magnetism	2
3.1	Gauss' Laws	2
3.2	Cyclotron	3
3.3	Conductivity / Current Density	3
3.4	Potential and Electric Field	3
3.4.1	Example Ring of Charge	3
3.5	Electrostatics	3
3.6	Classic E Examples	4
3.7	Parallel Plate Capacitor	4
3.7.1	Limits	4
3.8	Coulomb	4
3.9	Motion Through a Capacitor or Uniform Field	5
3.10	E of a Dipole	5
3.11	Gauss	5
3.12	Current Density	5
3.13	Drift Speed	5
3.14	Capacitors	5
3.15	Non-ohmic Materials	5
3.16	Convention of Battery	5
3.17	RC Circuit	6
3.18	Work	6
3.19	Magnetostatics	6
3.19.1	Magnetic Field	6
3.19.2	Current	6
3.19.3	Force	6
3.19.4	Cyclotron Motion	6
3.19.5	Cycloid	6
3.19.6	Solenoid	7
3.19.7	Ring of Current	7
3.19.8	Infinite Wire	7
3.19.9	Surface Current	7
3.19.10	Toroid	7
3.19.11	Dipole	7
3.19.12	Inductance	7
3.19.13	Maxwell's Equations in Matter	8
3.20	Dielectrics	8
3.20.1	Dipoles and Bound Charges	8
3.20.2	Dielectrics	8
3.20.3	Linear Dielectrics	8
3.21	Radiation	9
3.21.1	Electric Dipole	9
3.21.2	Point Charge	9
3.21.3	An Oscillating Sphere with Changing Radius	9
3.21.4	Magnetic Dipole Radiation	9
3.22	Maxwell Equations	9

3.23	Ampere's Law	9
3.24	Current	10
3.25	Boundary Conditions E&M Waves	10
3.26	E&M Fields	10
3.27	Energy Stored in E&M	10
3.28	Poynting Vector	10
3.29	Irradiance	10
3.30	Relativistic E&M	11
3.30.1	Example: Parallel Plate Capacitor	11
3.30.2	Special Cases	11
3.31	Coordinate Systems	11
3.32	Vectors	11
3.33	Diamagnetism	11
3.34	Paramagnetism	11
3.35	Ferromagnetism	12
3.36	Radiation Pressure	12
4	Circuits	12
4.1	Resistivity	12
4.2	Types of Cells	12
4.3	Band Pass	12
4.4	Low Pass	13
4.5	High Pass	13
5	Quantum Mechanics	13
5.1	Operators	13
5.2	Hermitian Operators	14
5.3	Transmission / Reflection / Tunneling Through Barrier	14
5.4	Hyperfine Splitting	14
5.5	Fine Structure	14
5.6	Zeeman Effect	15
5.7	Stark Effect	15
5.8	Degenerate Perturbation Theory	15
5.9	Non degenerate Perturbation Theory	15
5.10	Particle in a Box - Infinite Square Well	16
5.11	Schrödinger's Equation	16
5.12	Free Particle	16
5.12.1	Wave Packet Solutions	16
5.13	Traveling Wave Formalism	16
5.14	Finite Potential Well	17
5.15	Fundamental Particles	17
5.16	Single Slit Diffraction	17
5.17	Diffraction Grating	17
5.18	Double Slit Interference	17
5.19	Bragg Diffraction	17
6	Harmonics	18
6.1	Harmonic Oscillator Potential	18
6.2	Damped-Driven Oscillator	18
6.2.1	Underdamped $\omega_0 > \beta$	18
6.2.2	Overdamped $\omega_0 < \beta$	18
6.2.3	Critically Damped $\omega_0 = \beta$	18
6.3	Springs and Simple Harmonic Oscillators	18
6.4	Beats	19
7	Kinematics	19
7.1	Linear \rightarrow Rotational Kinematics	19
7.2	Lagrangian	19
7.2.1	EOMS:	19
7.3	Rocket Motion	19

7.4	Collisions	20
7.5	Central Force Motion	20
7.6	Moments of Inertia	20
8	Statistical Thermodynamics	21
8.1	Laws of Thermodynamics	21
8.1.1	1st Law	21
8.1.2	2nd Law	21
8.1.3	3rd Law	21
8.2	Maxwell Velocity Distribution	21
8.3	Mean Free Path	21
8.4	Particle Diffusion	21
8.5	Thermal Diffusion	21
8.5.1	Thermodynamic Identity	22
8.6	Heat Capacity	22
8.7	Isothermal Compression (Slow)	22
8.8	Adiabatic Compression (Fast)	22
8.9	Heat	23
8.10	Multiplicity/States	23
8.11	Boltzmann Statistics	23
8.12	Density of State Distributions	23
8.13	Blackbody Radiation	23
8.13.1	Wein's Law	23
8.13.2	Stephan-Boltzmann	23
8.14	Heat Engines	24
8.15	Refrigerators	24
8.16	Big People	24
9	Relativity	24
9.1	Space-Time Diagram	24
9.2	Special Relativity	25
9.2.1	Time Dilation	25
9.2.2	Length Contraction	25
9.2.3	Invariant Interval	25
9.2.4	Energy	25
10	Atomic Physics	25
10.1	Hydrogen Spectral Series	25
10.2	Atomic Notation	26
11	Particle Physics	26
11.1	Fermi	26
11.2	Degeneracy Pressure of a Solid	26
12	Misc	26
12.1	Water Density	26
12.2	Fundamental Law of Statistical Mechanics	26
12.3	Irreversible Process	27
12.4	Reversible Process	27

Introduction

By no means comprehensive, this list is meant to serve as additional study material to re-reading textbooks, practicing GRE tests, and nagging your physics friends for study help.

1 Waves

1.1 Doppler Effect

$$f = f_0 \left(\frac{v + v_s}{v + v_0} \right) \quad (1)$$

$$v_0 = \begin{cases} + & \text{away} \\ - & \text{towards} \end{cases}$$

$$v_s = \begin{cases} + & \text{towards} \\ - & \text{away} \end{cases}$$

$$\frac{\lambda}{\lambda_0} = \sqrt{\frac{1 + \beta}{1 - \beta}}, \beta = \frac{v}{c} \text{ for relativistic doppler shift} \quad (2)$$

$$= \frac{f_0}{f} \quad (3)$$

2 Optics

2.1 Thin Lenses

$$d_i = \frac{f \cdot d_o}{f - d_o} \quad (4)$$

$$M = \frac{h_i}{h_o} = -\frac{d_i}{d_o} \quad (5)$$

2.2 Ray Diagrams

1. Through f , || other side
2. Through center, continues along path
3. ||, goes through f on other side

2.3 Index of Refraction

$$n = \frac{c}{v} \quad (6)$$

$$v = v_Q = \frac{\omega}{k} = \sqrt{\frac{1}{\epsilon\mu}} \quad (7)$$

$$\lambda = \frac{\lambda_0}{n} \text{ inside a medium} \quad (8)$$

$$n = \frac{c}{v} = \sqrt{\frac{\epsilon\mu}{\epsilon_0\mu_0}} \quad (9)$$

2.4 Telescope and Magnification

2 lenses share a common focal point

$$M = -\frac{f_o}{f_e} = \frac{\theta_{\text{eye}}}{\theta_{\text{object}}} \quad (10)$$

$$d_o + d_e = f_o + f_e \quad (11)$$

2.5 Thin Films

$$\Delta\phi = \begin{cases} 0 & n_2 < n_1 \\ \pi & n_2 > n_1 \end{cases}$$

$$2d = \begin{cases} n\lambda/2 & \Delta\phi_{\text{tot}} = \pi \\ n\lambda & \Delta\phi_{\text{tot}} = 0, 2\pi \end{cases}$$

2.6 Resistance

$$R = \frac{\rho L}{A} = \frac{L}{\sigma A} \quad (12)$$

3 Electricity and Magnetism

3.1 Gauss' Laws

$$\int E \cdot da = \frac{Q_{\text{encl}}}{\epsilon_0} \quad (13)$$

$$\nabla \cdot E = \frac{\rho}{\epsilon_0} \quad (14)$$

$$\int B \cdot dl = \mu_0 I_{\text{encl}} (\text{in Amperes}) \rightarrow \int B \cdot da = 0 \quad (15)$$

$$\nabla \cdot B = 0 \quad (16)$$

$$\int g \cdot da = 4\pi MG \quad (17)$$

$$\nabla \cdot g = -4\pi G s \rho \quad (18)$$

3.2 Cyclotron

$$\omega = \frac{qB}{m} \quad (19)$$

$$F_c = F_B \rightarrow \frac{mv^2}{r} = qvB \quad (20)$$

$$v = \frac{qBr}{m} = r\omega \quad (21)$$

$$\omega = \frac{qB}{m} \quad (22)$$

3.3 Conductivity / Current Density

$$J = nq\bar{v} \quad (23)$$

$$J = \frac{ne^2\tau}{m}E = \sigma E, \quad \sigma = \frac{ne^2\tau}{m} \quad (24)$$

3.4 Potential and Electric Field

$$E = \int \frac{k dQ}{r^2} = k \int \frac{\sigma dA}{r^2} = k \int \frac{\rho dv}{r^2} = k \int \frac{\lambda dl}{r^2} \quad (25)$$

$$V = \int \frac{k dq}{r} \quad (26)$$

3.4.1 Example Ring of Charge

Imagine a ring with radius R and a point P above the ring at a height z making an angle θ above the ring plane.

$$E = k \int \frac{dQ}{r^2} \quad (27)$$

$$r^2 = R^2 + z^2 \quad (28)$$

$$dq = \lambda dl = Q \quad (29)$$

$$\sin \theta = \frac{z}{r} = \frac{E_z}{E} \quad (30)$$

$$E = \frac{kQ}{R^2 + z^2}, \text{ But } E = \hat{E}_z \quad (31)$$

$$E = \frac{kQ}{R^2 + z^2} \sin \theta = \frac{kQ}{R^2 + z^2} \frac{z}{r} = kQz \quad (32)$$

3.5 Electrostatics

$$F = \frac{kq_1q_2}{r^2}, \quad k_{\text{vacuum}} = \frac{1}{4\pi\epsilon_0}, \quad k_{\text{medium}} = \frac{1}{4\pi\epsilon}, \quad \epsilon = k\epsilon_0 \quad (33)$$

$$\nabla \cdot E = \frac{\rho_{\text{in}}}{\epsilon_0} \quad (34)$$

$$\nabla \times E = 0 \quad (35)$$

3.6 Classic E Examples

1. Sphere $\propto \frac{1}{r^2}$
2. Infinite Line $\propto \frac{1}{r}$
3. Infinite Plane doesn't fall off
 $E_{\text{plane}} = \frac{\sigma}{2\epsilon_0} \hat{n}$
4. Ring of charge:
 $E \propto \frac{x}{d^3} = \frac{x}{(x^2 + R^2)^{3/2}}$
5. Disk of charge:
 $E = \frac{\sigma}{2\epsilon_0} \left(1 - \frac{z}{(z^2 + R^2)^{1/2}} \right)$, σ = area charge density

3.7 Parallel Plate Capacitor

Model as infinite planes:

$$E_{\text{out}} = 0 \quad (36)$$

$$E_{\text{in}} = \frac{\sigma}{\epsilon_0} \quad (37)$$

3.7.1 Limits

1. As $x \rightarrow \infty$, all finite objects look like point charges
2. Sometimes must use binomial approximation to get behavior at ∞ . Disk of charge $\rightarrow 0$ if you don't use it.
3. $(1 + X)^n \sim 1 + nX$ for small x

3.8 Coulomb

$$E = k \int \frac{dQ}{r^2} \quad (38)$$

$$dQ = \lambda dl \sim \sigma dA \sim \rho dV \text{ , be careful of symmetry when integrating!} \quad (39)$$

For a ring of radius R in the x - y plane, a point is a distance r from the ring, making an angle θ on the z -axis. For ring of charge, must integrate by saying:

$$E = E_{\hat{z}} \quad (40)$$

$$\cos \theta = \frac{z}{r} \quad (41)$$

$$r = \sqrt{z^2 + R^2} \quad (42)$$

$$\lambda = \frac{Q}{2\pi R} \quad (43)$$

$$dl = R d\theta \quad (44)$$

$$dE_{\hat{z}} = dE \cos \theta = \frac{k\lambda dl}{r^2} \cos \theta \quad (45)$$

$$E = k\lambda \int \frac{dl}{r^2} \cos \theta = \frac{kQ}{2\pi R} (R) \int_0^{2\pi} \frac{dQ}{r^2} \frac{z}{r} \quad (46)$$

$$E = \frac{kQ}{2\pi} (2\pi) \frac{z}{r^3} = \frac{kQz}{(R^2 + z^2)^{3/2}} \quad (47)$$

3.9 Motion Through a Capacitor or Uniform Field

Kinematics equation: $F = ma = eE$. Find v_c, a, t to get θ deflection

3.10 E of a Dipole

$$\bar{p} = qd \quad (48)$$

$$E_{\text{dipole}} = \begin{cases} \frac{k2\bar{p}}{r^3} & \text{on axis of } \hat{d} \\ \frac{k\bar{p}}{r^3} & \text{plane perpendicular to } \hat{d} \end{cases}$$

3.11 Gauss

$$\Phi_E = \int E \cdot dA = \frac{Q_{\text{enclosed}}}{\epsilon_0} \quad (49)$$

3.12 Current Density

Continuity Equation:

$$\frac{\partial \rho}{\partial t} = -\nabla \cdot \bar{J} \quad (50)$$

$$J = q_\alpha n_\alpha v_\alpha, I = JA = \frac{\text{current}}{m^2} \text{ of the cross section} = \frac{A}{m^2} \quad (51)$$

3.13 Drift Speed

$$v_{\text{drift}} = \frac{e\tau E}{m} \quad (52)$$

3.14 Capacitors

C depends upon geometry of electrodes

3.15 Non-ohmic Materials

Do not obey $V = IR$: batteries, semiconductors, capacitors, inductors

3.16 Convention of Battery

Long side of battery is positive.

3.17 RC Circuit

$$Q = Q_0 e^{-t/\tau} \quad (53)$$

$$I = I_0 e^{-t/\tau}, \tau = RC \quad (54)$$

$$Q = VC \quad (55)$$

$$V = V_0 e^{-t/\tau}, \text{ decay} \quad (56)$$

$$= V_0(1 - e^{-t/\tau}), \text{ charging up} \quad (57)$$

3.18 Work

$$W = F \cdot d = eE \cdot d = e\Delta V \quad (58)$$

3.19 Magnetostatics

3.19.1 Magnetic Field

Biot-Savart Law:

$$B = \frac{\mu_0}{4\pi} \frac{\bar{I} \times \hat{r}}{r^2} = \frac{\mu_0}{4\pi} I \frac{d\bar{l} \times \hat{r}}{r^2}, d\bar{l} = \text{the actual length of the segment, not just the direction} \quad (59)$$

$$\text{Tesla} = T = \frac{N}{A \cdot m} \quad (60)$$

3.19.2 Current

$$I = \int J da_{\perp} \quad (61)$$

$$\text{if } J = Kr, I = \int_0^{2\pi} \int_0^r kr'(r' dr' d\phi) = \frac{2\pi}{3} kr^3 \quad (62)$$

3.19.3 Force

$$\bar{F} = q\bar{v} \times B = I(d\bar{l} \times \bar{B}) \quad (63)$$

3.19.4 Cyclotron Motion

$$v_{\parallel} B \rightarrow \text{helical} \quad (64)$$

$$\frac{mv^2}{r} = qvB \quad (65)$$

3.19.5 Cycloid

E in $+z$ direction and B in $+x$ direction with particle traveling in $+y$ direction make a cycloid.

3.19.6 Solenoid

$$B = \begin{cases} \mu_0 n I \hat{z} & \text{inside, } n = \frac{N}{L} \\ 0 & \text{outside} \end{cases}$$

3.19.7 Ring of Current

$$B = \frac{\mu_0 I}{2R} \quad (66)$$

Any displacement along center of ring should reduce to this equation as $x \rightarrow 0$, as $x \rightarrow \infty$ should be field of dipole.

3.19.8 Infinite Wire

$$B = \frac{\mu_0 I}{2\pi r}, \text{ in limits, } \theta_1 = -\frac{\pi}{2}, \theta_2 = \frac{\pi}{2}, \text{ and } r \text{ is the distance from the wire} \quad (67)$$

3.19.9 Surface Current

$$B = \begin{cases} -\frac{\mu_0}{2} & z > 0 \\ \frac{\mu_0}{2} & z < 0 \end{cases}, \text{ use Amperian square loop}$$

3.19.10 Toroid

$$B = \begin{cases} \frac{\mu_0 I N}{2\pi r} \hat{\phi} & \text{inside} \\ 0 & \text{outside} \end{cases}$$

3.19.11 Dipole

$$B \propto \frac{\mu}{r^3} \quad (68)$$

$$\mu = IA, x \rightarrow \infty \text{ limit looks like this} \quad (69)$$

Field far away from a current loop = field of a dipole.

Magnetic fields do no work.

3.19.12 Inductance

$$\Phi = LI \quad (70)$$

$$\epsilon = -L \frac{dI}{dt} \quad (71)$$

$$\Phi_B = \int B \cdot dA \quad (72)$$

$$\text{Henry} = H = \frac{Vs}{A} \quad (73)$$

Inductor in serial with resistor: $\tau = \frac{L}{R}$.

$$W = \frac{1}{2}LI^2 = U_{\text{stored}} \quad (74)$$

L is like mass, the greater the L the harder it is to try and change the current.

VLR Circuit: Voltage log's to V.

Ohms':

$$\epsilon_0 - L \frac{dI}{dt} = IR = V \quad (75)$$

Solution to differential equation:

$$I(t) = \frac{\epsilon_0}{R} + ke^{-(R/L)t}, \tau = \frac{L}{R} \quad (76)$$

If $t = 0, V = 0$, just plugged in, $k = -\frac{\epsilon_0}{R}$

$$I(t) = \frac{\epsilon_0}{R} \left(1 - e^{-(k/L)t}\right) \quad (77)$$

3.19.13 Maxwell's Equations in Matter

$$\begin{array}{llll} \nabla \cdot D = \rho_f & \nabla \times \bar{E} = -\frac{\partial B}{\partial t} & D = \epsilon E & \epsilon = \epsilon_0(1 + \chi_e) \\ \nabla \cdot B = 0 & \nabla \times H = \bar{J}_f + \frac{\partial D}{\partial t} & B = \mu H & \mu = \mu_0(1 + \chi_m) \end{array}$$

3.20 Dielectrics

3.20.1 Dipoles and Bound Charges

$$\begin{array}{lll} \rho_b = -\bar{\nabla} \cdot \bar{p} & \sigma_b = \bar{p} \cdot \hat{n} & \bar{p} = q\bar{d} \\ \tau = \bar{p} \times \bar{E} & \bar{U} = -\bar{p} \cdot \bar{E} & \end{array}$$

3.20.2 Dielectrics

1. Electric Displacement: $\bar{D} = \epsilon_0 \bar{E} + \bar{p}$
2. Gauss' Law:

$$\nabla \cdot D = \rho_{\text{free}} \quad (78)$$

$$\int D \cdot da = Q_{\text{enclosed}} \quad (79)$$

3.20.3 Linear Dielectrics

Conduction:

$$\bar{p} = \epsilon_0 \underbrace{\chi_e}_{\text{electric susceptibility}} \bar{E} \quad (80)$$

$$F = \frac{1}{4\pi \underbrace{\epsilon}_{\text{only thing that changes } (\epsilon_0 \rightarrow \epsilon)}} \frac{qQ}{r^2} \quad (81)$$

only thing that changes ($\epsilon_0 \rightarrow \epsilon$)

$$= \frac{1}{4\pi \epsilon_0 \epsilon_r} \frac{qQ}{r^2} = \frac{F_{\text{vac}}}{\epsilon_r} = F_{\text{medium}} \quad (82)$$

$$E_{\text{medium}} = \frac{E_{\text{vac}}}{\epsilon_r} \rightarrow E = \frac{E_0}{k} \quad (83)$$

1. Permittivity = ϵ
2. Dielectric Constant: $\epsilon_r = \frac{\epsilon}{\epsilon_0}$
3. Displacement: $\bar{D} = \epsilon \bar{E}$

3.21 Radiation

3.21.1 Electric Dipole

$$P \propto q^2 \omega^4 d^2 \quad (84)$$

$$\langle s \rangle \propto \frac{q^2 d^2 \omega^4}{r^2} \sin^2 \theta \quad (85)$$

Where the $\sin^2 \theta$ component is so we don't see along the direction of motion.

3.21.2 Point Charge

$$P \propto q^2 a^2 \quad (86)$$

$$\langle s \rangle \propto \frac{q^2 a^2 \sin^2 \theta}{r^2} \quad (87)$$

Once again, no power radiated along motion direction. $\langle s \rangle_{\max}$ @ $\theta = 90$ to motion.

3.21.3 An Oscillating Sphere with Changing Radius

...emits no radiation. Use Gauss' law for symmetry problems, E is constant. an uncharged particle accelerates more than a charged particle because the charged particle emits radiation, $\bar{F}_{\text{in}} - \hat{d}$.

3.21.4 Magnetic Dipole Radiation

1. Model a wire loop with alternating current
2. $P \propto b^4 I_0^2 \omega^4$
3. $\langle s \rangle \propto \frac{b^4 I_0^2 \omega^4 \sin^2 \theta}{r^2}$

3.22 Maxwell Equations

$$\begin{aligned} \nabla \cdot E &= \frac{\rho}{\epsilon_0} & \nabla \times E &= -\frac{\partial B}{\partial t} \\ \nabla \cdot B &= 0 & \nabla \times B &= \mu_0 J + \mu_0 \epsilon_0 \frac{\partial E}{\partial t} \end{aligned}$$

Magnetic monopoles would symmetrize the equations... *wings hands*

$$\begin{aligned} \oint E \cdot dA &= \frac{Q_{\text{in}}}{\epsilon_0} & \oint E \cdot dl &= -\frac{\partial \Phi_B}{\partial t} \\ \oint B \cdot dA &= 0 & \oint B \cdot dl &= \mu_0 I + \mu_0 \epsilon_0 \frac{\partial \Phi_E}{\partial t} \end{aligned}$$

3.23 Ampere's Law

$$\oint B \cdot dl = \mu_0 I_{\text{enclosed}} \quad (88)$$

3.24 Current

$$I = \int J \cdot dA \quad (89)$$

3.25 Boundary Conditions E&M Waves

1. $E_{\parallel} = 0 \quad B_{\perp} = 0 \rightarrow$ reflections
2. For reflection, $E_{\text{tot}} = 0 \quad B_{\text{tot}} = 2B_{\text{wave}}$
3. E_{\perp} is always discontinuous by $\frac{\sigma}{\epsilon_0}$ @ boundary
4. E_{\parallel} is always continuous

$$\begin{aligned} \epsilon_1 E_1 - \epsilon_2 E_2^{\perp} &= \sigma_p & E_1^{\parallel} &= E_2^{\parallel} \\ B_1^{\perp} &= B_2^{\perp} & \frac{1}{\mu_1} B_1^{\parallel} - \frac{1}{\mu_2} B_2^{\parallel} &= \underbrace{k_f}_{\text{free current}} x \hat{n} \end{aligned}$$

3.26 E&M Fields

E/B are in phase and perpendicular

1. $B_0 = \frac{k}{\omega} E_0 = \frac{1}{c} E_0$
2. Radiation Pressure: $p = \frac{\langle s \rangle}{c}$
3. Energy Density: $\langle U \rangle = \frac{1}{2} \epsilon_0 E^2$
4. $\vec{s} = \frac{1}{\mu_0} (\vec{E} \times \vec{B})$
5. Intensity: $I = \langle s \rangle = \frac{1}{2} c \epsilon_0 E^2$
6. \hat{s} = propagation of E&M field

3.27 Energy Stored in E&M

$$U = \epsilon_0 E^2 = \frac{1}{\mu_0} B^2, U_E = U_B \quad (90)$$

3.28 Poynting Vector

$$\vec{S} = \frac{1}{\mu_0} \vec{E} \times \vec{B} \quad (91)$$

3.29 Irradiance

$$I = \langle s \rangle \quad (92)$$

$$= c \epsilon_0 \langle E^2 \rangle \quad (93)$$

$$= \frac{c}{\mu_0} \langle B^2 \rangle \quad (94)$$

3.30 Relativistic E&M

1. E&M consistent with relativity
2. Between reference frames the E&M processes change but particle motion and outcome is always the same
3. Charge is invariant

3.30.1 Example: Parallel Plate Capacitor

$$S : E^\perp = \frac{\sigma_0}{\epsilon_0} \hat{y} \quad (95)$$

$$S' : E^\perp = \frac{\sigma}{\epsilon_0} \hat{y}, \text{ only } \sigma \text{ changes} \quad (96)$$

Charge on each plate is invariant, width is unchanged, but the length (along direction of motion) is contracted.

$$l = \frac{l_0}{\gamma} \rightarrow \sigma = \frac{\sigma_0}{\gamma} \quad (97)$$

For motion in \hat{x} , $E_{\hat{y}}$ is changed while $E_{\hat{x}}$ is unchanged since $E = E_{\hat{y}}$.

$$E_\perp = \gamma E_\perp \quad (98)$$

$$E_\parallel = E_\parallel \quad (99)$$

3.30.2 Special Cases

If $B = 0$ in any one reference frame,

$$\bar{B} = -\frac{1}{c^2}(\bar{V} \times \bar{E}) \quad (100)$$

If $E = 0$ in any one reference frame,

$$\bar{E} = \bar{V} \times \bar{B} \quad (101)$$

3.31 Coordinate Systems

1. Cartesian: $dl = dx\hat{x} + dy\hat{y} + dz\hat{z}$, $dV = dxdydz$
2. Spherical: $dl = dr\hat{r} + r d\theta\hat{\theta} + r \sin\theta d\phi\hat{\phi}$, $dV = r^2 \sin\theta dr d\theta d\phi$
3. Cylindrical: $dl = ds\hat{s} + s d\phi\hat{\phi} + dz\hat{z}$, $dV = s ds d\phi dz$

3.32 Vectors

$$\nabla \times (\nabla \times A) = \nabla(\nabla \cdot A) - \nabla^2 A \quad (102)$$

3.33 Diamagnetism

Caused by change in orbital moment (μ) induced by B . Acts to negate B , anti-parallel to B .

3.34 Paramagnetism

In a magnetic field, breaking of energy levels by spin/spin or spin/orbit coupling induced along B .

3.35 Ferromagnetism

Any material that exhibits a spontaneous B . (A net magnetic moment in the absence of an external B)

3.36 Radiation Pressure

Energy Density of the wave

$$P = U = V_e + U_B \quad (103)$$

$$\langle p \rangle = \frac{\langle s \rangle}{c} \quad (104)$$

$$(105)$$

1. Perfect reflection: light enters with $+c$ and exits with $-c$

$$\text{so } \Delta v = 2c \rightarrow \langle p \rangle = \frac{2\langle s \rangle}{c}$$

Curl-less Fields: \vec{E}

$$\nabla \times \vec{F} = 0 \text{ everywhere}$$

$$\int_b \vec{F} \cdot d\vec{l} = \text{pattern independent}$$

$$\oint_a \vec{F} \cdot d\vec{l} = 0 \text{ closed loop}$$

$$\vec{F} = -\nabla V$$

Div-less Fields: \vec{B}

$$\nabla \cdot \vec{F} = 0$$

$$\int \vec{F} \cdot d\vec{A} = \text{independent of any bound line}$$

$$\oint \vec{F} \cdot d\vec{A} = 0 \text{ for all surfaces}$$

$$\vec{F} = \vec{\nabla} \times \vec{A}$$

4 Circuits

4.1 Resistivity

$$\rho(T_2) = \rho(T_1)(1 + \alpha\Delta T) \quad (106)$$

For metals:

1. $\alpha = (+)$
2. $\rho \uparrow \quad T \uparrow$
3. Doping increases ρ

while for semiconductors:

1. $\alpha = (-)$
2. $\rho \downarrow \quad T \uparrow$
3. Doping decreases ρ

4.2 Types of Cells

1. Conventional Cell: Contains more than 1 lattice point.
2. Primitive Cell: Contains 1 lattice point. $V_{cc}/N_{cc \text{ lattice points}} = V_{pc}$

4.3 Band Pass

$$\omega_0 \rightarrow \omega_0 L = \frac{1}{\omega_0 C} \rightarrow \omega_0 = \frac{1}{\sqrt{LC}} \quad (107)$$

4.4 Low Pass

Either looks like a RC or LR circuit, but perpendicular to each other. Purpose is to cut out high frequencies, essentially letting low frequencies pass through.

$$T_1 = \frac{\frac{1}{j\omega C}}{R + \frac{1}{j\omega C}} \quad (108)$$

$$= \frac{\frac{1}{j\omega C}}{\frac{Rj\omega C + 1}{j\omega C}} \quad (109)$$

$$= \frac{1}{1 + j\omega CR} \quad (110)$$

1. As $\omega \rightarrow \infty$, $T_1 \rightarrow 0$

2. As $\omega \rightarrow 0$, $T_1 \rightarrow 1$

$$T_2 = \frac{R}{j\omega C + R} \quad (111)$$

1. As $\omega \rightarrow \infty$, $T_2 \rightarrow 0$

2. As $\omega \rightarrow 0$, $T_2 \rightarrow 1$

4.5 High Pass

Either looks like a CR or RL circuit, like a low pass filter configuration but with elements reversed. Cuts out low frequencies.

$$T_1 = \frac{R}{R + \frac{1}{j\omega C}} \quad (112)$$

$$= \frac{R}{\frac{j\omega CR + 1}{j\omega C}} \quad (113)$$

$$= \frac{j\omega CR}{j\omega CR + 1} \quad (114)$$

1. As $\omega \rightarrow \infty$, $T_1 \rightarrow 1$

2. As $\omega \rightarrow 0$, $T_1 \rightarrow 0$

$$T_2 = \frac{j\omega L}{R + j\omega L} \quad (115)$$

1. As $\omega \rightarrow \infty$, $T_2 \rightarrow 1$

2. As $\omega \rightarrow 0$, $T_2 \rightarrow 0$

5 Quantum Mechanics

5.1 Operators

$$\hat{x} = x \quad (116)$$

$$\hat{H} = -\frac{\hbar^2}{2m}\nabla^2 + V \quad (117)$$

$$\hat{p} = -i\hbar\frac{\partial}{\partial x} \quad (118)$$

5.2 Hermitian Operators

1. Represent observables
2. $\langle \hat{Q} \rangle = \langle \Psi | \hat{Q} | \Psi \rangle = \text{real} \#$
3. Conditions: $\langle f | \hat{Q} g \rangle = \langle \hat{Q} f | g \rangle \Rightarrow \hat{a}^\dagger = \hat{a}^* = \hat{a}$
4. Determinant states are eigenfunctions of \hat{Q}
5. $\left(\frac{\partial}{\partial x}\right)^\dagger = -\frac{\partial}{\partial x}$, note

5.3 Transmission / Reflection / Tunneling Through Barrier

1. Incident: Ae^{ikx}
2. Reflection: Re^{-ikx}
3. Transmission: Te^{-ikx}
4. Limits:
 $v_0 \rightarrow 0$, $R \rightarrow 0$
 $v_0 \rightarrow \infty$, $T \rightarrow 0$
5. Probability(Transmission) = $|T/A|^2$
6. Probability(Reflection) = $|R/A|^2$
7. Probability(Transmission) + Probability(Reflection) = 1
 $T^2 + R^2 = A^2$
8. Tunneling Depth $d \propto \frac{1}{\sqrt{V-E}}$

5.4 Hyperfine Splitting

1. Spin/spin of e^- nucleus
2. Responsible for 21 cm line

$$\mu_p = \frac{ge}{Zm_p} \overline{s_p} \quad (119)$$

$$\mu_e = \frac{-e}{m_e} \overline{s_e} \quad (120)$$

$$E_{n'_f} = \frac{\mu_0 g_p e^2}{3\pi m_p m_e a^3} \langle \overline{s_p} \cdot \overline{s_e} \rangle \quad (121)$$

$$E_{n'_f} \propto \frac{e^2}{m_p m_e a^3} \langle \overline{s_p} \cdot \overline{s_e} \rangle \quad (122)$$

5.5 Fine Structure

1. Spin/orbit coupling + relativistic correction
2. Breaks l degeneracy, retains j degeneracy
3. Why $E_{2s} < E_{2p}$

5.6 Zeeman Effect

1. Atom in external \vec{B}
2. Spin+orbital angular momentum/B coupling
3. $H_{z'} = (-\bar{\mu}_e + \bar{\mu}_s) \cdot \vec{B}_{\text{ext}}$
4. Weak $B_{\text{ext}} \ll B_{\text{int}} \rightarrow E' = \mu_b g_j \underbrace{m_j}_{\text{breaks } m_i \text{ degeneracy into } 2j+1 \text{ levels}} B_{\text{ext}}$
5. Strong $B_{\text{ext}} \gg B_{\text{int}} \rightarrow E' = \mu_b B_{\text{ext}}(m_l + 2m_s)$

5.7 Stark Effect

1. External \vec{E}
2. not spin dependent
3. $H' = eE_z$ if $E = \hat{E}_z$
4. Hydrogen, $E'_1 = \langle H' \rangle = eE \int_0^\infty d^3r \underbrace{z}_{\text{odd}} \underbrace{|\Psi_{100}|^2}_{\text{even}} = 0$

5.8 Degenerate Perturbation Theory

1. A state when n degenerate states breaks into n distinct E levels
2. Tensor, $w_{aa}, w_{bb}, w_{cc} = E_a, E_b, E_c$ of unperturbed states
- 3.

$$\begin{pmatrix} w_{aa} & w_{ab} \\ w_{ba} & w_{bb} \end{pmatrix} \Rightarrow w_{ab} = w_{ba}^*$$

5.9 Non degenerate Perturbation Theory

1. $H = H' + H^0$
2. First order: $E'_n = \langle \Psi_n | H' | \Psi_n \rangle = \langle H' \rangle$

$$\Psi_{n'} \sum_{m,n} \frac{\langle \Psi_m^0 | H' | \Psi_n^0 \rangle}{E_n^0 - E_m^0} \Psi_m^0 \quad (123)$$

1. If E introduced

$$H' = eE \rightarrow E' = 0 \quad (124)$$

1. Potential raised by constant

$$H' = v_0 \rightarrow E' = v_0 \quad (125)$$

5.10 Particle in a Box - Infinite Square Well

$$E_n = n^2 E_0 \quad (126)$$

$$E_0 = \frac{\hbar^2 k_0^2}{2m} = \frac{p_0^2}{2m} \quad (127)$$

$$k_n = \frac{n\pi}{a} \quad (128)$$

$$p_n = \hbar k_n \quad (129)$$

$$\psi = \sqrt{\frac{2}{a}} \sin(k_n x) \quad (130)$$

$$\text{3D: } E = \frac{\hbar^2}{2m} [k_x^2 + k_y^2 + k_z^2] \quad (131)$$

5.11 Schrödinger's Equation

$$\left(-\frac{\hbar^2}{2m} \nabla^2 + V \right) \Psi = i\hbar \frac{\partial \Psi}{\partial t} \quad (132)$$

Separable Solutions:

$$\Psi = \Phi(t)\Psi(x) \quad (133)$$

$$\Phi(t) = e^{-iE_n t/\hbar} \quad (134)$$

5.12 Free Particle

$$\Psi = Ae^{i(kx - \omega t)} \quad (135)$$

5.12.1 Wave Packet Solutions

$$\Psi = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Phi(k) e^{ikx} dk \quad (136)$$

$$\Phi = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \Psi(x) e^{-ikx} dx \quad (137)$$

1. Packet moves at group velocity, $v_g = \frac{\partial \omega}{\partial k}$

2. $\Delta x \Delta k \sim 1$

3. $\Delta x \Delta p \sim \hbar$, $p = \hbar k$

5.13 Traveling Wave Formalism

$$\frac{\partial^2 \Psi}{\partial x^2} = \frac{1}{v^2} \frac{\partial^2 \Psi}{\partial t^2}, v = \sqrt{\frac{\text{restoring force}}{\text{density}}} \quad (138)$$

$$v_\phi = \frac{\omega}{k} \quad (139)$$

$$\Psi = A \cos(k(vt - X)) = A \cos(\omega t - kx) \quad (140)$$

In one period, $x - vT = 2\pi$

5.14 Finite Potential Well

$$E \propto n^2 \quad (141)$$

$$d \propto \frac{1}{\sqrt{V - E_n}}, d = \frac{\hbar}{\sqrt{2m(V - E_N)}} \quad (142)$$

$$d \propto n \quad (143)$$

5.15 Fundamental Particles

1. Bosons: Force carriers
 Gauge Boson: Gluon-strong
 W,Z Boson - a.k.a Weak Boson
 photons - E&M
 other: Higgs, graviton, pion
2. Fermions: Associated with matter
 Quarks: up, down, top, bottom, strange, charm
 Leptons: electron, muon, tauon, neutrino flavors of each
3. Composite Fermions: Protons and Neutrons, etc.

5.16 Single Slit Diffraction

$$w \sin \theta = n\lambda, \quad \tan \theta = \frac{y}{L} \quad (144)$$

Central maximum width:

$$\frac{2L\lambda}{d} = \Delta y_{\max} \quad (145)$$

5.17 Diffraction Grating

$$d \sin \theta = n\lambda \quad (146)$$

$$y = L \tan \theta = L \frac{\sin \theta}{\cos \theta} = \frac{Ln\lambda}{d \cos \theta} \quad (147)$$

5.18 Double Slit Interference

$$d \sin \theta = n\lambda, d \sin \theta = n \left(\lambda + \frac{\lambda}{2} \right) \quad (148)$$

5.19 Bragg Diffraction

$$2d \sin \theta = n\lambda \quad (149)$$

$$d = \frac{\overbrace{a}^{\text{lattice spacing}}}{\underbrace{\sqrt{h^2 + k^2 + l^2}}_{\text{miller indices}}} \quad (150)$$

6 Harmonics

6.1 Harmonic Oscillator Potential

$$E_n = \hbar\omega \left(n + \frac{1}{2} \right) , \text{ lowest } n = 0 \quad (151)$$

$$\langle v \rangle = \langle T \rangle = \frac{1}{2} \hbar\omega \left(n + \frac{1}{2} \right) \quad (152)$$

$$\omega = \sqrt{\frac{k}{m}} \quad (153)$$

$$X = A \sin \omega t + B \cos \omega t \quad (154)$$

$$\Psi_n \propto e^{-\frac{m\omega x^2}{2\hbar}} H_n(x) \quad (155)$$

6.2 Damped-Driven Oscillator

$$F = -\underbrace{kx}_{\text{Hooke's}} - \overbrace{b\dot{x}}^{\text{dampening} \propto v} + \underbrace{A \cos \theta}_{\text{driver}} \quad (156)$$

$$\omega_0 = \sqrt{\frac{k}{m}} \quad (157)$$

$$\beta = \frac{b}{2m} \quad (158)$$

6.2.1 Underdamped $\omega_0 > \beta$

$$X_u = Ae^{-\beta t} \cos(\omega' t + \phi) , \omega' = \sqrt{\omega_0^2 - \beta^2} \quad (159)$$

6.2.2 Overdamped $\omega_0 < \beta$

$$X_o = Ae^{-\beta t} e^{-\omega'' t} , \omega'' = \sqrt{\beta^2 - \omega_0^2} \quad (160)$$

6.2.3 Critically Damped $\omega_0 = \beta$

$$X_c = A_1 e^{-\omega_0 t} + A_2 t e^{-\omega_0 t} \quad (161)$$

6.3 Springs and Simple Harmonic Oscillators

$$F = -kx \Rightarrow U = \frac{1}{2} kx^2 , \omega = \sqrt{\frac{k}{m}} \quad (162)$$

$$ma = -kx \quad (163)$$

$$\ddot{x} = -\omega_0^2 x = -\frac{k}{m} x \quad (164)$$

Solutions: sines and cosines

1. A = max amplitude

2. $E_{tot} = \frac{1}{2}KA^2$
3. $KE = \frac{1}{2}KA^2 \cos^2(\omega_0 t)$
4. $PE = \frac{1}{2}KA^2 \sin^2(\omega_0 t)$

To find oscillations about the minimum of E in an arbitrary u :

1. Find equilibrium value: $\frac{\partial u}{\partial x} = 0 \rightarrow x_0 = ?$
2. 2nd derivative of taylor series gives $\omega_a \rightarrow \frac{1}{2}v''(x_0) = \frac{1}{2}m\omega^2$

6.4 Beats

$$f_b = f_1 - f_2 \quad (165)$$

$$T_b = \frac{1}{f_1 - f_2} \quad (166)$$

7 Kinematics

7.1 Linear \rightarrow Rotational Kinematics

$$\begin{array}{lll} x \rightarrow \theta & s_{\text{arc}} = r\theta & \Delta x = v_0 t + \frac{1}{2}at^2 \rightarrow \Delta\theta = \omega_0 t + \frac{1}{2}\alpha t^2 \\ v \rightarrow \omega & v_{\perp} = r \times \omega & v = v_0 + at \\ a \rightarrow \alpha & a_{\perp} = r \times \alpha & v^2 = v_0^2 + 2a\Delta x \\ p \rightarrow L & L = r \times p & L = I\omega \quad (p = mv) \\ F \rightarrow \tau & \tau = r \times F & \tau = \frac{\partial L}{\partial t} \quad (F = \frac{\partial p}{\partial t}) \\ m \rightarrow I & I \propto mr^2 & \end{array}$$

7.2 Lagrangian

$$\begin{array}{ll} L = T - U & H = T + U \text{ if } U \neq U(v) \neq U(t) \\ \underbrace{\frac{\partial L}{\partial q} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right)}_{\text{EOMS}} = 0 & p = \frac{\partial L}{\partial \dot{q}} \end{array}$$

7.2.1 EOMS:

$$\dot{q}_k = \frac{\partial H}{\partial p_k} \quad (167)$$

$$-\dot{p}_k = \frac{\partial H}{\partial q_k} \quad (168)$$

7.3 Rocket Motion

$$U \frac{dm}{dt} + M \frac{dv}{dt} = 0 \quad (169)$$

$$v_f = v_0 + u \ln \left(\frac{M_i}{M_f} \right) \quad (170)$$

7.4 Collisions

1. Momentum + mass are always conserved classically
2. Use p equalities for before/after collisions even if elastic
3. Elastic \rightarrow conservation of kinetic energy

$$\epsilon = 1 = \frac{\overbrace{|v_1| + |v_2|}^{\text{final}}}{\underbrace{|u_1| + |u_2|}_{\text{before}}} \quad (171)$$

Don't forget to include $(-)$ and $(+)$ for direction of velocity in momentum equations!

1. Only use kinetic energy for conservation of total energy either before or after the collision
2. Impulse $J = F\Delta t = \Delta p = \Delta L$
3. Cross section:

$$N_{\text{scat}} = \frac{N_{\text{target}}}{\text{area}} N_{\text{incident}} \sigma$$

7.5 Central Force Motion

$$\mu = \frac{m_1 \cdot m_2}{m_1 + m_2} \quad (172)$$

$$r_{\text{CM}} = \frac{\sum_i m_i r_i}{\sum_i m_i} \quad (173)$$

$$T = \frac{1}{2} \mu |\dot{r}|^2 \quad (174)$$

$$\bar{r}_1 = \frac{m_2}{m_1 + m_2} \bar{r} \quad (175)$$

$$\bar{r}_2 = \frac{m_1}{m_1 + m_2} \bar{r} \quad (176)$$

$$\bar{r} = \bar{r}_1 - \bar{r}_2 \quad (177)$$

7.6 Moments of Inertia

1. $I = CMR^2$, where C is a constant
2. $I_{\text{hoop}} = MR^2$
3. $I_{\text{disk}} = \frac{1}{2} MR^2$
4. $I_{\text{hollow sphere}} = \frac{2}{3} MR^2$
5. $I_{\text{solid sphere}} = \frac{2}{5} MR^2$
6. $I_{\text{point mass}} = MR^2$
7. $I_{\text{rod end}} = \frac{1}{3} ML^2$
8. $I_{\text{rod center}} = \frac{1}{12} ML^2$
9. $L_{\text{rot}} = I\omega$
10. $T_{\text{rot}} = \frac{1}{2} I\omega^2$

11. $\tau = I dv = \frac{dL}{dt}$
12. $I_{\text{parallel axis}} = I_{\text{CM}} + MR_{\text{displaced}}^2$

8 Statistical Thermodynamics

8.1 Laws of Thermodynamics

8.1.1 1st Law

$$\Delta U = Q + W \quad (178)$$

8.1.2 2nd Law

E flows spontaneously until the system is at the most likely microstate \Rightarrow entropy tends to increase

8.1.3 3rd Law

$$S(T = 0) = 1, \text{ so } C_v \rightarrow 0 \text{ as } T \rightarrow 0 \quad (179)$$

8.2 Maxwell Velocity Distribution

Speed of molecules in ideal gas:

$$D(v) \propto v^2 e^{-E/k_b T} \quad (180)$$

8.3 Mean Free Path

$$l = \frac{1}{n\sigma} \quad (181)$$

$$n = \frac{\text{particles}}{\text{volume}} \quad (182)$$

$$\sigma = \text{scattering cross section} \quad (183)$$

8.4 Particle Diffusion

Fick's Law:

$$J_p = - \underbrace{D}_{\text{constant}} \nabla n \quad (184)$$

8.5 Thermal Diffusion

Fourier's Law:

$$J_q = \Phi_q = - \underbrace{\sigma}_{\text{conductivity (Thermal)}} \nabla T \quad (185)$$

$$\underbrace{\Phi_q}_{\text{Flow of energy/time} \cdot \text{area, units of } \frac{\text{W}}{\text{m}^2}} = -k \nabla T, k = \text{thermal conductivity with units} = \frac{\text{W}}{\text{m degrees K}} \quad (186)$$

Type of Interaction	Quantity	Variable	Formula
Mechanical	volume	P	$P = -\left(\frac{\partial U}{\partial V}\right)_{U,N} = T\left(\frac{\partial S}{\partial V}\right)_{U,N}$
Thermal	Temperature/Energy	T	$T = \left(\frac{\partial U}{\partial S}\right)_{V,N}$
Diffusive	Particles	μ	$\mu = -\left(\frac{\partial U}{\partial N}\right) = T\left(\frac{\partial S}{\partial N}\right)$

8.5.1 Thermodynamic Identity

$$dU = TdS - PdV + \mu dN \quad (187)$$

8.6 Heat Capacity

$$c = \frac{dQ}{dt} \quad (188)$$

$$c_p = \left(\frac{\partial Q}{\partial T}\right)_P = T\left(\frac{\partial S}{\partial T}\right) \quad (189)$$

$$c_v = \left(\frac{\partial Q}{\partial T}\right)_V = \left(\frac{\partial U}{\partial T}\right)_V, \quad U = \text{total } E \quad (190)$$

$c_P > c_V$ since at constant P the system loses E in the form of work \Rightarrow for the same Q , $dT_P < dT_V$, thus $c_P > c_V$.

8.7 Isothermal Compression (Slow)

$$P_1 V_1 = P_2 V_2 \quad (191)$$

$$W = Nk \ln(V_i/V_f), \quad W = - \int_{V_i}^{V_f} P dV \quad (192)$$

$$\Delta U = 0 \text{ since } \Delta T = 0, \quad \Delta U = \frac{f}{2} Nk \Delta T \quad (193)$$

8.8 Adiabatic Compression (Fast)

If no heat flows,

$$\Delta Q = 0 \rightarrow \Delta U = W \quad (194)$$

Equipartition Theorem,

$$\Delta U = Nk \Delta T = W \quad (195)$$

$$V_f T_f^{f/2} = V_i T_i^{f/2}, \quad f = \text{Degrees of Freedom} \quad (196)$$

$$V_f^\gamma P_f = V_i^\gamma P_i, \quad \gamma = \frac{f+2}{f} \quad (197)$$

$$W = \frac{P_f V_f - P_i V_i}{1 - \gamma} \quad (198)$$

$$PV^\gamma = C \rightarrow P = \frac{C}{V^\gamma} \quad (199)$$

$$W = \int P dV = C \frac{dV}{V^\gamma} = \frac{1}{\gamma} \frac{C}{V^{\gamma-1}} \Big|_{V_1}^{V_2} \quad (200)$$

8.9 Heat

$$Q = TdS \quad (201)$$

$$= mc\Delta T \quad (202)$$

$$= \text{Power} \cdot t \quad (203)$$

8.10 Multiplicity/States

$$\text{Probability}(\Omega_n) = \Omega(n)/\Omega(\text{all}) \quad (204)$$

1. Ω = multiplicity = how many different microstates yield a macrostate
2. Total number of macrostates = ($\#$ states thing can be in)^($\#$ of things)
3. e.g., 3 coins $\rightarrow 2^3 = 8 = \Omega$
4. $\#$ of ways to choose n things from N : $\Omega\binom{N}{n} = \frac{N!}{(N-n)!n!}$

8.11 Boltzmann Statistics

$$P(s) = \frac{g_s e^{E_s/kT}}{Z}, \quad Z = \sum_i g_i e^{-E_i/kT}, \quad g_i = \text{degeneracy of } i \quad (205)$$

$$P(A)/P(B) = \frac{g_a}{g_b} e^{(-A+B)/kT} \quad (206)$$

$$\langle \bar{x} \rangle = \frac{1}{Z} \sum_s x_s e^{-E_s/kT} \quad \text{average of any value} \quad (207)$$

$$\langle \bar{E} \rangle = \frac{1}{Z} \sum_i E_i e^{-E_i/kT} \quad (208)$$

$$U = N\bar{E} : \text{total energy of the system} \quad (209)$$

8.12 Density of State Distributions

Fermions:

$$N_i = \frac{g_i}{e^{(E_i - \mu)/kT} + 1} \quad (210)$$

Mesons/Bosons:

$$N_i = \frac{g_i}{e^{[E_i - \mu]/kT} - 1} \quad (211)$$

Boltzmann:

$$N_i = g_i e^{(E_i - \mu)/kT} - 1 \quad (212)$$

8.13 Blackbody Radiation

8.13.1 Wein's Law

$$T \cdot \lambda_{\max} = 3 \text{ mm} \cdot K \quad (213)$$

8.13.2 Stephan-Boltzmann

$$P \propto aT^4 \quad (214)$$

8.14 Heat Engines

$$e \leq 1 - \frac{T_c}{T_h} \quad (215)$$

$$e = \frac{\text{benefit}}{\text{cost}} = \frac{W}{Q_h}, W = Q_h - Q_c \quad (216)$$

8.15 Refrigerators

$$e \leq \frac{T_c}{T_h - T_c} \quad (217)$$

$$W = Q_h - Q_c \quad (218)$$

$$\Delta S = 0, \text{ independent of working substances} \quad (219)$$

8.16 Big People

1. Onnes: Superconductivity in Hg
2. Anderson: Positron
3. Yukawa: Strong Nuclear
4. Fermi: First nuclear reactor
5. Mann + Zweig: Quarks
6. Rontgen: X-rays
7. Penzias & Wilson: Background Radiation
8. Huygens: Wavefronts
9. Cavendish: G
10. Oersted: Connection between E&M
11. Ampere: B force law
12. Hertz: Showed E&M waves existed

9 Relativity

9.1 Space-Time Diagram

$\Delta S > 0$ Spacelike

1. Ordering of events depends on reference frame
2. There exists a reference frame where 2 events occur simultaneously, but they can't occur at the same location in space

$\Delta S < 0$ Timelike

1. Ordering of events is absolute
2. Casual relationships are timelike
3. Two events can occur at same point in space

9.2 Special Relativity

v/c	γ
.1	1.005
.25	1.033
.5	1.151
.75	1.55
.9	2.29

$$x = \gamma(x' + vt')$$
 (220)

$$t = \gamma\left(t' + \frac{vx}{c^2}\right)$$
 (221)

$$u'_x = \frac{u_x + v}{1 + \frac{u_x v}{c^2}}$$
 (222)

$$u'_z = \frac{u_z}{\gamma\left(1 + \frac{u_x v}{c^2}\right)}$$
 (223)

9.2.1 Time Dilation

$$t' = \gamma t_0, t_0 = \text{rest time}$$
 (224)

9.2.2 Length Contraction

$$x' = \frac{x_0}{\gamma}, x_0 = \text{rest length}$$
 (225)

9.2.3 Invariant Interval

$$\Delta s^2 = \Delta x^2 - (ct)^2 \leftarrow \text{transform between 2 moving frames}$$
 (226)

9.2.4 Energy

$$E_{\text{rel}} = \gamma E_0$$
 (227)

$$p = \gamma p = \gamma mv$$
 (228)

$$E_{\text{rel}}^2 = E_0^2 + (pc)^2$$
 (229)

$$E_{\text{rel}} \neq \frac{p_{\text{rel}}^2}{2m}$$
 (230)

$$p_x = \gamma\left(p_{x'} + \frac{v}{c^2}E'\right)$$
 (231)

$$E = \gamma(E' + vp_{x'})$$
 (232)

Last 2 lines employ the invariant 4-vector, where $p_{y'} = p_y$.

10 Atomic Physics

10.1 Hydrogen Spectral Series

$$\frac{1}{\lambda} = R_y \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right), \quad R_y \approx 1 \times 10^7 \text{m}^{-1}$$
 (233)

1. Lyman: $n_f = 1$
2. Balmer: $n_f = 2$
3. Paschen: $n_f = 3$

$$\Delta E = E_0 \left(\frac{1}{n_f^2} - \frac{1}{n_i^2} \right) \quad (234)$$

10.2 Atomic Notation

$${}_Z^AX \quad (235)$$

1. A = mass number = $p^+ + n^0$
2. Z = number of protons = chemical number

11 Particle Physics

11.1 Fermi

$$E_F = k_b T_F \quad (236)$$

$$p_F = \hbar k_F \rightarrow E_F = \frac{p^2}{2mn} = \frac{\hbar^2 k^2}{2m}, \quad v_F = \frac{p_F}{m} \quad (237)$$

$$k_F \left(\frac{3\pi^2 N}{\text{volume}} \right)^{1/3}, \quad p_F = \frac{2}{3} \frac{E_F}{v} \quad (238)$$

Degenerate Fermi gas, so cold that nearly all states below E_F are occupied and above states are unoccupied.

11.2 Degeneracy Pressure of a Solid

$P = \frac{3}{2} \frac{E}{V}$: The stabilizing internal pressure that comes from the anti-symmetrization requirement for the wave functions of identical fermions.

12 Misc

12.1 Water Density

$$1 \text{ liter} = 1 \text{ kg}, \rho = 1 \text{ g/cm}^3 \quad (239)$$

Beats occur when f_1 and f_2 are close together

12.2 Fundamental Law of Statistical Mechanics

All accessible microstates are equally likely

12.3 Irreversible Process

Creates new entropy

12.4 Reversible Process

Creates no new entropy