

Forestry in the Pacific Northwest: How important are riparian buffers to stream-associated amphibians?

The *Western Forester* has asked Washington Wildlife Biologists Marc Hayes and A.J. Kroll to provide their perspectives on the importance of riparian buffers to stream-associated amphibians.

Both of their perspectives were printed in the March/April/May 2010 issue of the *Western Forester*. Below is Literature Cited for both articles.

A.J. Kroll Literature Cited

- Brattstrom, B. H. 1963. A preliminary review of the thermal requirements of amphibians. *Ecology* 44:238-255.
- Brosofske, K. D., J. Chen, R. J. Naiman, and J. F. Franklin. 1997. Harvesting effects on microclimatic gradients from small streams to uplands in western Washington. *Ecological Applications* 7:1188-1200.
- Claussen, D. L. 1973. The thermal relations of the tailed frog, *Ascaphus truei* and the Pacific tree frog, *Hyla regilla*. *Comparative Biochemistry and Physiology* 44A:137-153.
- Corn, P. S. and R. B. Bury. 1989. Logging in western Oregon: responses of headwater habitats and stream amphibians. *Forest Ecology and Management* 29:39-57.
- Diller, L. V. and R. L. Wallace. 1996. Distribution and habitat of *Rhyacotriton variegatus* in managed, young growth forests in north coastal California. *Journal of Herpetology* 30:184-191.
- Dupuis, L. and D. Steventon. 1999. Riparian management and the tailed frog in northern coastal forests. *Forest Ecology and Management* 124:35-43.
- Hawkins, C. P., M. L. Murphy, N. H. Anderson, and M. A. Wilzbach. 1983. Density of fish and salamanders in relation to riparian canopy and physical habitat in streams of the northwestern United States. *Canadian Journal of Fisheries and Aquatic Sciences* 40:1173-1185.
- Huff, D. D., S. L. Hubler, and A. N. Borisenko. 2005. Using field data to estimate the realized thermal niche of aquatic vertebrates. *North American Journal of Fisheries Management* 25:346-360.
- Jackson, C. R., D. P. Batzer, S. S. Cross, S. M. Haggerty, and C. A. Sturm. 2007. Headwater streams and timber harvest: channel, macroinvertebrate, and amphibian response and recovery. *Forest Science* 53:356-370.

- Kiffney, P. M., J. S. Richardson, and J. P. Bull. 2004. Establishing light as a causal mechanism structuring stream communities in response to experimental manipulation of riparian buffer width. *Journal of the North American Benthological Society* 23:542-555.
- Mallory, M. A. and J. S. Richardson. 2005. Complex interactions of light, nutrients, and consumer density in a stream periphyton-grazer (tailed frog tadpoles) system. *Journal of Animal Ecology* 74:1020-1028.
- Murphy, M. L. and J. D. Hall. 1981. Varied effects of clear-cut logging on predators and their habitat in small streams of the Cascade Mountains, Oregon. *Canadian Journal of Fisheries and Aquatic Science* 38:137-145.
- Stoddard, M. A. and J. P. Hayes. 2005. The influence of forest management on headwater stream amphibians at multiple spatial scales. *Ecological Applications* 15:811-823.
- Vesely, D. G. and W. C. McComb. 2002. Salamander abundance and amphibian species richness in riparian buffer strips in the Oregon Coast Range. *Forest Science* 48:291-297.
- Welsh, H. H., Jr. and A. J. Lind. 1996. Habitat correlates of the southern torrent salamander, *Rhyacotriton variegatus* (Caudata:Rhyacotritonidae), in northwestern California. *Journal of Herpetology* 30:385-398.
- Welsh, H. H., Jr. and L. M. Ollivier. 1998. Steam amphibians as indicators of ecosystem stress: a case study from California's redwoods. *Ecological Applications* 8:1118-1132.
- Wilkins, R. N. and N. P. Peterson. 2000. Factors related to amphibian occurrence and abundance in headwater streams draining second-growth Douglas fir forests in southwestern Washington. *Forest Ecology and Management* 139:79-91.

MARC P. HAYES LITERATURE CITED

- BROWN, H.A. 1975. Temperature and development of the tailed frog, *Ascaphus truei*. *Comparative Biochemistry and Physiology* 50A:397-405.
- BURY, R.B. AND P.S. CORN. 1988. Responses of aquatic and streamside amphibians to timber harvest: a review. Pp. 165-181. *In*: Raedeke, K.J. (editor), *Streamside management: riparian wildlife and forestry interactions: Proceedings of a symposium*; 1987 Feb. 11-13; Institute of Forest Resources, University of Washington, Seattle, Contribution 59.
- CORN, P.S. AND R.B. BURY. 1989. Logging in western Oregon: responses of headwater habitats and stream amphibians. *Forest Ecology and Management* 29(1):39-57.

- HAWKINS, C.P., J. FEMINELLA, AND C.M. CRISAFULLI. 1997. Responses and recovery of stream-dwelling tailed frog and giant salamander populations following the 1980 eruption of Mount St. Helens. *In*: Richardson, J.S. and T.R. Wahbe (editors). Proceedings of the Tailed Frog Workshop. April 14-17, 1997. University of British Columbia. Vancouver, British Columbia, Canada. 49 p.
- HAWKINS, C.P., L.J. GOTTSCHALK, AND S.S. BROWN. 1988. Densities and habitat of Tailed Frog tadpoles in small streams near Mt. St. Helens following the 1980 eruption. *Journal of the North American Benthological Society* 7(3):246-252.
- KIFFNEY, P.M., AND J.S. RICHARDSON. 2001. Interactions among nutrients, periphyton, and invertebrates and vertebrate (*Ascaphus truei*) grazers in experimental channels. *Copeia* 2001(2):422-429.
- KREUTZWEISER, D.P., AND S.S. CAPELL. 2001. Fine sediment deposition in streams after selective forest harvesting without riparian buffers. *Canadian Journal of Forest Research* 31(12): 2134–2142.
- KROLL, A.J., K. RISENHOOVER, T. MCBRIDE, E. BEACH, B.J. KERNOHAN, J. LIGHT, AND J. BACH. 2007. Factors influencing stream occupancy and detection probability parameters of stream-associated amphibians in commercial forests of Oregon and Washington. *Forest Ecology and Management* 255:3726-3735.
- LISLE, T.E. AND S. HILTON. 1999. Fine bed material in pools of natural gravel bed channels. *Water Resources Research*. (35)4:1291-1304.