

# WILDLIFE RESPONSES TO FOREST STRUCTURAL RETENTION



Jake Verschuyl

Biodiversity Research Coordinator  
National Council for Air and Stream Improvement  
Anacortes, WA

# Forest Practices Rules require retention



Green Trees



Snags



Downed wood

Requirements per harvested acre:

- 2 snags or green trees  $\geq 30$  ft tall and  $\geq 11$  in. DBH
- 2 down logs  $\geq 6$  ft long with a total volume of  $\geq 10$  ft<sup>3</sup>; logs  $\geq 20$  ft<sup>3</sup> count as 2 logs



# Structural Retention



# Recent Research Topics

- Structural retention
  - What is out there on the landscape?
  - Bird structure use and response to patch size
- Effects of environmental gradients on structure use
  - Bird response to snag density and down wood across a productivity gradient

- Structural enrichment



Dr. A.J. Kroll



Dr. Dan Linden



Dr. Gary Roloff



# Structural Retention

## *Response of avian species richness to operational practices*



- California ( $n=23$ ), Oregon ( $n=20$ ), and Washington ( $n=42$ ); 2008-09
- Stand size (ha)
  - CA, 9 (4-11)
  - OR, 40 (22-61)
  - CWA, 8 (4-12)
  - SWA, 37 (24-57)
- Structural retention
  - Number of green trees
  - Number of snags
  - Distance to edge

# Objectives



- 1) Quantify the physical characteristics of retained structures found in clearcut harvest units



- 2) Summarize stand-level bird communities that use clearcut harvest units

# Objectives




- 1) Quantify the physical characteristics of retained structures found in clearcut harvest units



- 2) Summarize stand-level bird communities that use clearcut harvest units

# Selecting Structures for Sampling

Photo Mapping



Field Reconnaissance




Sample





# Types of Structures



Green Trees



Snags



Downed Wood



Single



Patches

## Quantity

- Patches (includes green trees and snags)\*
  - Range = 0.25 to 4.9 patches/ac (0.1 to 2.0 patches/ha)
  - Average size = 0.75 ac (range = 0.25 to 10.9 ac)  
(0.3 ha [range = 0.1 to 4.4 ha])

\* Only includes patches isolated from adjacent forest and riparian zones

## Quantity (Stand-level)

- Patches (includes green trees and snags)\*
  - Range = 0.25 to 4.9 patches/ac (0.1 to 2.0 patches/ha)
  - Average size = 0.75 ac (range = 0.25 to 10.9 ac)  
(0.3 ha [range = 0.1 to 4.4 ha])
- Single Green Trees
  - Range = 0.25 to 2.5 trees/ha (0.1 – 1.0 trees/ha)




# Quantity


- Patches (includes green trees and snags)\*
  - Range = 0.25 to 4.9 patches/ac (0.1 to 2.0 patches/ha)
  - Average size = 0.75 ac (range = 0.25 to 10.9 ac)  
(0.3 ha [range = 0.1 to 4.4 ha])
- Single Green Trees
  - Range = 0.25 to 2.5 trees/ha (0.1 – 1.0 trees/ha)
- Snags
  - Range = 0.25 to 1.7 snags/ac (0.1 – 0.7 snags/ha)

# Green Trees


**NCA**




**SOR**



**SWA**




**CWA**




# Snags


**NCA**




**SOR**



**SWA**




**CWA**





# Downed Wood



1: 10 – 25 cm, 2: 26 – 51 cm, 3: >51 cm

# Objective 1 - Summary

- Structures do not occur at high densities
- Trees and snags encompass a range of sizes
  - 10 to >125 cm (4 to >49 in) dbh, with majority <51 cm (< 20 in)
- Subtle differences between patches and single structures
- Legacy structures present
- Abundant downed wood

# Objectives



- 1) Quantify the physical characteristics of retained structures found in clearcut harvest units



- 2) Summarize stand-level bird communities that use clearcut harvest units




# Analytical Methods



- Estimate species richness at *individual structures* or *patches of structures* within each harvest unit

# Results:

Positive relationship between patch size and species richness



## Objective 2 - Summary

- Large number of bird species
- Detection for bird community is complex
- Occupancy for region = no clear pattern with retention
- Within study area occupancy can be linked to retention attributes
- Recent clearcuts are being used by guilds not typically associated with clearcutting



# Does energy mediate the relationship between avian functional diversity and within stand structure?

**Objective:** Investigate the role of available energy in determining the relationship between bird richness and abundance and elements of forest structure and vegetation cover at stand scales.



- Large snag density, coarse wood volume, canopy cover
- Response data: Species richness and species' abundance

# Does energy mediate the relationship between avian functional diversity and within stand structure?






# Does energy mediate the relationship between avian functional diversity and within stand structure?




# Snags and Coarse Woody Debris

Abundance of  
primary guilds  
by snag density  
and CWD  
volume

Springfield




Cle Elum







# Canopy Cover


Species richness  
and abundance  
by canopy cover



Springfield



Cle Elum



# Management Implications

- Avian species respond to variation in density of snags and volume of coarse wood differently on the west and east slopes of the Cascades
- Responses of species' abundance were more defined than species richness in both models
- Canopy closure reduces the abundance of bird species, especially in high energy landscapes
- Many other factors are likely in play

# Structural Enrichment






# Creating Snags

## Created Snags Project

- Initiated 1997→1999
- 1123 snags created & monitored
- 5 year revisit / 10 year revisit complete: 20 year revisit planned
- Monitoring use by wildlife
- Examining how spatial distribution & landscape context influences use











# Methods


## *Experimental Design and Analysis*

- Ground-based units, primarily Douglas-fir
- Random assignment to one of 6 treatments: dispersion  $\times$  density
- ~1,200 snags were created with a harvester on 28 harvest units
- Landscape was characterized in a 1000 m buffer around each stand
- Dispersion as median NN distance

Density	Clumped created snags	Dispersed created snags
<i>Low</i> (0.5 created snag/ha)		
<i>Medium</i> (1 created snag/ha)		
<i>High</i> (2 created snags/ha)		

# Does enrichment benefit wildlife?

## *Demographic responses*




- **CBCH**
  - <0.50; control and partial harvest; Mahon and Martin (2006)
  - <0.50 ; buffer strips <250 m or > 350 m; Sperry et al. (2008)
- **HOWR**
  - 0.63 in natural cavities to 0.83 in nest boxes; Finch, 1989, 1990; Li and Martin, 1991; Johnson and Kermott, 1994; Purcell et al., 1997
- **NOFL**
  - 0.65 in both unlogged burned forests and partially logged burned forests; Saab et al. 2007
  - 0.50 (low severity burn) and 1.0 (high severity burn); Vierling et al. 2008
  - 0.65 in an unmanaged forest; Fisher and Wiebe (2007)
  - 0.41 in unburned forests and 0.80 in burned forests; Kozma and Kroll (2012)


# How does landscape composition interact with local structural enrichment?

## *Habitat surplus and habitat source hypotheses*

*Habitat surplus hypothesis*



*Habitat source hypothesis*




Percent of mature forest in landscape


# Results

## *Support for the habitat surplus hypothesis*

### Early successional (ES)



### Mature forest (MF)



Percent of surrounding landscape (1 km radius) in mature forest (> 40 years of age)



# Future Research Topics

What other features (e.g., remnant green trees) are required before species will use snags in clearcut stands?

Green tree longevity


How big must retention patches be to promote nesting/roosting occupancy of cavity-dependent species?

Does the proportion of older forest (>80 years old) in the landscape influence the use of snags in harvest units?

# Evaluating Alternative Retention/Enrichment Practices

*NCASI and Michigan State University*

- Bird and bat responses
- Control for landscape structure
- Retention and enrichment
- Reduce uniformity



Current  
operational

The diagram shows a dense cluster of green tree icons of varying heights, all pointing upwards, representing a uniform forest structure.



Treatment 1

The diagram shows a dense cluster of green tree icons of varying heights, all pointing upwards, representing a uniform forest structure.



Treatment 2

The diagram shows two distinct groups of green tree icons, each with varying heights and pointing upwards, representing a non-uniform forest structure.



Treatment 4

The diagram shows four distinct groups of green tree icons, each with varying heights and pointing upwards, representing a non-uniform forest structure.



Treatment 3

The diagram shows two distinct groups of green tree icons, each with varying heights and pointing upwards, representing a non-uniform forest structure.

# Study Objectives

1. Determine the degree to which use of green trees and created snags by birds varies with aggregation and slope position of retained structures.

2.



use of  
regat  
create

3.

aggreg  
cavity



ures

“Responses of bat populations to various forest management practices... and habitat variables... are virtually unknown.”



“Conservation of bat species... should center on preserving suitable roost sites and foraging areas, but attaining this goal is hampered by our rudimentary knowledge of these requirements.”

Christy and West 1993



# Factors Influencing Patterns of Habitat Use by Bats



Amount of clutter

Roost Sites



Use and distribution of water



# The Future

## *Reorganization of the Forest Estate*

- Increased upslope retention
  - Modification of unstable slope buffers
- Reduced riparian buffers
  - Reflect regional variation
- Monetization of other values
  - Ecosystem services, including carbon storage and water
- Landscape planning
  - Across jurisdictions



# Partners

- AJ Kroll, Weyerhaeuser
- Gary Roloff, Michigan State University
- Dan Linden, MSU and USGS
- Ed Arnett, Theodore Roosevelt Conservation Partnership
- Mike Lacki, University of Kentucky
- Matt Betts, Oregon State University
- Steve Duke, Matt Hane, Josh Johnson, & Mike Rochelle, Weyerhaeuser
- Oregon Forest Industry Council
- NCASI Western Wildlife Task Group

