




QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

FERROUS SULFATE

DRUG CLASS: Oral iron supplement



✓ MAIN USES



Prevention and treatment of iron-deficiency anaemia

⚠ COMMON SIDE EFFECTS



Constipation



Nausea



Abdominal discomfort



Dark stools

⚠ IMPORTANT WARNINGS



Iron overdose can be extremely dangerous, particularly in children.

★ KEY POINT

Store iron products safely and use only as directed.



🛡 PRECAUTIONS



Keep iron products securely away from children.



Other medicines and foods can interfere with absorption.



Iron should ideally be used when deficiency or increased need is appropriately established.

★ SEEK MEDICAL HELP IF



Suspected overdose



Severe abdominal symptoms



Vomiting blood



ADDITIONAL INFORMATION



Take on an empty stomach or as advised for best absorption.



Follow the dose, timing and duration as recommended.



Do not self-medicate or use for longer than recommended.



KEY POINT

Ferrous sulfate treats iron deficiency. Do not use for non-iron deficiencies.

CALCIUM + VITAMIN D

DRUG CLASS: Mineral/vitamin supplementation



✓ MAIN USES



Prevention or treatment of calcium/vitamin-D deficiency



Support of bone health when supplementation is indicated

⚠ COMMON SIDE EFFECTS



Constipation



Bloating



Nausea

⚠ IMPORTANT WARNINGS



Excessive supplementation can result in high blood calcium and kidney complications.

★ KEY POINT

Calcium and Vitamin D work together to support strong bones and overall health.



🛡 PRECAUTIONS



Use carefully with kidney stones, kidney disease or existing high calcium.



Calcium can interfere with absorption of several medicines.

★ SEEK MEDICAL HELP IF



Confusion



Severe weakness



Persistent vomiting



Abnormal heartbeat



ADDITIONAL INFORMATION



Take with or after food as advised.



Take regularly at the same time for best results.



Do not self-medicate or use for longer than recommended.



KEY POINT

Use only as directed by a healthcare professional.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY



Not a Prescription

ORAL REHYDRATION SALTS (ORS)

CATEGORY:

Glucose-electrolyte replacement solution




MAIN USES



Prevention and treatment of dehydration caused by diarrhoea and other appropriate fluid-loss states.



IMPORTANT NOTE



ORS replaces water and electrolytes; **it does not** directly stop diarrhoea or eliminate the underlying infection.



PRECAUTIONS



Prepare exactly according to package instructions.



Incorrect dilution can be dangerous.



Severe dehydration requires urgent medical treatment.



SEEK MEDICAL HELP IF



Inability to drink



Persistent vomiting



Marked lethargy



Very little urine



Signs of severe dehydration



ADDITIONAL INFORMATION



Drink small, frequent amounts of ORS.



Continue feeding normally for infants and children.



Store prepared ORS properly and use within 24 hours.



ORS is safe, effective and recommended worldwide for the management of dehydration.

DEXTROSE IV SOLUTIONS

CATEGORY:

Intravenous fluid/glucose preparation



EDUCATIONAL REFERENCE ONLY



Not a Prescription

MAIN USES



Specific fluid, glucose or nutritional indications in medical settings depending on concentration.



IMPORTANT WARNINGS



- Different concentrations such as 5% and 25% are **not interchangeable**.
- Improper administration can cause serious glucose, electrolyte and fluid disturbances.



PRECAUTIONS



Intravenous dextrose should be selected and administered by trained healthcare professionals.



ADDITIONAL INFORMATION



- Concentration, rate and total volume depend on patient condition.



- Monitor blood glucose, electrolytes and fluid status as indicated.



- Use appropriate sterile technique.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine or IV preparation.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

ALPRAZOLAM

DRUG CLASS: Benzodiazepine



✓ MAIN USES



Anxiety disorders



Panic disorder

⚠ COMMON SIDE EFFECTS



Drowsiness



Dizziness



Tiredness



Problems with coordination or concentration

⚠ IMPORTANT WARNINGS



Can cause dependence and withdrawal.



Abrupt discontinuation after regular use can cause serious withdrawal symptoms, including seizures.



Combining benzodiazepines with opioids, alcohol or other powerful sedatives can cause dangerous respiratory depression.

★ KEY POINT

Use only as directed by a healthcare professional for the shortest time required.



🛡 PRECAUTIONS



Use only under medical supervision.



Do not stop regular treatment suddenly without professional guidance.



Avoid driving when impaired.

★ SEEK MEDICAL HELP IF



Very slow breathing



Extreme sedation



Seizure



Severe confusion

i ADDITIONAL INFORMATION



Take only as prescribed. Do not share this medicine with others.



Take at the same time each day as advised for best results.



Do not self-medicate or use for longer than recommended.

💡 KEY POINT

Alprazolam works by calming excessive brain activity and reducing anxiety.



FOLIC ACID

DRUG CLASS: Vitamin B9



✓ MAIN USES



Prevention and treatment of folate deficiency



Certain types of megaloblastic anaemia caused by folate deficiency



Recommended supplementation around pregnancy according to clinical guidance

⚠ COMMON SIDE EFFECTS



Usually well tolerated



Occasional gastrointestinal discomfort

⚠ IMPORTANT WARNINGS



High-dose folic acid can mask some signs of vitamin-B12 deficiency while neurological injury progresses.

★ KEY POINT

Folic acid is an essential nutrient that supports healthy cell growth and red blood cell production.



🛡 PRECAUTIONS



The cause of unexplained anaemia should be appropriately assessed.

★ SEEK MEDICAL HELP IF



Significant allergic reaction

i ADDITIONAL INFORMATION



Take with or without food, as advised.



Adequate intake of balanced diet is important.



Do not self-medicate. Use only as directed by a healthcare professional.

💡 KEY POINT

Folic acid helps prevent deficiency-related anaemia and supports overall health.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

BETAHISTINE

DRUG CLASS: Histamine analogue /
antivertigo medicine



✓ MAIN USES



Symptoms associated with Ménière's disease, such as vertigo

⚠ COMMON SIDE EFFECTS



Headache



Nausea



Indigestion

⚠ IMPORTANT WARNINGS



It does not treat every possible cause of dizziness.

★ KEY POINT

Betahistine helps relieve symptoms associated with Ménière's disease.



🛡 PRECAUTIONS



Vertigo should be appropriately diagnosed.



Use cautiously in people with peptic-ulcer history or asthma when advised by a clinician.

★ SEEK MEDICAL HELP IF



Severe allergic reaction



New neurological symptoms associated with dizziness

i ADDITIONAL INFORMATION



Take with food if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.

💡 KEY POINT

It does not treat every possible cause of dizziness. Use only as advised.

CINNARIZINE

DRUG CLASS: Antihistamine with
antivertigo properties



✓ MAIN USES



Motion sickness



Vertigo and vestibular symptoms in some countries

⚠ COMMON SIDE EFFECTS



Drowsiness



Weight gain



Indigestion

⚠ IMPORTANT WARNINGS



Can impair alertness.



Prolonged use may produce movement-related adverse effects, especially in older adults.

★ KEY POINT

Use caution when driving and avoid alcohol when it increases sedation.



🛡 PRECAUTIONS



Avoid alcohol when it increases sedation.



Use caution when driving.



Parkinson's disease requires particular caution.

★ SEEK MEDICAL HELP IF



New tremor



Muscle stiffness



Abnormal movements



Severe allergic reaction

i ADDITIONAL INFORMATION



Take with food if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.

💡 KEY POINT

Cinnarizine helps relieve motion sickness and vertigo symptoms in selected cases.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY



Not a Prescription

CIPROFLOXACIN – SYSTEMIC

DRUG CLASS:

Fluoroquinolone antibiotic



MAIN USES

Selected susceptible bacterial infections when ciprofloxacin is appropriate.



COMMON SIDE EFFECTS



Nausea



Diarrhoea



Headache



IMPORTANT WARNINGS



Systemic fluoroquinolones can cause serious tendon, nerve and central-nervous-system adverse effects.



Tendon rupture can occur.



They should not be used casually for uncomplicated illnesses when safer alternatives are appropriate.



PRECAUTIONS



Tell a clinician about tendon disorders, neurological disease, myasthenia gravis and interacting medicines.



SEEK MEDICAL HELP IF



Tendon pain/swelling



New numbness or burning



Severe mental/neurological symptoms



Serious allergic reaction



ADDITIONAL INFORMATION



- Take exactly as prescribed.
- Complete the full course even if you feel better.



- Appropriate dose and duration depend on the infection.



- Avoid unnecessary use to reduce risk of resistance and harm.

OFLOXACIN + ORNIDAZOLE

DRUG CLASS:

Fluoroquinolone antibiotic + nitroimidazole antimicrobial



MAIN USES

Combination products exist in some markets for selected mixed bacterial/protozoal gastrointestinal or other infections.



IMPORTANT WARNING



A fixed combination should **not** be described as a general medicine for diarrhoea.



Most acute diarrhoeal illnesses do not automatically require antibiotics.



PRECAUTIONS



Appropriate diagnosis is important.



Fluoroquinolone and nitroimidazole warnings both apply.



Prescription supervision is appropriate.



ADDITIONAL INFORMATION



Use only when prescribed by a healthcare professional.



Complete the prescribed course.



Do not share antibiotics with others.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE


EDUCATIONAL REFERENCE ONLY




Not a Prescription

GENTAMICIN INJECTION

DRUG CLASS: Aminoglycoside antibiotic



✓ MAIN USES



Serious susceptible bacterial infections.

⚠ IMPORTANT WARNINGS



Kidney toxicity



Hearing/balance toxicity



Neuromuscular toxicity

🛡 PRECAUTIONS



Professional dosing and monitoring are necessary.



Kidney function is an important determinant of therapy.

★ SEEK MEDICAL HELP IF



Hearing loss



Vertigo



Markedly reduced urination



Muscle weakness

i ADDITIONAL INFORMATION



Therapy usually requires drug level monitoring.



Avoid other kidney or ear-toxic medicines when possible.



Duration of therapy is kept as short as appropriately possible.



Report any unusual symptoms promptly to your healthcare professional.

CEFTRIAXONE INJECTION


DRUG CLASS: Third-generation cephalosporin antibiotic




EDUCATIONAL REFERENCE ONLY



Not a Prescription



✓ MAIN USES



Numerous serious susceptible bacterial infections.

⚠ COMMON SIDE EFFECTS



Injection-site pain



Diarrhoea



Rash

⚠ IMPORTANT WARNINGS



Severe allergic reactions and antibiotic-associated colitis are possible.



Special precautions apply to neonates, particularly around calcium-containing intravenous solutions.

🛡 PRECAUTIONS



Administer under professional care.



Severe beta-lactam allergy should be reviewed.

i ADDITIONAL INFORMATION



Given by injection (I.M. or I.V.) as prescribed.



Use only when clearly needed for bacterial infections.



Report persistent or bloody diarrhoea.



Appropriate cultures and sensitivity testing help guide therapy.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



TRAMADOL INJECTION

DRUG CLASS: Opioid analgesic



MAIN USES



Moderate-to-severe pain where parenteral opioid therapy is clinically appropriate.

COMMON SIDE EFFECTS



Nausea



Dizziness



Sedation



Vomiting

IMPORTANT WARNINGS



Respiratory depression



Dependence



Seizures



Serotonin syndrome

PRECAUTIONS



Should be administered under professional supervision.

ADDITIONAL INFORMATION



Use the lowest effective dose for the shortest necessary duration.



Avoid alcohol and other CNS depressants (e.g., benzodiazepines).




Do not drive or operate machinery when sedated.




Keep out of reach of children.

AMIKACIN INJECTION

DRUG CLASS: Aminoglycoside antibiotic



MAIN USES



Serious susceptible bacterial infections.

COMMON SIDE EFFECTS



Injection-related effects may occur.

(e.g., pain, redness, swelling at injection site)

IMPORTANT WARNINGS



Can cause serious kidney toxicity and hearing/balance damage.



Neuromuscular blockade is another important risk.



Use with caution in patients with existing kidney problems.

PRECAUTIONS



Kidney function and drug exposure may require monitoring.



Other nephrotoxic or ototoxic medicines increase risk.

SEEK MEDICAL HELP IF



Hearing changes



Ringing in ears



Balance problems



Reduced urine output (less urination)



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

VITAMIN E

DRUG CLASS: Fat-soluble vitamin



MAIN USES



Prevention or treatment of vitamin-E deficiency when medically indicated.



COMMON SIDE EFFECTS



Usually well tolerated at appropriate amounts.



Higher doses may cause gastrointestinal symptoms.



IMPORTANT WARNINGS



High-dose supplementation may increase bleeding risk.



KEY POINT

Use only as directed and at the recommended dose.



PRECAUTIONS



Discuss significant supplementation with a clinician if taking anticoagulant or antiplatelet medicines.



SEEK MEDICAL HELP IF



- Unusual bleeding
- Serious allergic reaction



QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

VITAMIN C

DRUG CLASS: Water-soluble vitamin



MAIN USES



Prevention and treatment of vitamin-C deficiency, including scurvy.



COMMON SIDE EFFECTS



Nausea



Abdominal cramps



Diarrhoea at higher amounts.



IMPORTANT WARNINGS



Very high supplemental intake may contribute to kidney-stone formation in susceptible people.



KEY POINT

Use only as directed and at the recommended dose.



PRECAUTIONS



Extra caution with recurrent kidney stones or certain iron-overload conditions.



SEEK MEDICAL HELP IF



Unusual bleeding



Serious allergic reaction



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY




Not a Prescription

DICYCLOMINE INJECTION

DRUG CLASS: Anticholinergic antispasmodic



MAIN USES



Injectable treatment of gastrointestinal spasm in limited circumstances where appropriate.



IMPORTANT WARNINGS



Anticholinergic adverse effects can be significant.



Injectable administration requires professional assessment.



KEY POINT

Use only as directed by a healthcare professional.



POSSIBLE ANTICHOLINERGIC EFFECTS



Dry mouth



Blurred vision



Fast heartbeat



Confusion



Difficulty urinating



Flushing



ADDITIONAL INFORMATION



Administration must be under the care of appropriately trained healthcare personnel based on clinical need.



Not for self-administration.
Not a treatment for the cause of abdominal pain.

DROTAVERINE

DRUG CLASS: Antispasmodic



MAIN USES



Smooth-muscle spasm in countries where it is approved.



COMMON SIDE EFFECTS



Nausea



Dizziness



Headache



Possible lowering of blood pressure



IMPORTANT WARNINGS



Evidence, approved indications and formulations vary by country.



PRECAUTIONS



Use should follow local product information rather than a universal dosing recommendation.



ADDITIONAL INFORMATION



Availability, strengths and approved use may differ between countries.



Follow the product information and healthcare professional advice in your region.



Report any concerning symptoms to a healthcare professional.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

TETANUS VACCINE / TETANUS-CONTAINING VACCINE

DRUG CLASS: Vaccine



EDUCATIONAL REFERENCE ONLY



Not a Prescription



MAIN USES



- Prevention of tetanus as part of recommended immunisation.



- Used in wound-management protocols depending on vaccination history and wound type.



COMMON SIDE EFFECTS



Injection-site pain



Redness/swelling



Mild fever



Fatigue



IMPORTANT WARNINGS



This is a preventive vaccine, not a treatment for established tetanus.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Previous severe allergic reaction to a vaccine component requires medical evaluation.



SEEK MEDICAL HELP IF



Severe allergic reaction



Significant neurological symptoms



Stay up to date with your recommended vaccination schedule for ongoing protection.



ADDITIONAL INFORMATION



Booster doses are important to maintain protection.



For wounds, timing and need for vaccine depend on history and wound type.



Inform the healthcare professional about any past vaccine reactions or health conditions.



Vaccines help protect you and the community from tetanus.

ONDANSETRON INJECTION

DRUG CLASS: 5-HT₃ antagonist antiemetic



EDUCATIONAL REFERENCE ONLY



Not a Prescription



MAIN USES



Prevention/treatment of significant nausea and vomiting in medical settings.



COMMON SIDE EFFECTS



Headache



Constipation



Injection-site reaction



IMPORTANT WARNINGS



Cardiac QT prolongation is an important consideration.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Administered under medical supervision when injection is required.



SEEK MEDICAL HELP IF



Irregular heartbeat or fainting



Severe allergic reaction or significant neurological symptoms



Inform the healthcare team about all medicines you are taking and any heart conditions.



ADDITIONAL INFORMATION



Usually given by IV or IM route in clinical settings.



Works within minutes to help control nausea and vomiting.



Dose and frequency depend on the condition, patient factors and medical decision.



Keep all healthcare appointments and follow medical advice.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



PHENIRAMINE INJECTION

DRUG CLASS: First-generation antihistamine



MAIN USES



Allergic reactions in appropriate clinical circumstances.

COMMON SIDE EFFECTS



Sedation



Dizziness



Dry mouth



Blurred vision

IMPORTANT WARNINGS



Injectable antihistamines can cause significant sedation and cardiovascular effects.

KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Should be given by appropriately trained healthcare personnel.

SEEK MEDICAL HELP IF



Severe allergic reaction



Rapid or irregular heartbeat



Breathing difficulty

ADDITIONAL INFORMATION




May impair alertness.

Avoid driving or operating machinery after injection.

DEXAMETHASONE INJECTION

DRUG CLASS: Systemic corticosteroid



MAIN USES



Severe inflammatory or allergic conditions.



Various specialist indications.

COMMON SIDE EFFECTS



Increased blood glucose



Mood changes



Indigestion



Sleep disturbance

IMPORTANT WARNINGS



Infection and systemic steroid risks apply.

Use only when the benefits outweigh the risks.

KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Injectable corticosteroids require an appropriate clinical indication and professional administration.

SEEK MEDICAL HELP IF



Fever or signs of infection



Severe mood changes



Severe muscle weakness or pain

ADDITIONAL INFORMATION



Do not stop suddenly after prolonged use. Follow your doctor's advice for tapering if needed.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





LEVOCETIRIZINE

DRUG CLASS:

Second-generation antihistamine

**MAIN USES**

Allergic rhinitis



Sneezing and runny nose due to allergies



Itching and hives

**COMMON SIDE EFFECTS**

Sleepiness



Fatigue



Dry mouth

**IMPORTANT WARNINGS**

May cause drowsiness in some people.

**PRECAUTIONS**

Avoid driving until you know how it affects you.



Alcohol and other sedatives can increase drowsiness.



Dose adjustment may be necessary in significant kidney impairment.

**SEEK MEDICAL HELP IF**

Swelling of the face/throat



Difficulty breathing



Severe allergic reaction

**ADDITIONAL INFORMATION**

Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.

**KEY POINT**

Levocetirizine helps relieve allergy symptoms such as sneezing, runny nose, itching and hives. Use only as advised.



CETIRIZINE

DRUG CLASS:

Second-generation antihistamine

**MAIN USES**

Hay fever and other allergic rhinitis symptoms



Itching



Urticaria (hives)

**COMMON SIDE EFFECTS**

Drowsiness



Fatigue



Dry mouth



Headache

**IMPORTANT WARNINGS**

Although less sedating than older antihistamines, it can still cause sleepiness.

**PRECAUTIONS**

Use care when driving.



Alcohol or sedating medicines may increase drowsiness.



Kidney impairment may affect how the medicine is used.

**SEEK MEDICAL HELP IF**

Severe allergic reaction



Difficulty breathing or swallowing



Significant facial or throat swelling

**ADDITIONAL INFORMATION**

Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.

**KEY POINT**

Cetirizine helps relieve allergy symptoms like runny nose, itching and hives. Use only as advised.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

CIPROFLOXACIN OPHTHALMIC

DRUG CLASS: Fluoroquinolone
ophthalmic antibiotic



✓ MAIN USES



Susceptible
bacterial
eye infections.

⚠ COMMON SIDE EFFECTS



Temporary
eye irritation



Burning/stinging



Unpleasant taste

⚠ IMPORTANT WARNINGS



Antibiotic
eye drops
do not treat
viral or allergic
conjunctivitis.

★ KEY POINT

Use only as directed
by a healthcare
professional.



🛡 PRECAUTIONS



Avoid contaminating
the dropper tip.



Contact-lens use should
follow professional advice
during infection.

★ SEEK MEDICAL HELP IF



Severe eye pain



Vision deterioration



Significant swelling



Symptoms worsen
despite treatment



Complete the full
course even if you
feel better unless
advised otherwise.



Do not share
this medicine
with others.



ADDITIONAL INFORMATION



Wash hands before
and after use.



Use at evenly spaced
intervals for best results.



Store as directed on the label.
Do not freeze.

CARBOXYMETHYLCELLULOSE OPHTHALMIC

DRUG CLASS: Ocular lubricant /
Artificial tears



✓ MAIN USES



Relief of
dry-eye
symptoms.

⚠ COMMON SIDE EFFECTS



Temporary
blurred vision



Mild stinging
or irritation

⚠ IMPORTANT WARNINGS



Persistent eye
pain or vision
changes require
evaluation rather
than repeated
self-treatment.

★ KEY POINT

Use only as directed
by a healthcare
professional.



PRECAUTIONS



Avoid touching the
bottle tip to the eye.



ADDITIONAL INFORMATION



Close the bottle
tightly after use.



Use as needed or
as advised.



Store at room
temperature.



This content is for educational and general awareness only.
It is not a prescription or substitute for professional medical advice,
diagnosis or treatment. Always consult a qualified healthcare
professional before using any medicine.



PARACETAMOL

DRUG CLASS:

Analgesic and antipyretic



MAIN USES



Relief of mild to moderate pain



Reduction of fever



Commonly used for headache, body aches, toothache and similar short-term pain conditions

COMMON SIDE EFFECTS



Paracetamol usually causes few side effects when used correctly.



Rarely, allergic skin reactions may occur.

IMPORTANT WARNINGS



Taking too much can cause **serious liver damage**.



Avoid taking multiple medicines containing paracetamol at the same time.

PRECAUTIONS



Speak with a healthcare professional if you have liver disease.



Extra caution is needed with regular or heavy alcohol consumption.



Children's products and doses differ from adult products.

SEEK MEDICAL HELP IF



An overdose is suspected, even if there are no symptoms.



Yellow skin/eyes, dark urine, severe rash, swelling or breathing difficulty occurs.

ADDITIONAL INFORMATION



Complete the full course of treatment.



Do not stop the medicine without doctor's advice.



Consult your doctor if symptoms persist or worsen.

KEY POINT

Paracetamol is not an antibiotic. It works by relieving pain and reducing fever. It is not effective against infections like cold or flu.



IBUPROFEN

DRUG CLASS:

Non-steroidal anti-inflammatory drug (NSAID)



MAIN USES



Mild to moderate pain



Fever



Inflammation



Headache, dental pain, muscle/joint pain and menstrual pain

COMMON SIDE EFFECTS



Indigestion



Heartburn



Stomach discomfort



Nausea



Dizziness

IMPORTANT WARNINGS



NSAIDs can cause stomach ulcers or gastrointestinal bleeding.



They may increase the risk of heart attack or stroke, particularly with higher doses or prolonged use.



Kidney problems can occur in susceptible patients.

PRECAUTIONS



Use cautiously in people with a history of stomach ulcers, kidney disease, heart disease or asthma triggered by NSAIDs.



Ask a clinician before combining with blood thinners, aspirin or another NSAID.



NSAIDs require particular caution or caution during pregnancy.

SEEK MEDICAL HELP IF



Black stools or vomiting blood



Chest pain or sudden weakness



Severe allergic reaction



Markedly reduced urination

KEY POINT

Ibuprofen reduces pain, fever and inflammation but may affect the stomach, heart and kidneys.



ADDITIONAL INFORMATION



Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate for long periods without medical advice.



Reference Note: Educational information only; not individual treatment advice.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



QUICK MEDICINE REFERENCE

SALBUTAMOL / ALBUTEROL

DRUG CLASS: Short-acting beta-2 agonist
bronchodilator



EDUCATIONAL REFERENCE ONLY

Not a Prescription



✓ MAIN USES



Relief/prevention of bronchospasm in asthma and other reversible airway disease.

⚠ COMMON SIDE EFFECTS



Tremor



Nervousness



Fast heartbeat



Headache

⚠ IMPORTANT WARNINGS



Increasing need for rescue bronchodilator can indicate poorly controlled asthma and requires medical review.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Heart disease, hyperthyroidism and low potassium risk may require additional care.



SEEK MEDICAL HELP IF



Severe breathlessness not relieved by prescribed rescue treatment



Chest pain



Severe or persistent rapid heartbeat



Do not share this medicine with others.

Regular medical review for good asthma control is important.



ADDITIONAL INFORMATION



Use as a rescue medicine or as advised.



Rinse mouth after inhalation if using a steroid combination.



Do not exceed recommended dose.



Monitor symptoms and inhaler use. Consult a doctor if control worsens.

CODEINE

DRUG CLASS: Opioid



✓ MAIN USES



Pain treatment in selected circumstances.



Cough suppression in limited settings depending on local regulation.

⚠ COMMON SIDE EFFECTS



Drowsiness



Constipation



Nausea



Dizziness

⚠ IMPORTANT WARNINGS



Can cause dependence and dangerous respiratory depression.



Children's use has major restrictions because metabolism can be unpredictable.



Combining with alcohol, benzodiazepines or other sedatives increases risk.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Prescription supervision is essential.



Avoid driving when sedated.



SEEK MEDICAL HELP IF



Slow/shallow breathing



Extreme sleepiness



Blue lips



Inability to wake



Do not share this medicine with others.

Store safely out of reach of children.



ADDITIONAL INFORMATION



Take with water as advised.



Use the lowest effective dose for the shortest time.



Never take more than prescribed.



Report any severe side effects to your doctor.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY




Not a Prescription

DEXTROMETHORPHAN

DRUG CLASS: Cough suppressant



MAIN USES



Temporary suppression of dry/non-productive cough.



COMMON SIDE EFFECTS



Dizziness



Drowsiness



Nausea



IMPORTANT WARNINGS



Excessive amounts can cause serious toxicity and misuse.



Dangerous interactions may occur with MAO inhibitors and some serotonergic medicines.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Do not automatically suppress cough associated with significant mucus, breathing difficulty or persistent illness.



SEEK MEDICAL HELP IF



Confusion



Severe agitation



Breathing difficulty



Persistent unexplained cough



Do not share this medicine with others.



ADDITIONAL INFORMATION



Use only for short-term cough relief.



Follow the dose recommended on the label or by your doctor.



Do not take more than the recommended dose.



Keep out of reach of children.

GUAIFENESIN

DRUG CLASS: Expectorant



MAIN USES



Helps loosen mucus and make productive cough easier to clear.



COMMON SIDE EFFECTS



Nausea



Dizziness



Headache



PRECAUTIONS



Adequate hydration may help.



Persistent cough requires evaluation by a healthcare professional.



KEY POINT

Use only as directed by a healthcare professional.



ADDITIONAL INFORMATION



Drink plenty of fluids while taking this medicine.



Follow the dose recommended on the label or by your doctor.



Do not use for cough with difficulty breathing unless advised by a doctor.



Keep out of reach of children.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE


EDUCATIONAL REFERENCE ONLY



Not a Prescription

METRONIDAZOLE

DRUG CLASS: Nitroimidazole
antibacterial/antiprotozoal medicine



✓ MAIN USES



• Anaerobic bacterial infections



• Bacterial vaginosis



• Trichomoniasis



• Amoebiasis and giardiasis in appropriate situations

⚠ COMMON SIDE EFFECTS



Nausea



Metallic taste



Abdominal discomfort



Headache

⚠ IMPORTANT WARNINGS



Neurological adverse effects are possible with prolonged/high exposure.



Significant interactions occur with some medicines.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Follow the specific product instructions concerning alcohol.



Tell a clinician about liver disease or neurological disorders.



SEEK MEDICAL HELP IF



Numbness/tingling



Seizures



Severe rash



Confusion



Complete the full course even if you feel better unless advised otherwise.

Do not share this medicine with others.



ADDITIONAL INFORMATION



Take at evenly spaced intervals for best results.



Take with food if it upsets your stomach.



Drink plenty of water unless advised otherwise.



Do not stop the medicine early unless advised by your clinician.

CHLORPHENIRAMINE

DRUG CLASS: First-generation antihistamine



✓ MAIN USES



Allergy symptoms such as sneezing, runny nose and itching.



⚠ COMMON SIDE EFFECTS



Drowsiness



Dry mouth



Blurred vision



Constipation

⚠ IMPORTANT WARNINGS



Can significantly impair alertness.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Avoid driving or operating machines if sleepy.



Alcohol and other sedatives may worsen drowsiness.



Use caution with glaucoma and urinary retention.



ADDITIONAL INFORMATION



Most drowsiness occurs shortly after taking the medicine.



Use for the shortest duration necessary for symptom relief.



Do not take more than recommended or more frequently than advised.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.




CLOBETASOL

DRUG CLASS: Very potent topical corticosteroid



MAIN USES



Short-term treatment of steroid-responsive inflammatory skin diseases such as severe eczema or psoriasis when clinically appropriate.

COMMON SIDE EFFECTS



Burning or irritation



Skin dryness

IMPORTANT WARNINGS



Prolonged, extensive or inappropriate use can thin the skin and cause systemic corticosteroid effects.



It can worsen or mask untreated infections.

KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Avoid casual use on the face, groin or large body areas.



Use in children needs particular caution.



Do not use merely because a rash is itchy.

SEEK MEDICAL HELP IF



Worsening infection



Severe skin thinning



Significant systemic symptoms

ADDITIONAL INFORMATION



Use for the shortest duration needed.



Apply a thin layer to the affected area only as directed.



Do not self-medicate or use for longer than recommended.



KEY POINT
Potent medicine. Incorrect use can cause harm.

PERMETHRIN

DRUG CLASS: Topical antiparasitic



MAIN USES



Scabies



Head lice, using the appropriate formulation.

COMMON SIDE EFFECTS



Itching



Mild burning



Redness



Tingling

IMPORTANT WARNINGS



Itching may persist temporarily even after successful scabies treatment and does not automatically mean treatment failure.

KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Follow formulation-specific instructions.



Household/contact management may also be required for scabies or lice.

ADDITIONAL INFORMATION



Wash hands before and after application.



Use the recommended amount and duration only.



Wash clothing, bedding and towels as advised.

KEY POINT

Do not use for any other condition unless directed by a clinician.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.






QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY


Not a Prescription

AMOXICILLIN + CLAVULANATE

DRUG CLASS: Penicillin-class antibiotic + beta-lactamase inhibitor



✓ MAIN USES



Selected susceptible bacterial infections of the respiratory tract, skin, urinary tract and other sites.

⚠ COMMON SIDE EFFECTS



Diarrhoea



Nausea



Vomiting



Rash

⚠ IMPORTANT WARNINGS



Serious allergic reactions can occur in people with penicillin allergy.



Liver injury and C. difficile-associated diarrhoea are uncommon but important risks.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Tell a clinician about previous penicillin/cephalosporin allergy and liver disease.



Antibiotics should not be used for viral illnesses.



SEEK MEDICAL HELP IF



Difficulty breathing



Facial/throat swelling



Severe skin rash



Persistent watery/bloody diarrhoea



Jaundice



Complete the full course even if you feel better unless advised otherwise.



Do not share this medicine with others.



ADDITIONAL INFORMATION



Take at evenly spaced intervals for best results.



Take with food to reduce stomach upset.



Drink plenty of fluids unless advised otherwise.




Report any unusual symptoms promptly to your clinician.



Always consult a qualified healthcare professional before using any medicine.

CEFIXIME

DRUG CLASS: Third-generation cephalosporin antibiotic



✓ MAIN USES



Selected susceptible bacterial infections.

⚠ COMMON SIDE EFFECTS



Diarrhoea



Nausea



Abdominal discomfort

⚠ IMPORTANT WARNINGS



Severe allergic reactions and antibiotic-associated colitis can occur.



★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Tell a healthcare professional about serious beta-lactam antibiotic allergy.



Kidney impairment may affect prescribing.



SEEK MEDICAL HELP IF



Severe allergic reaction



Persistent/bloody diarrhoea



Severe rash



Complete the full course even if you feel better unless advised otherwise.



Do not share this medicine with others.



ADDITIONAL INFORMATION



Take at the same time each day.



Take with food if it upsets your stomach.



Do not stop the medicine early without medical advice.



Alcohol may increase stomach irritation in some people.



Always consult a qualified healthcare professional before using any medicine.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY



Not a Prescription

TADALAFIL

DRUG CLASS: PDE-5 inhibitor



MAIN USES



Erectile dysfunction.



Benign prostatic hyperplasia in appropriate patients.



Certain formulations/indications for pulmonary arterial hypertension.



COMMON SIDE EFFECTS



Headache



Indigestion



Flushing



Back pain



Muscle aches



IMPORTANT WARNINGS



Nitrates must **NOT** be combined with tadalafil.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Cardiovascular disease, kidney/liver impairment and interacting medicines require consideration.



Alcohol may increase the risk of side effects such as low blood pressure, dizziness or headache.



SEEK MEDICAL HELP IF



Chest pain



Prolonged erection (priapism)



Sudden vision or hearing loss



Do not take more than prescribed.

This medicine does not protect against sexually transmitted infections.

Regular medical follow-up is recommended.



ADDITIONAL INFORMATION



Take with water as directed.



Food may delay the onset but does not reduce the effect.




Store in a cool, dry place away from direct sunlight.



Always consult a qualified healthcare professional before using any medicine.

AZITHROMYCIN

DRUG CLASS: Macrolide antibiotic



MAIN USES



Selected susceptible bacterial respiratory, skin and other infections.



Certain sexually transmitted bacterial infections.



COMMON SIDE EFFECTS



Nausea



Diarrhoea



Abdominal pain



Vomiting



IMPORTANT WARNINGS



Does not treat viral colds or influenza.



Can affect cardiac rhythm in susceptible patients.



Unnecessary use contributes to antimicrobial resistance.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Tell a clinician about long-QT syndrome, heart-rhythm problems, liver disease and interacting medicines.



Complete the full course even if you feel better.



SEEK MEDICAL HELP IF



Fainting



Irregular heartbeat



Severe or bloody diarrhoea



Yellowing skin/eyes



Do not share this medicine with others.

Report any unusual symptoms promptly.

This medicine does not protect against sexually transmitted infections.



ADDITIONAL INFORMATION



Take with or without food as directed.



Take doses at evenly spaced intervals.



Store below 25°C in a cool, dry place.



Always consult a qualified healthcare professional before using any medicine.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.






QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

FLUCONAZOLE

DRUG CLASS: Azole antifungal



✓ MAIN USES



Treatment of susceptible fungal infections, including some *Candida* infections.

⚠ COMMON SIDE EFFECTS



Headache



Nausea



Abdominal pain



Diarrhoea

⚠ IMPORTANT WARNINGS



Can cause liver injury and serious skin reactions in rare cases.



Important drug interactions are possible.



Pregnancy-related risks depend on dose and treatment circumstances.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Tell a clinician about liver disease, heart-rhythm conditions and other medicines.



Some medicines can interact with fluconazole.



SEEK MEDICAL HELP IF



Yellowing skin/eyes



Severe rash



Fainting or abnormal heartbeat



Do not stop the medicine without medical advice.

Complete the full course even if you feel better, unless advised otherwise.



ADDITIONAL INFORMATION



Take at the same time each day.



Take with water.



Avoid alcohol as it may increase the risk of liver problems.



Report any unusual symptoms promptly to your clinician.

SILDENAFIL

DRUG CLASS: PDE-5 inhibitor



✓ MAIN USES



Erectile dysfunction.



Certain formulations are also used for pulmonary arterial hypertension under specialist care.

⚠ COMMON SIDE EFFECTS



Headache



Flushing



Indigestion



Nasal congestion



Visual colour changes

⚠ IMPORTANT WARNINGS



Must **NOT** be combined with nitrate medicines because blood pressure can fall dangerously.

★ KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Cardiovascular suitability should be considered.



Discuss alpha-blockers and other interacting medicines.



SEEK MEDICAL HELP IF



Chest pain



Sudden vision or hearing loss



Erection lasting more than four hours (priapism)



Do not take more than prescribed.

This medicine does not protect against sexually transmitted infections.



ADDITIONAL INFORMATION



Take with water as directed.



High-fat meals may delay the effect.



Avoid excessive alcohol.



Store in a cool, dry place away from direct sunlight.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.






QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

NORETHISTERONE / NORETHINDRONE

DRUG CLASS: Progestogen



MAIN USES



- Various menstrual disorders depending on formulation and indication.
- Some formulations are used for contraception.



COMMON SIDE EFFECTS



Irregular bleeding



Headache



Breast tenderness



Nausea



Mood changes



IMPORTANT WARNINGS



Hormonal treatment is not appropriate for every patient.



Some preparations/indications carry thrombotic and other hormonal risks.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Pregnancy should be excluded where appropriate.



Tell a clinician about blood-clot history, migraine, liver disease and unexplained vaginal bleeding.



SEEK MEDICAL HELP IF



Chest pain



Sudden shortness of breath



One-sided leg swelling



Sudden neurological symptoms



ADDITIONAL INFORMATION



Take at the same time each day as advised.



Do not stop the medicine without medical advice.



Does not protect against STIs (sexually transmitted infections).



Regular follow-up with your healthcare professional is important.

TRANEXAMIC ACID

DRUG CLASS: Antifibrinolytic



MAIN USES



Reduction of excessive bleeding, including heavy menstrual bleeding in appropriate patients.



COMMON SIDE EFFECTS



Nausea



Diarrhoea



Abdominal discomfort



IMPORTANT WARNINGS



Blood-clot risk must be considered in susceptible people.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Tell a clinician about previous thrombosis, kidney disease and hormonal contraceptive use.



SEEK MEDICAL HELP IF



Chest pain



Shortness of breath



Pain/swelling in one leg



Vision changes



ADDITIONAL INFORMATION



Take with water as directed.



Use for the duration advised by your doctor.



Do not use to prevent clots or for routine long-term use unless advised.



Report any unusual symptoms promptly.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY



Not a Prescription

DICLOFENAC TOPICAL

DRUG CLASS: Topical NSAID



MAIN USES



Localised musculoskeletal or joint pain depending on formulation.



COMMON SIDE EFFECTS



Skin irritation



Itching



Dryness



Rash at application site



IMPORTANT WARNINGS



Systemic NSAID adverse effects remain possible, although exposure is generally lower than with oral diclofenac.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Do not apply to broken skin unless the specific product permits it.



Avoid eyes and mucous membranes.



Pregnancy and other NSAID use require consideration.



ADDITIONAL INFORMATION



Apply a thin layer to affected area and wash hands after application.



Use for the shortest duration needed.



Do not use more than directed or cover the area unless advised.

CLOTRIMAZOLE

DRUG CLASS: Azole antifungal



MAIN USES



Susceptible fungal skin infections.



Vaginal candidiasis when using appropriate vaginal formulations.



COMMON SIDE EFFECTS



Burning



Stinging



Redness



Local irritation



IMPORTANT WARNINGS



Not every rash is fungal.



Steroid-containing combination creams can mask or worsen infections if used incorrectly.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Use the formulation intended for the affected area.



Persistent or recurrent infection warrants medical assessment.



SEEK MEDICAL HELP IF



Severe swelling



Extensive blistering



Symptoms persist or repeatedly return



ADDITIONAL INFORMATION



Use regularly and complete the full course as advised, even if symptoms improve.



Keep the affected area clean and dry.



Do not self-medicate or use for longer than recommended.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





DICLOFENAC TOPICAL



DRUG CLASS: Topical NSAID



MAIN USES



Localised musculoskeletal or joint pain depending on formulation.



COMMON SIDE EFFECTS



Skin irritation



Itching



Dryness



Rash at application site



IMPORTANT WARNINGS



Systemic NSAID adverse effects remain possible, although exposure is generally lower than with oral diclofenac.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Do not apply to broken skin unless the specific product permits it.



Avoid eyes and mucous membranes.



Pregnancy and other NSAID use require consideration.



ADDITIONAL INFORMATION



Apply a thin layer to affected area and wash hands after application.



Use for the shortest duration needed.



Do not use more than directed or cover the area unless advised.

CLOTRIMAZOLE

DRUG CLASS: Azole antifungal



MAIN USES



Susceptible fungal skin infections.



Vaginal candidiasis when using appropriate vaginal formulations.



COMMON SIDE EFFECTS



Burning



Stinging



Redness



Local irritation



IMPORTANT WARNINGS



Not every rash is fungal.



Steroid-containing combination creams can mask or worsen infections if used incorrectly.



KEY POINT

Use only as directed by a healthcare professional.



PRECAUTIONS



Use the formulation intended for the affected area.



Persistent or recurrent infection warrants medical assessment.



SEEK MEDICAL HELP IF



Severe swelling



Extensive blistering



Symptoms persist or repeatedly return



ADDITIONAL INFORMATION



Use regularly and complete the full course as advised, even if symptoms improve.



Keep the affected area clean and dry.



Do not self-medicate or use for longer than recommended.





TRAMADOL

DRUG CLASS: Opioid analgesic



✓ MAIN USES



Management of pain when an opioid analgesic is considered appropriate

⚠ COMMON SIDE EFFECTS



Nausea



Dizziness



Drowsiness



Constipation



Headache

⚠ IMPORTANT WARNINGS



Can cause dependence and misuse.



Excessive doses or combination with other sedating medicines can cause dangerous breathing suppression.



May increase the risk of seizures and serotonin syndrome in susceptible people or with interacting medicines.



PRECAUTIONS



Do not combine with alcohol or sedative medicines without medical advice.



Avoid driving if drowsy or dizzy.



Tell a clinician about antidepressants, seizure disorders and breathing problems.



SEEK MEDICAL HELP IF



Slow or difficult breathing



Extreme sleepiness or inability to wake



Seizure



Severe confusion, agitation, fever or muscle rigidity



ADDITIONAL INFORMATION



Take exactly as prescribed by your doctor.



Do not stop the medicine suddenly without medical advice.



Consult your doctor if symptoms persist or worsen.



KEY POINT

Tramadol is effective for moderate to severe pain but must be used safely under medical supervision.



KETOROLAC

DRUG CLASS: NSAID



✓ MAIN USES



Short-term management of moderately severe acute pain

⚠ COMMON SIDE EFFECTS



Stomach discomfort



Nausea



Headache



Dizziness



Drowsiness

⚠ IMPORTANT WARNINGS



Intended only for **short-term treatment**.



Carries significant risks of gastrointestinal bleeding and kidney injury.



Should not be used together with another NSAID.



PRECAUTIONS



Avoid in people with active gastrointestinal bleeding or significant kidney problems unless specifically managed by a clinician.



Extra caution is required in older adults and people who are dehydrated.



Tell a clinician about anticoagulant medicines.



SEEK MEDICAL HELP IF



Blood in vomit or stools



Severe abdominal pain



Reduced urine output



Chest pain



Severe allergic reaction



ADDITIONAL INFORMATION



Take with food to minimize stomach upset.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Ketorolac relieves acute pain but carries higher risks; follow dosage and duration instructions carefully.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY



Not a Prescription

SILVER SULFADIAZINE

DRUG CLASS: Topical antimicrobial



✓ MAIN USES



Prevention/treatment of infection in selected burn wounds under clinical care.

⚠ COMMON SIDE EFFECTS



Burning



Itching



Skin discomfort

⚠ IMPORTANT WARNINGS



Can cause serious skin reactions or blood-cell abnormalities in rare cases.

★ KEY POINT

Use only as directed by a healthcare professional.



🛡 PRECAUTIONS



Sulfonamide allergy should be discussed.



Large burn areas require professional monitoring.



Pregnancy and very young infants require special consideration.

★ SEEK MEDICAL HELP IF



Severe rash/blistering



Fever



Unusual bruising or weakness

i ADDITIONAL INFORMATION



Clean the area gently before application as advised by your healthcare professional.



Use regularly and for the full duration as directed.



Do not self-medicate or use for longer than recommended.




KEY POINT


For external use only. Do not apply in eyes, mouth or inside the body.

FRAMYCETIN

DRUG CLASS: Aminoglycoside antibiotic for topical use



✓ MAIN USES



Certain susceptible bacterial skin infections depending on formulation.

⚠ COMMON SIDE EFFECTS



Local irritation



Redness



Allergic/contact dermatitis

⚠ IMPORTANT WARNINGS



Antibiotics should not be used for fungal or viral skin conditions.



Prolonged indiscriminate use can encourage resistance or sensitisation.

★ KEY POINT

Use only as directed by a healthcare professional.



🛡 PRECAUTIONS



Avoid unnecessary use on large damaged areas unless medically directed.

★ SEEK MEDICAL HELP IF



Severe skin reaction



Swelling or difficulty breathing

i ADDITIONAL INFORMATION



Apply to clean, dry skin. Complete the full course as advised.



Follow the directions on the label or as advised by a healthcare professional.



Do not self-medicate or use for longer than recommended.



KEY POINT

For external use only. Avoid contact with eyes and mucous membranes.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





ALBENDAZOLE

DRUG CLASS: Anthelmintic



MAIN USES



Treatment of susceptible parasitic worm infections.



COMMON SIDE EFFECTS



Abdominal pain



Nausea



Headache



Dizziness



IMPORTANT WARNINGS



Longer or high-intensity courses can affect liver function and blood-cell counts.



KEY POINT

Use only as directed. Do not share this medicine with others.



PRECAUTIONS



Pregnancy considerations are important.



Longer courses may require blood counts and liver monitoring.



SEEK MEDICAL HELP IF



Fever or signs of infection



Unusual bleeding



Yellow skin/eyes



Severe rash



ADDITIONAL INFORMATION



Take at the same time(s) each day as advised for best results.



Complete the full course even if you feel better.



Do not self-medicate or use for longer than recommended.



KEY POINT

Albendazole is used to treat worm infections. Do not use without medical advice.

POVIDONE-IODINE

DRUG CLASS: Topical antiseptic



MAIN USES



Antisepsis of skin



Cleaning/preparation of appropriate wounds or skin depending on formulation.



COMMON SIDE EFFECTS



Local irritation



Redness



Dryness



IMPORTANT WARNINGS



Intended for external use unless a specific formulation states otherwise.



Significant iodine absorption can occur with extensive/prolonged application.



KEY POINT

Use only as directed. Avoid contact with eyes.



PRECAUTIONS



Use cautiously over large wounds, severe burns or prolonged periods.



Thyroid disease and iodine sensitivity require consideration.



SEEK MEDICAL HELP IF



Severe skin reaction



Swelling or difficulty breathing



ADDITIONAL INFORMATION



Wash hands after use unless applying to the hands.



Follow the directions on the label or as advised by a healthcare professional.



Do not self-medicate or use for longer than recommended.



KEY POINT

Povidone-iodine helps clean and protect skin from infection.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY



Not a Prescription

BISACODYL

DRUG CLASS: Stimulant laxative



✓ MAIN USES



Short-term constipation



Bowel preparation before certain procedures

⚠ COMMON SIDE EFFECTS



Abdominal cramps



Stomach discomfort



Diarrhoea

⚠ IMPORTANT WARNINGS



Frequent or prolonged unsupervised use may lead to dependence on laxatives or disruption of normal bowel function.

★ KEY POINT

Use only as directed. Not for regular or long-term use without medical supervision.



🛡 PRECAUTIONS



Persistent constipation requires evaluation.



Do not use simply to promote weight loss.



Ask for medical advice with severe abdominal pain, nausea or vomiting.

★ SEEK MEDICAL HELP IF



Rectal bleeding



Severe abdominal pain



No bowel movement despite treatment

i ADDITIONAL INFORMATION



Take with plenty of water unless advised otherwise.



Follow the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.


💡 KEY POINT

Bisacodyl works in the colon to help relieve constipation by stimulating bowel movement.



LACTULOSE

DRUG CLASS: Osmotic laxative



✓ MAIN USES



Constipation



Reduction of blood ammonia in certain patients with significant liver disease

⚠ COMMON SIDE EFFECTS



Gas



Bloating



Abdominal cramps



Diarrhoea



Nausea

⚠ IMPORTANT WARNINGS

Excessive use may cause significant diarrhoea and disturbances in body electrolytes.



Na⁺

K⁺

Cl⁻

★ KEY POINT

Lactulose draws water into the bowel, softens stool and helps it pass more easily.



🛡 PRECAUTIONS



Adequate fluid intake is important unless medically restricted.



Persistent constipation requires evaluation.

★ SEEK MEDICAL HELP IF



Severe diarrhoea



Symptoms of dehydration

Severe abdominal pain

i ADDITIONAL INFORMATION



Take with or without food, as advised.



Take regularly at the same time for best results.



Do not self-medicate or use for longer than recommended.

💡 KEY POINT

Lactulose is safe and effective when used as directed under medical advice.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.





QUICK MEDICINE REFERENCE

EDUCATIONAL REFERENCE ONLY

Not a Prescription

PANTOPRAZOLE

DRUG CLASS: Proton-pump inhibitor



✓ MAIN USES



GERD



Acid-related
oesophageal damage



Conditions involving
excessive gastric-acid
secretion

⚠ COMMON SIDE EFFECTS



Headache



Diarrhoea



Nausea



Abdominal discomfort



Gas

⚠ IMPORTANT WARNINGS



Prolonged use should have
an appropriate medical
indication.



Long-term PPI treatment can
be associated with nutrient
deficiencies and certain
gastrointestinal complications.



PRECAUTIONS



Review prolonged treatment with a
healthcare professional.



Tell a clinician about liver disease
and other medicines.



SEEK MEDICAL HELP IF



Severe persistent diarrhoea



Serious allergic reaction



Unexplained muscle spasms, weakness
or abnormal heartbeat



ADDITIONAL INFORMATION



Take with food or milk
if stomach upset
occurs.



Use the lowest effective dose
for the shortest time
necessary.



Do not self-medicate
or use for longer than
recommended.



KEY POINT

Pantoprazole reduces stomach acid
and helps treat acid-related
conditions. Use only as advised.

DOMPERIDONE

DRUG CLASS: Dopamine-antagonist/
prokinetic antiemetic



✓ MAIN USES



Used in some countries
for nausea and vomiting.

⚠ COMMON SIDE EFFECTS



Dry mouth



Headache

⚠ IMPORTANT WARNINGS



Domperidone can increase the
risk of serious abnormal heart
rhythms in susceptible patients.



It should not be treated as a
routine medicine for every case
of acidity or indigestion.



PRECAUTIONS



Extra caution with existing
heart-rhythm disorders.



Important interactions occur with
medicines that affect cardiac rhythm
or domperidone metabolism.



Use should follow applicable local
prescribing guidance.



SEEK MEDICAL HELP IF



Fainting



Palpitations



Fast or irregular heartbeat



ADDITIONAL INFORMATION



Take with food if it
causes stomach
discomfort.



Use the lowest effective dose
for the shortest time
necessary.



Do not self-medicate
or use for longer than
recommended.



KEY POINT

Domperidone helps relieve
nausea and vomiting by
improving stomach emptying.



This content is for educational and general awareness only.
It is not a prescription or substitute for professional medical advice,
diagnosis or treatment. Always consult a qualified healthcare
professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



PROCHLORPERAZINE

DRUG CLASS: Phenothiazine antiemetic



MAIN USES



Severe nausea and vomiting



Vertigo in selected circumstances



COMMON SIDE EFFECTS



Drowsiness



Dizziness



Dry mouth



Constipation



IMPORTANT WARNINGS



Can cause extrapyramidal/movement reactions.



Rarely, a severe reaction called neuroleptic malignant syndrome can occur.



KEY POINT

Use with caution as it can cause movement-related reactions and severe rare complications.



PRECAUTIONS



Extra caution in Parkinson's disease, older adults and patients taking other sedating medicines.



Avoid driving if drowsy.



SEEK MEDICAL HELP IF



Severe muscle stiffness



Fever/confusion



Uncontrolled movements



Fainting



ADDITIONAL INFORMATION



Take with food if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Prochlorperazine is used only when clearly needed and under medical supervision.

LOPERAMIDE

DRUG CLASS: Antidiarrhoeal medicine



MAIN USES



Short-term control of acute diarrhoea



Selected chronic diarrhoeal conditions under medical supervision



COMMON SIDE EFFECTS



Constipation



Dizziness



Nausea



Abdominal cramps



IMPORTANT WARNINGS



Taking excessive amounts can cause life-threatening heart-rhythm disturbances.



It is not appropriate for every type of infectious diarrhoea.



KEY POINT

Use only as directed. Do not take more than the recommended amount.



PRECAUTIONS



Do not self-treat bloody diarrhoea, black stools or diarrhoea associated with high fever without assessment.



Avoid exceeding the recommended amount.



SEEK MEDICAL HELP IF



Bloody stool



Severe abdominal swelling



Fainting



Fast/irregular heartbeat



Diarrhoea persists or worsens



ADDITIONAL INFORMATION



Take with food if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Loperamide helps control diarrhoea symptoms, but underlying cause must be addressed if severe.





ONDANSETRON

DRUG CLASS:

5-HT3 receptor antagonist antiemetic

**MAIN USES**

Prevention and treatment of nausea and vomiting associated with chemotherapy



Radiation-related nausea/vomiting



Prevention/treatment of postoperative nausea and vomiting in appropriate settings

**COMMON SIDE EFFECTS**

Headache



Constipation



Weakness or tiredness

**IMPORTANT WARNINGS**

May affect the QT interval and heart rhythm in susceptible patients.



Serotonin syndrome is possible when combined with certain serotonergic medicines.

**PRECAUTIONS**

Tell a clinician about long-QT syndrome, electrolyte abnormalities and heart-rhythm problems.



Review antidepressants and other interacting medicines.

**SEEK MEDICAL HELP IF**

Fainting



Irregular heartbeat



Severe agitation, fever or muscle rigidity



Serious allergic reaction

**ADDITIONAL INFORMATION**

Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.

**KEY POINT**

Ondansetron helps prevent and treat nausea and vomiting. Use only as advised.

METOCLOPRAMIDE

DRUG CLASS:

Dopamine-antagonist antiemetic and prokinetic

**MAIN USES**

Certain cases of gastroparesis



Prevention/treatment of nausea and vomiting in selected clinical situations

**COMMON SIDE EFFECTS**

Drowsiness



Restlessness



Fatigue



Diarrhoea

**IMPORTANT WARNINGS**

Can cause movement disorders, including tardive dyskinesia.



Risk increases with longer exposure.

**PRECAUTIONS**

Treatment duration should be limited according to the indication.



Extra caution with Parkinson's disease, seizure disorders and medicines affecting dopamine.

**SEEK MEDICAL HELP IF**

Uncontrolled facial or body movements



Severe muscle stiffness



Fever and confusion



Seizure

**ADDITIONAL INFORMATION**

Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.

**KEY POINT**

Metoclopramide helps improve stomach emptying and reduces nausea/vomiting in selected cases. Use only as prescribed.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



DICYCLOMINE

DRUG CLASS: Anticholinergic antispasmodic



MAIN USES



Relief of intestinal muscle spasms



Symptomatic treatment of irritable bowel syndrome (IBS)



COMMON SIDE EFFECTS



Dry mouth



Dizziness



Blurred vision



Drowsiness



Constipation



Nausea



IMPORTANT WARNINGS



Can reduce sweating and increase the risk of overheating.



May cause confusion or other anticholinergic effects, particularly in older adults.



PRECAUTIONS



Use cautiously in glaucoma, urinary problems, enlarged prostate, intestinal obstruction, myasthenia gravis and certain heart conditions.



Alcohol may increase drowsiness.



Avoid driving if vision or alertness is affected.



SEEK MEDICAL HELP IF



Difficulty urinating



Severe confusion



Fainting



Rapid heartbeat



Difficulty breathing



ADDITIONAL INFORMATION



Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Dicyclomine helps relieve intestinal spasms and IBS symptoms but can cause anticholinergic side effects.



OMEPRAZOLE

DRUG CLASS: Proton-pump inhibitor (PPI)



MAIN USES



Gastroesophageal reflux disease (GERD)



Acid-related oesophageal injury



Peptic ulcers



Part of some H. pylori eradication regimens



Conditions involving excessive gastric-acid production



COMMON SIDE EFFECTS



Headache



Abdominal pain



Nausea



Diarrhoea



Constipation



Gas



IMPORTANT WARNINGS



Long-term or inappropriate use may be associated with vitamin/mineral deficiencies and other complications.



Persistent symptoms should not automatically be treated indefinitely without evaluation.



PRECAUTIONS



Tell a healthcare professional about other medicines because clinically important interactions can occur.



Long-term users may require medical review.



New alarm symptoms such as difficulty swallowing or unexplained weight loss require assessment.



SEEK MEDICAL HELP IF



Severe or persistent diarrhoea



Bloody/black stools



Severe allergic reaction



New difficulty swallowing



ADDITIONAL INFORMATION



Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Omeprazole reduces stomach acid and helps treat acid-related conditions. Use only as advised.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



HYDROXYZINE

DRUG CLASS: First-generation antihistamine



MAIN USES



Relief of itching caused by allergic conditions



May also be prescribed for anxiety in certain circumstances



COMMON SIDE EFFECTS



Drowsiness



Dry mouth



Dizziness



Headache



IMPORTANT WARNINGS



Can cause significant sedation.



In susceptible patients, hydroxyzine can affect heart rhythm.



PRECAUTIONS



Avoid alcohol and other sedative medicines unless approved by a clinician.



Do not drive when affected by drowsiness.



Tell a clinician about heart-rhythm problems or medicines that can prolong the QT interval.



SEEK MEDICAL HELP IF



Fainting



Abnormal heartbeat



Severe allergic reaction



Extreme sedation



ADDITIONAL INFORMATION



Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Hydroxyzine is used for allergy-related itching and anxiety in some cases, but it can cause strong drowsiness and affect heart rhythm.



DEXAMETHASONE

DRUG CLASS: Corticosteroid



MAIN USES



Severe inflammatory conditions



Allergic disorders



Certain autoimmune and immune-related conditions



Used for many other specialist indications depending on the formulation



COMMON SIDE EFFECTS



Increased appetite



Indigestion



Mood or sleep changes



Increased blood glucose



Fluid retention, particularly with systemic treatment



IMPORTANT WARNINGS



Corticosteroids can suppress the immune system and increase susceptibility to infection.



Longer courses may suppress the body's normal adrenal steroid production.



Long-term use has additional risks including bone, eye and metabolic complications.



PRECAUTIONS



Tell a clinician about diabetes, active infections, ulcers, osteoporosis or eye problems.



Long-term systemic treatment should generally not be stopped abruptly without medical guidance.



Important interactions exist with several medicines.



SEEK MEDICAL HELP IF



Signs of serious infection



Severe mood or behavioural changes



Vision problems



Severe weakness or allergic reaction



ADDITIONAL INFORMATION



Take with food or milk if stomach upset occurs.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Dexamethasone is a powerful steroid used for many serious conditions and should be used only as prescribed by a clinician.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



MEFENAMIC ACID

DRUG CLASS: NSAID



MAIN USES



Short-term mild to moderate pain



Menstrual pain



COMMON SIDE EFFECTS



Diarrhoea



Nausea



Stomach discomfort



Heartburn



Dizziness



IMPORTANT WARNINGS



May cause gastrointestinal bleeding, ulcers and cardiovascular complications like other NSAIDs.



Kidney problems may occur.



PRECAUTIONS



Discuss use if you have ulcers, inflammatory bowel disease, kidney disease, heart disease or NSAID-sensitive asthma.



Avoid using multiple NSAIDs together.



SEEK MEDICAL HELP IF



Black/bloody stools



Vomiting blood



Chest pain



Severe skin or allergic reaction



Reduced urination



ADDITIONAL INFORMATION



Take with food or milk to reduce stomach upset.



Use the lowest effective dose for the shortest time necessary.



Consult your doctor if symptoms persist or worsen.



KEY POINT

Mefenamic Acid is used for pain relief including menstrual pain but should be taken carefully due to possible gut and kidney risks.



NIMESULIDE

DRUG CLASS: NSAID



MAIN USES



Used in some countries for short-term acute pain and primary dysmenorrhoea.



COMMON SIDE EFFECTS



Gastrointestinal discomfort



Nausea



Diarrhoea



Dizziness



IMPORTANT WARNINGS



Nimesulide has particular concern regarding **liver toxicity**.



Its systemic use has been restricted in several jurisdictions and is generally limited to short-term use where authorised.



PRECAUTIONS



Avoid in patients with liver disease unless specifically determined appropriate by a qualified clinician.



Do not combine casually with other potentially liver-toxic medicines.



It should not be presented in this book as a routine first-choice painkiller.



SEEK MEDICAL HELP IF



Loss of appetite



Persistent nausea



Abdominal pain



Dark urine



Yellow skin or eyes



ADDITIONAL INFORMATION



Take with food to minimize stomach upset.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate or use for longer than recommended.



KEY POINT

Nimesulide may be effective for short-term pain relief but has important liver safety concerns and restricted use.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice, diagnosis or treatment. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION



DICLOFENAC

DRUG CLASS: NSAID



MAIN USES



Pain and inflammation



Arthritis-related pain



Musculoskeletal pain



Acute painful inflammatory conditions



COMMON SIDE EFFECTS



Indigestion



Stomach pain



Nausea



Headache



Dizziness



IMPORTANT WARNINGS



Can cause gastrointestinal ulceration or bleeding.



May increase cardiovascular risk, including heart attack and stroke.



Kidney and liver problems can occur.



PRECAUTIONS



Extra caution is required with heart disease, high blood pressure, kidney/liver disease or previous stomach ulcers.



Avoid taking another NSAID at the same time unless specifically instructed.



Tell a clinician about blood thinners and corticosteroids.



SEEK MEDICAL HELP IF



Black or bloody stools



Vomiting blood



Chest pain or breathing difficulty



Yellowing of the skin/eyes



Severe swelling or allergic reaction



ADDITIONAL INFORMATION



Take with food or milk to reduce stomach upset.



Use the lowest effective dose for the shortest time necessary.



Do not self-medicate for long periods without medical advice.



KEY POINT

Diclofenac reduces pain and inflammation but carries stomach, heart, kidney and liver risks.



ACECLOFENAC

DRUG CLASS: NSAID



MAIN USES



Pain and inflammation associated with conditions such as osteoarthritis, rheumatoid arthritis and ankylosing spondylitis



COMMON SIDE EFFECTS



Indigestion



Abdominal discomfort



Nausea



Diarrhoea



Dizziness



IMPORTANT WARNINGS



Like other NSAIDs, it may cause gastrointestinal bleeding or ulceration.



Cardiovascular, kidney and liver risks should be considered.



PRECAUTIONS



Use cautiously with a history of ulcers, kidney/liver disease or cardiovascular disease.



Avoid combining with other NSAIDs unless prescribed.



Discuss use during pregnancy or breastfeeding with a healthcare professional.



SEEK MEDICAL HELP IF



Gastrointestinal bleeding



Severe allergic reaction



Chest pain



Marked swelling or reduced urine output



ADDITIONAL INFORMATION



Take with food to minimize stomach upset.



Do not use for longer than recommended without advice.



Consult your doctor if symptoms persist or worsen.



KEY POINT

Aceclofenac helps relieve pain and inflammation but should be used carefully to reduce serious side effects.



This content is for educational and general awareness only. It is not a prescription or substitute for professional medical advice. Always consult a qualified healthcare professional before using any medicine.



AI-ASSISTED CONTENT
VERIFY WITH CURRENT OFFICIAL
MEDICINE INFORMATION