

# Lecture 13: Continuity

September 02, 2019

Sunil Kumar Gauttam

Department of Mathematics, LNMIIT

## 13.1 $\epsilon - \delta$ Definition of Continuity

**Definition 13.1** Let  $D \subseteq \mathbb{R}$ . Consider a function  $f : D \rightarrow \mathbb{R}$  and a point  $c \in D$ . We say that  $f$  is continuous at  $c$  if for every  $\epsilon > 0$ , there is  $\delta > 0$  such that

$$x \in D \text{ and } |x - c| < \delta \implies |f(x) - f(c)| < \epsilon.$$

**Theorem 13.2** Sequential definition of continuity, i.e., Definition 12.1 is equivalent to  $\epsilon - \delta$  definition of continuity, i.e., Definition 13.1.

**Example 13.3** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by

$$f(x) = \begin{cases} x & \text{if } x \text{ is rational} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$

Show that  $f$  is continuous at  $x = 0$  using  $\epsilon - \delta$  definition.

**Solution:** Let  $\epsilon > 0$  be given. Take  $\delta = \epsilon$ . If  $|x - 0| = |x| < \delta$  then

$$|f(x) - f(0)| = |f(x)| \leq |x| < \epsilon.$$

■

**Example 13.4** Let  $f(x) = x^2$  for all  $x \in [a, b]$ , where  $a, b \in \mathbb{R}, a < b$ . Show that  $f$  is continuous on  $[a, b]$  using  $\epsilon - \delta$  definition.

**Solution:** Let  $c \in [a, b]$  and  $\epsilon > 0$  be given. Define  $R = \max\{|a|, |b|\}$ . Choose  $\delta < \frac{\epsilon}{2R}$ . Let  $x \in [a, b]$  be such that  $|x - c| < \delta$ . Then  $|x + c| \leq |x| + |c| < R + R$ . Consider

$$\begin{aligned} |f(x) - f(c)| &= |x^2 - c^2| \\ &= |x + c||x - c| \\ &\leq (|x| + |c|)|x - c| \\ &< 2R|x - c| \\ &< 2R\delta \\ &< \epsilon. \end{aligned}$$

Thus,  $f$  is continuous at  $c \in [a, b]$  and hence on  $\mathbb{R}$ , since  $c$  was an arbitrary element of  $\mathbb{R}$ . ■

**Example 13.5** Let  $f(x) = x^2$  for all  $x \in \mathbb{R}$ . Show that  $f$  is continuous on  $\mathbb{R}$  using  $\epsilon - \delta$  definition.

Solution: Let  $c \in \mathbb{R}$  and  $\epsilon > 0$  be given. Choose  $\delta < \min\{1, \frac{\epsilon}{2+|c|}\}$ . Let  $x \in \mathbb{R}$  be such that  $|x - c| < \delta$ . Then  $|x - c| < 1$  so that  $|x| = |x - c + c| \leq |x - c| + |c| < 1 + |c|$ . Consider

$$\begin{aligned} |f(x) - f(c)| &= |x^2 - c^2| \\ &= |x + c||x - c| \\ &\leq (|x| + |c|)|x - c| \\ &< (2 + |c|)|x - c| \\ &< (2 + |c|)\delta \\ &< \epsilon. \end{aligned}$$

Thus,  $f$  is continuous at  $c \in \mathbb{R}$  and hence on  $\mathbb{R}$ , since  $c$  was an arbitrary element of  $\mathbb{R}$ .

**Remark 13.6** Let us see that how easy is working with sequences. We prove the continuity of the function  $f(x) = x^2$  using Definition 12.1. Let  $c \in \mathbb{R}$  and  $(x_n)$  be a sequence in  $\mathbb{R}$  such that  $x_n \rightarrow c$ . Then by part 3 of Limit Theorem for sequences we have  $x_n^2 \rightarrow c^2$  which is same as saying  $f(x_n) \rightarrow f(c)$ .