

Empirical seed transfer zones require conventions for data sharing to increase their utilization by practitioners

Brianna Wieferich^{2,3}, Reed Clark Benkendorf^{1,2*}

¹ Program in Plant Biology and Conservation, Northwestern University, Evanston,

² Negaunee Institute for Plant Conservation Science & Action, The Chicago Botanic Garden, 1000 Lake

³ Dorena Genetic Resource Center, 34963 Shoreview Dr, Cottage Grove, Oregon

Abstract

Empirical seed transfers zones (ESTZs) are becoming increasingly developed to help guide both the development of native plant materials, and the most desirable seed sources for restoration projects. Despite the growing popularity of ESTZs, standards for distributing these data are nascent. In order to maximize the utilization of ESTZs we propose a set of standards to guide the distribution of ESTZs which should allow for making them easier to interact with, and increase the focus of seed collection efforts and foster utilization of the most appropriate seed source available. Further we propose that sharing of metrics of uncertainty for these data, which can help practitioners identify best alternatives for a seed transfer zone, should become common practice. Finally, we briefly introduce an R package eSTZwritR which implements our core suggestions for data dissemination.

IMPLICATIONS FOR PRACTICE:

- Developing a restoration plan in a short time period can be a stressful process. To decrease the chances of simple human induced mistakes we develop standards to increase consistency between eSTZs to make using them in GIS software more consistent.
- We implement these suggestions in an R package ‘eSTZwritR’ which should make adherence to the guidelines more simple for the scientists developing eSTZ products, allowing for a rapid uptake of these conventions.

*Correspondence: rbenkendorf@chicagobotanic.org

- We further suggest the incorporation of estimates of uncertainty for spatial eSTZ data products so practitioners can determine the most suitable alternative seed zones when material from a preferred zone is not available.
- Finally we present a collation of known eSTZ products for the Western United States as a single raster stack allowing easy use by practitioners.

INTRODUCTION

[Figure 1 about here.]

Empirical seed transfer zones (ESTZ's) are gaining in popularity among restoration practitioners as a tool to help ensure that the most appropriate locally adapted seed source is utilized at restoration sites (McKay et al. 2005). The development of empirical seed transfer zones (ESTZ's) for a species is a costly and time consuming process, most often involving common garden studies, or genetic studies, with many individual populations being represented as samples (Kramer et al. (2015)). To our knowledge, in Western North America, a couple research groups are regularly developing ESTZ's for multiple species at a time, while intermittently an eSTZ for a single species are being developed by various labs focused on restoration ecology more broadly. Further these zones are being developed via funding from multiple agencies with multiple different final use cases and implementation strategies in mind. While the suggestions for best practices during the development of ESTZs are becoming more available, the results of ESTZs need to be conveyed via spatial products, however standardization or guidance on the best practices for accomplishing this are absent.

We argue that the dissemination of data during restoration is a vital portion of any projects success. . .

Here, using 23 sets of ESTZs produced for 22 taxa, we show that most of the data developed and disseminated, to share the results of an ETZ, are incongruous Massatti et al. (2020). Subsequently, using any consensus (wisdom of the masses) from these data, combined with standard conventions of data sharing, we present a set of guiding standards for researchers to employ to make results more congruous.

Current Condition

We conducted a review of all ESTZs on the Western Wildland Environmental Threat Assessment Center WWETAC website (as of May 1, 2024). Each data product was analyzed for its file name structure (using 5

categories), metadata, naming conventions, and directory structure. All scoring was done by hand, and all analyses were carried out in R version 4.2.1.

Herein we present only what we consider to be the most inconsistent, and likely to interfere with practitioners workflows, results in Panel 1. In summary ...

Recommendations

Consensus exists among the developers of ESTZs for a range of attributes related to distribution of spatial products. Combining those opinions with our perceived best practices for data sharing, and experience as users of each of the existing empirical products, results in the recommendations below.

Directory Structure

[Figure 2 about here.]

ESTZ's should be distributed using a predictable directory structure allowing practitioners to be immediately familiar with where to find relevant contents (Figure XX). This predictable nature should decrease the time required to find particular attributes of the data.

We recommend (FIGURE X) that all directories have two main subdirectories, one containing the essential Data products, preferably in both raster and vector data formats. The other directory contains Information relating to the product, including a formatted citation for data use, a map for quick reference, and any materials describing the production of the product both as a paper, and a text file of quick metadata attributes.

File Naming

[Figure 3 about here.]

The individual files within the directory should follow a simple naming format which is easy for users of various softwares to interpret and readily import for use, while also containing key parameters of the data product. We recommend (FIGURE) that each filename has three main components, in addition to the file extension. The first component is the USDA PLANTS code, the specific taxon, and the second is the type of data used to develop the STZ, the final is the two main regions which the product overlaps. We strongly

recommend the use of the 12 Department of Interior regions as they cover considerable geographic expanses and reflect some degree of ecological patterns.

Maps

Maps should be included within the Information directory. Many questions about ESTZs can be answered quickly and simply from a practitioner consulting a map saved as a PDF with the essential cartographic components. We recommend (FIGURE X) that each map contains the following elements: north arrow, scale bar, state borders, geographically relevant cities, coordinate reference system information, sensible categorical color schemes for the seed zones, a legend, the taxons name as a title, and the maps theme ('Seed Transfer Zones') as a subtitle.

Data Formats

We recommend that the spatial data associated with an ESTZ be distributed using both popular spatial data models, vector and raster. For vector data we advocate for the continued usage of data using the shapefile format, while for raster data we propose the usage of geoTIFFs ('tifs', the .tif extension). In our experience tifs seem to be the most widely used of the raster data models in ecology for non-time series data, they are widely supported by a variety of geographic information systems, and generally seem to perform better than ASCII.

Vector Data Field Attributes

[Figure 4 about here.]

We believe that the fields, or columns, of the vector data should follow a predictable pattern. This will allow humans visualizing the data in a GUI to quickly visually detect their field of interest, and while it's bad practice – allow code to subset columns by position rather than field name. We further recommend the standardization of field names to allow for code and scripts to retrieve these values without more complicated coding techniques.

We recommend that each shapefile has a bare minimum of four fields in the following order and of the following data types. 1) ID (numeric - integer) a unique number associated with each individual polygon in the file, we do not recommend combining polygons into multipolygon units, as individual polygons can

retain information about their Area, and are easier for users to subset. 2) Seed Zone (numeric - integer) a unique identifier for each of the eSTZs delineated by the practitioners, these allow for quick subsetting of the data based on a simple value which is hard to mispecify. 3) SZName (character) a human developed name for the zone this may refer to a components of a principal component analysis, e.g. LOW MEDIUM LOW, or be defined by the analysts. We opine that semi-informative names should be developed before data distribution to help practitioners more easily convey important attributes without having to rely on numeric values which may be more difficult to remember due to their non-descript nature.

In addition to these standard field naming and placement conventions, we further recommend a series of standards for the contents within these essential fields, and how to format any additional fields relevant to the project.

Recommendations for new products

Metadata

Metadata offers a way of rapidly distilling many parameters of the ESTZ development process for quick reference by practitioners. The inclusion of metadata may make it easier for a practitioner to find relevant details which they may otherwise have to peruse the literature files to glean. Given that the results of eSTZS are published in a variety of journals requiring different amounts of methodological reporting this is a fruitful format for sharing data in a standardized way.

Database Tracking Results

We have created a Github page for tracking the results of all ESTZ papers, along with their citations to foster future synthesis, and to ensure practitioners are aware of all available results. We ask that developers of eSTZs make user submitted updates to the database to help ensure their results are found by relevant parties.

Estimating Uncertainty

We have observed considerable consternation from seed collection crews, curators, and restoration practitioners alike over the ‘proper’ classification for both a seed source, and the selection of a seed source for a restoration. In most instances these hesitations relate to a seed source which is present from a population

which ‘straddles’ two or more seed zones. We predict that with the increasing availability of fine resolution spatial data, and the wider availability of ecological relevant variables - which more accurately reflects local ecological heterogeneity - individual portions of seed zones will become more fine, increasing the perimeter to surface area ratio and thereby increasing the prevalence of this already common phenomenon (Gibson et al. 2019).

Based on our literature review we believe that currently eSTZs are distributed only as vector data (e.g. shapefiles) polygons. Vector data convey a sense of separation between the objects they represent, i.e. they are used to represent discrete classes with meaningful borders between them. Common examples of polygon vector data model usage include administrative units (e.g. zip codes, states, and countries), hydrologic basins, and the geographic range of a species. On the other hand raster data, or gridded surfaces, are used for representing continuous phenomena, i.e. gradients. Common examples of raster data model usages include climate variables, land cover classes, and predictions of species habitat suitability.

While we agree with the current prevailing census that in most applications, the use of the polygon vector data model is generally the preferred method of sharing data, we have witnessed enough scenarios where a population crosses multiple seed zones, that we believe the usage of raster data is warranted. Raster data come with an enormous benefit in that they can readily incorporate multiple layers (individual raster files) for each pixel across a domain, thus allowing for a first layer of consensus predictions for each cell (the data conveyed in a vector data set), and a few other levels of prediction, e.g. the second and third most likely classification of seed zone in the following layers. An alternative scenario is that a first raster layer can contain the final assignments, while a second layer has the probability of those assignments contained in it, thereby denoting the confidence in the assignment.

We believe that conveying these uncertainties will behoove data users to understand and explore the caveats with model predictions more. This practice is further grounded in best scientific practice as the spatial data used to develop the initial zones are imperfect, the study itself was imperfect, and the classification process is itself imperfect. Further on an ecological level we believe that a porosity exists between these populations of species – they are by virtue of being components of a species connected at least marginally via gene flow, and the expression of this continuity is the most appropriate course of action for data dissemination.

IMPLEMENTATION

The suggestions above may seem relatively onerous to carry out at the end of a multi year study, especially when considering manuscripts are being prepared for publication and further funding opportunities are being

154 applied for, and staff (e.g. Postdocs) may be leaving the group at the end of the project. For these reasons
155 we have created an R package, eSTZwritR ('pronounced easy rider'), which can implement all of them, less
156 the statistical processing, with minimal user inputs. The package is installable from GitHub via devtools at
157 <https://github.com/sagesteppe/eSTZwritR>.

158 FOR DEVELOPERS

159 The package requires only 4-5 functions to produce a directory with the contents discussed above, with
160 minimal data entry. Most importantly the entries are well outlined and easily entered without requiring
161 close attention to detail, an omnipresent scenario when processing standards by hand.

162 FOR PRACTITIONERS

163 These results should allow for simple utilization of existing empirical seed transfer zone resources. We have
164 re-processed all eSTZ data products we are aware of to follow these standards, with the exception of creating
165 the uncertainty raster layers. We have provided some sample code which showcases loading these data into
166 a GIS which utilizes either R or python coding elements, as well as the freely available QGIS which is set up
167 with an advanced graphical user interface (GUI), which allows a user to navigate via a computer mouse.

168 CONCLUSIONS

169 Seed based active restoration will always be a relatively expensive, yet necessary, option for restoration.
170 Here we present simple standards for the scientists developing eSTZs to use in order to standardize the
171 data products they are developing to ease their implementation. While these conventions should be easy to
172 implement for a sufficiently detail oriented and interested individual, we also present an R package which
173 can quickly achieve these results.

174 LITERATURE CITED

175 Bradley St. Clair J, Kilkenny FF, Johnson RC, Shaw NL, Weaver G (2013) Genetic variation in adaptive
176 traits and seed transfer zones for *pseudoroegneria spicata* (bluebunch wheatgrass) in the northwestern
177 united states. *Evolutionary Applications* 6:933–948

- Doherty KD, Butterfield BJ, Wood TE (2017) Matching seed to site by climate similarity: Techniques to prioritize plant materials development and use in restoration. *Ecological Applications* 27:1010–1023
- Erickson VJ, Mandel NL, Sorensen FC (2004) Landscape patterns of phenotypic variation and population structuring in a selfing grass, *elymus glaucus* (blue wildrye). *Canadian Journal of Botany* 82:1776–1789
- Gibson A, Nelson CR, Rinehart S, Archer V, Eramian A (2019) Importance of considering soils in seed transfer zone development: Evidence from a study of the native *bromus marginatus*. *Ecological Applications* 29:e01835
- Horning ME, McGovern TR, Darris DC, Mandel NL, Johnson R (2010) Genecology of *holodiscus discolor* (rosaceae) in the pacific northwest, USA. *Restoration Ecology* 18:235–243
- Johnson RC, Cashman M, Vance-Borland K (2012) Genecology and seed zones for indian ricegrass collected in the southwestern united states. *Rangeland Ecology & Management* 65:523–532
- Johnson RC, Erickson VJ, Mandel NL, St Clair JB, Vance-Borland KW (2010) Mapping genetic variation and seed zones for *bromus carinatus* in the blue mountains of eastern oregon, USA. *Botany* 88:725–736
- Johnson RC, Hellier BC, Vance-Borland KW (2013) Genecology and seed zones for tapertip onion in the US great basin. *Botany* 91:686–694
- Johnson RC, Horning ME, Espeland EK, Vance-Borland K (2015) Relating adaptive genetic traits to climate for sandberg bluegrass from the intermountain western united states. *Evolutionary Applications* 8:172–184
- Johnson RC, Leger E, Vance-Borland K (2017) Genecology of thurber’s needlegrass (*achnatherum thurberi* [piper] barkworth) in the western united states. *Rangeland Ecology & Management* 70:509–517
- Johnson RC, Vance-Borland K (2016) Linking genetic variation in adaptive plant traits to climate in tetraploid and octoploid basin wildrye [*elymus cinereus* (scribn. & merr.) A. Love] in the western US. *PLoS One* 11:e0148982
- Kramer AT, Larkin DJ, Fant JB (2015) Assessing potential seed transfer zones for five forb species from the great basin floristic region, USA. *Natural Areas Journal* 35:174–188
- Massatti R (2020) Genetically-informed seed transfer zones for *cleome lutea* and *machaeranthera canescens* across the colorado plateau and adjacent regions. Bureau of Land Management
- Massatti R (2019) Genetically-informed seed transfer zones for *pleuraphis jamesii*, *sphaeralcea parvifolia*, and *sporobolus cryptandrus* across the colorado plateau and adjacent regions. Bureau of Land Management
- Massatti R, Shriver RK, Winkler DE, Richardson BA, Bradford JB (2020) Assessment of population genetics and climatic variability can refine climate-informed seed transfer guidelines. *Restoration Ecology* 28:485–493
- McKay JK, Christian CE, Harrison S, Rice KJ (2005) ‘How local is local?’—a review of practical and

211 conceptual issues in the genetics of restoration. *Restoration Ecology* 13:432–440
212 Shryock DF, Havrilla CA, DeFalco LA, Esque TC, Custer NA, Wood TE (2017) Landscape genetic ap-
213 proaches to guide native plant restoration in the mojave desert. *Ecological Applications* 27:429–445

214 **List of Figures**

215	1	Dissemination	11
216	2	Directory Structure	12
217	3	File Naming	13
218	4	Vector Data Field Attributes	14




Figure 1: Dissemination




Figure 2: Directory Structure




Figure 3: File Naming

Example field names in a shapefile					
ID	SeedZone	SZName	AreaAcres	BIO1_R	BIO2_mean
1	1	Salt Desert	12340	20.2	5.1
2	2	Desert Scrub	14230	19.1	7.1
3	3	Pinyon-Juniper/Oak Brush	30142	15.1	10.1
4	4	Montane	9872	12.3	12.3
The first four (blue) fields should be in every file. More fields are optional.					

Figure 4: Vector Data Field Attributes