## Table of Contents

(i)	Table of Contents
(ii)	List of Figures
(iii)	List of Tables
(iv)	List of Acronyms
(v)	Abstract5
(vi)	Introduction
(vii)	Field Methods
1.	Sample Design, Stratification, and Plots Weights
2.	**STRATA RESAMPLE**
3.	Ecological Site Descriptions Completion Assessment
4.	Increase Range around Narrow Ecological Site Benchmarks
5.	Ecological Site Description and Ecological Site Group Comparison - Vegetation 'Benchmarks' $\dots$ pp
6.	An Assessment of Drought Conditions
7.	An Overview of the Analytical Sections
8.	Bare groundpp
9.	Soil Stabilitypp
10.	Noxious Species
11.	Plant Functional Diversity - Coverpp
12.	Plant Functional Diversity - Speciespp
13.	Rare Speciespp
14.	Floristic Quality Indexpp
15.	Glossarypp
16	Acknowledgements

# List of Figures

1.	Figure	1.1 All AIM plots sampled in the field office
2.	Figure	1.2 AIM plots sampled near Dominguez-Escalente NCA xx
3.	Figure	1.3 AIM plots sampled near Gunnison-Gorge NCA
4.	Figure	1.4 Summary of Plot Sampling Effort and Plot Fates
5.	Figure 2	2.1 Number of AIM plots per Ecological Site
6.	Figure 2	2.2 Number of plots which have Quantitative Benchmarks
7.	Figure 3	3.1 Imputed Ranges Around Mean Values
8.	Figure 3	<b>3.2</b> Estimates of Ranges of Quantitative Benchmarks
9.	Figure 4	<b>4.1</b> Proportion of Plots mapped to their Ecological Site Groups
10.	_	<b>4.2</b> Initial Relationship Between Field Verified ESD and ESG extracted from the griddedxx
11.	Figure 4	<b>4.3</b> Relationships Between ESDs and ESGs midway through the cleaning process $\dots \dots xx$
12.	Figure 4	<b>4.4</b> Relationship between ESDs and ESGs at end of classification process $\dots xx$
13.	Figure 4	4.5 Mean Benchmarks ESD and ESGxx
14.	Figure 4	4.6 Distribution of Percent Coversxx
15.	Figure	<b>5.1</b> Stylized topographic map of the area of analysis
16.	Figure	<b>5.2</b> SPEI of the Uncompangre Field Office Area
17.	Figure	<b>5.3</b> Drought Status of the Uncompangre Field Office Area
18.	Figure '	7.1 Predicting Bare ground from Foliar Cover
19.	Figure '	<b>7.2</b> ESD and ESG derived Benchmarks [bare ground]xx
20.	Figure '	<b>7.3</b> Bare ground
21.	Figure '	<b>7.4</b> Land Meeting Benchmarks (Bare Ground)xx
22.	Figure 8	8.1 Estimates of Median Soil Stability
23.	Figure 8	8.2 Land Meeting Benchmarks (Soil Stability)
24.	Figure 9	9.1 Composite "Invasive Index"
25.	Figure	9.2 Invasive Species Index
26.	Figure	9.3 Invasive Species in the NE Field Office
27.	Figure	9.4 Land Meeting Benchmarks (Invasive Species)
28.	Figure	10.1 Validation of Calculations
29.	Figure	10.2 Forb Cover [Benchmarks]xx
30.	Figure	10.3 Shrub Cover [Benchmarks]
31.	Figure 1	<b>10.4</b> Tree Cover [Benchmarks]

32.	Figure	10.5	Grass Cover [Benchmarks]xx
33.	Figure	10.6	Proportion of Plots Meeting Benchmark
34.	Figure	10.7	Total area of each stratum and the overall status of benchmarks
35.	Figure	11.1	Number of Species per Functional Group across ESD
36.	Figure	11.2	Species Observed per AIM Plot
37.	Figure	<b>12.1</b>	Species Occurrencesxx
38.	Figure	13.1	Measured Floristic Qualityxx
39.	Figure	13.2	Comparison of Median Values by Stratumxx
40.	Figure	13.3	Predicted Floristic Qualityxx
41.	Figure	13.4	Model Selection Tablexx
42.	Figure	14.1	A Sample Frame with Five Panelsxx

Note, digits refer to the section which the figure is located in, and the decimal to the figures number within that section.

### List of Tables

1.	Table	1.1 Original Sample Design for the Entire Sample Frame
2.	Table	1.2 Realized Weighted Sample Design for the Entire Sample Framexx
3.		1.3 Original Sample Design for Areas of Critical Environmental Concern and Wilderness Studyxx
4.	Table	1.4 Number of Plots Drawn per ACEC
5.	Table	<b>1.5</b> Realized Weighted Sample Design for the Entire Sample Frame
6.	Table	1.6 Original Sample Design for Dominguez-Escalante National Monument
7.	Table	1.7 Realized Weighted Sample Design for the Entire Sample Framexx
8.	Table	1.8 Original Sample Design for Gunnison Gorge National Monument
9.	Table	3.1 Variation of Benchmarksxx
10.	Table	<b>5.1</b> Variables to calculate Potential Evaporation via the Penman-Montieth equation xx
11.	Table	<b>5.2</b> SPI Values Interpretation
12.	Table	10.1 Land Meeting Benchmarks by Administrative Unit
13.	Table	12.1 Seven Forms of Rarity - Conceptualxx
14.	Table	12.2 Seven Forms of Rarity - Examplesxx

### List of Acronyms

ACEC Area of Critical Environmental Concern

AIM Terrestrial Assess, Inventory, and Monitor

**BLM** Bureau of Land Management

**DEM** Digital Elevation Model

**ES** Ecological Site

 $\mathbf{ESD}$  Ecological Site Description

 $\mathbf{ESG}$  Ecological Site Group

**MLRA** Major Land Resource Area

NCA National Conservation Area

NOC National Operations Center

 ${f NGO}$  Non-Governmental Organization

 $\mathbf{NPS}$ National Park Service

NRCS National Resource Conservation Service

 ${\bf RMP}$ Resource Management Plan

**USFS** United States Forest Service

 ${\bf USGS}$  United States Geological Survey

 $\mathbf{WSA}$  Wilderness Study Area

#### Abstract

The Uncompanier Field office completed the first Terrestrial Assess, Inventory, and Monitor panel which it initiated in summer of 2022. This report summarizes the status and conditions of several key indicators from the AIM data set, in comparison to the reference conditions contained in Ecological Site Descriptions, or when required Ecological Site Groups. All comparisons are made using spatially explicit inferential statistics which allow for the interpretation of the percent of the field office falling into different categories which are meeting management conditions as specified in the Resource Management Plan (RMP).

Virtually all work was conducted in the statistical Programming language R, with occasional use of 'bash' and python. Most project 'styling' and report elements were designed using Latex in an Rmarkdown/Rstudio environment. All work was tracked using the version control software 'git' and is stored on a website known as Github, which contains logs of all major incremental changes in the sections. We hope that these steps make it easy for others to replicate our work, and in short order build upon, and then readily surpass it.

#### Introduction

This document can be considered in two parts. The first four sections, i.e. the sections up to 'An Assessment of Drought Conditions' are materials which do not directly relate to the analyses in the remaining portion of the document. Rather they reflect import processes in establishing the framework which we used to compare plots, and interpret results. The first section, 'Sample Design, Stratification, and Plots Weights', relates to an initial oversight regarding the sample design which made us very slightly re-assign the weights of plots associated with the three different areas of the field office which have different management objectives under the Resource Management Plan. The section 'Ecological Side Descriptions Completion Assessment' was essential to evaluate the progress made, and remaining to be made, regarding the establishment of Ecological Sites, and their documentation via the formal Ecological Site Description process. Many of the plots had benchmarks which were very narrow, and did not reflect either uncertainty associated with sampling errors, or natural variation, we slightly adjusted the range of these benchmarks in 'Increase Range around Narrow Ecological Site Benchmarks'. While we intended to use benchmarks directly from Ecological Site Descriptions, the Natural Resource Conservation Service still has many to complete in our area, largely focused in a Major Land Resource Area which contains most of our higher elevation lands. In order to utilize benchmarks across the entirety of the field office, we turned to Ecological Site Groups, a framework recently developed by the United State Geological Survey for the Colorado Plateau. We had to undertake multiple steps to ensure that the NRCS and USGS approaches would be congruent for the report which are detailed in 'Ecological Site Description and Ecological Site Group Comparison - Vegetation 'Benchmarks'. The final section 'An Assessment of Drought Conditions' relates to an aspect of the AIM sample design regarding time series data. Some other AIM analysts have made note of using time as a predictor of variation in responses within a single 5-year panel, which allows them to weight plots sampled in years with normal of above normal precipitation more than plots in drier years. However, we believed that the drought conditions over the sample period were too great to consider years in isolation, and pooled these data. These sections all served to inform how we carried out the analysis of the AIM indicators in the remaining work.

The second part, the next several sections, up to 'Plant Functional Diversity - Species', form the main body of work, and deal with a number of the Assess, Inventory, and Monitor, Indicators, for which we had enough data to investigate in a meaningful manner. These sections proceed in fashion of increasing ecological and biological complexity, but are written to largely be independent of each other; i.e. modular. Notably, the earlier sections refer to more detailed phenomena which are described in the introductions to the later sections. The first section 'Bareground' documents how much soil across the field office is exposed to wind and precipitation. 'Soil Stability' contextualizes how these soils, based on their aggregate stability, have different potential to soil erosion, and the effects of this process on the field office and adjacent areas. 'Noxious Species' investigates the distribution of weeds across the field office, the taxonomic identities of them, and identifies areas with considerable presence of invasive species. We next turn to investigating the cover of each of the major plant functional types in comparison to reference conditions in Plant Functional Diversity - Cover, and believe this to be the most integral portion of the document. We then identify the number of species in the major, and some finer resolution, functional groups in 'Plant Functional Diversity - Species'.

The remaining sections, 'Rare Species' and 'Floristic Quality Index', show opportunistic applications of the AIM data set outside of the realm of Ecological Sites. In 'Rare Species' we identify the species of conservation concern, across a variety of agencies and Non-Governmental Organizations, as well as species rare under two other non-conservation related metrics. Finally we compute the 'Floristic Quality Index' and model it across the field office using Species Richness data, this metric is commonly used in Midwestern and Eastern states, and we believed it's results were largely congruent to the efforts from the combined AIM indicators.