

Noxious Plants

One of the first acts in the life of a plant, as a seed, is to move. Generally seeds travel only short distances, but less common long distance movements are known to occur naturally (Section X.X), and long distance movement occurs regularly via humans. The long distance movement of seeds by humans, in recent history *i.e.* the last couple thousand years, may result in the seed germinating and the population (if any) which results from it surviving for only a few generations (*waif*) (Nesom (2000)). A seed may also produce a population which is able to persist for many generations in the new ecosystem, and even disperse and spread out across their new landscape (*naturalized*); many of these species are readily incorporated into the existing vegetation and do little to alter it (Nesom (2000), Pysek & Richardson (2010)). A subset of these species which are able to naturalize may displace considerable amounts of the plant species already in the landscape, and in so doing alters the composition of species at ecological sites (*noxious*), hence also the functioning of these sites (*invasive*) (Davies (2011), Evans et al. (2001), Pysek et al. (2012), Land Management (n.d.), and reviewed in Ehrenfeld (2010), Pysek & Richardson (2010)). That a few species can alter the properties of landscapes is at the core of the Ecological Sites. While all invasive plants are noxious, not all noxious plants are invasive, as can be seen from a number of native species which act as aggressive weeds in disturbed settings and which prevent natural succession.

Noxious and invasive species adversely affect nearly all ecosystem services offered by natural areas, increase fire frequency and intensity, and have enormous economic impacts (reviewed in Ehrenfeld (2010), D’Antonio & Vitousek (1992), Duncan et al. (2004), Fantle-Lepczyk et al. (2022), Crystal-Ornelas et al. (2021)). Invasive species at landscape scales have been shown to decrease plant species richness, taxonomic, functional and structural diversity leading to declines in habitat heterogeneity and adversely affecting wildlife (Keeley & Brennan (2012), Ehrenfeld (2010), Klinger & Brooks (2017)). They pose serious threats to the well being of both wildlife, livestock, and humans, via interactions with historic land management alterations, to the fire cycle (D’Antonio & Vitousek (1992), Keeley & Brennan (2012), @). The economic impacts of invasive species include enormous amounts of funds being channeled into their treatment to reduce fuel loads (), treatments to curtail their spread into new areas (), and losses in economic activity e.g. by displacement of grasses more suitable as forage for livestock ().


Most land managers in the Western cold deserts (The Colorado Plateau, Great Basin, and Columbia Plateau) consider noxious annual grasses to pose the greatest challenge towards maintaining ecosystems which allow for multiple uses (Chambers et al. (2009)). Of concern to our office is the increasing adaption of cheatgrass (*Bromus tectorum*), which is already present throughout the field office, towards higher elevation sites (J. T. Smith et al. (2022)). Based on the limited evidence currently available the effect of encroachment of invasive species is of more adverse affect than is attributable to climate change or drought, although synergistic interactions between invasive species and drought still occur (Clarke et al. (2005), Lopez et al. (2022)).

Methods

Creation and maintenance of registries of noxious and invasive species often falls on the Department of Agriculture of the Federal and State governments (Quinn et al. (2013)). Given the focus of these agencies, these lists are generally focused on arable lands used for crop production, with minimal effort placed on natural settings (Quinn et al. (2013)). To better characterize the condition of invasive and noxious species in natural areas, BLM staff co-developed definitive lists of status of species regarding their Noxious or Invasive status in 2019.

To develop a list of invasive plant species for the study area ‘IRanks’, a semi-quantitative expert based assessment of introduced species, were extracted from the C-Values prepared by the Colorado Natural Heritage Program (Section X.X) (Morse et al. (2004), P. Smith et al. (2020)). This was combined with an AIM

species attribute table extract from the vicinity of the study area, the latter dataset contained a handful of synonyms which included codes not present in the former.



```
## During execution of the program, a warning message was generated. The warning
## message is stored in a data frame named 'warn_df'. Enter the following command
## to view the warning message: warnprnt()
```

To detect whether Ecological Sites differed in their resistance to noxious weed invasion a Kruskal-Wallis test was used (Kruskal & Wallis (1952)). Kruskal-Wallis was used due to non-normal data (heavily right skewed), and a small number of replicates for each Ecological Site. The Kruskal-Wallis test offered evidence of a difference ($p = 0.043$), to detect which Ecological Sites differed from others a two-sided Dunn's test with Holms correction for multiple testing was used as a *post-hoc* test (Dinno (2017), Ogle et al. (2022), Holm (1979), Dunn (1964)). Once p-values were adjusted for multiple testing, as we had no *a-priori* hypothesis for which sites would be more resistant, there was no strong statistical evidence that any pairs were significantly different than any others. However, this is most certainly in part due to the very few number of plots per sites (Mdn = 5.5), and the large number of sites (total = 24), rather than an actual lack of resistance in Ecological Sites.

Results

63 introduced, and of these 54 invasive species were detected across all **r** plots. 181 plots had invasive species on them, and 148 plots had more than one invasive species. When reducing these to species of immediate management concern **r** species were present at **r** plots (REVIEW THE LIST AND THIN IT). As expected, the species which occurred the most often were *Bromus tectorum*, *Chenopodium album*, *Lactuca serriola*, *Tragopogon dubius*, *Halogeton glomeratus*, *Sisymbrium altissimum* at 113, 54, 53, 51, 50, 43 of all plots respectively. Most of the species were present across the entirety of the field office, except for ca. a dozen species which were isolated to localities between Delta and Paonia (Figure XX). Several of these species generally occur in wetter, higher elevation, habitats than most terrestrial BLM Land and their spread to BLM is minimal; however a number of the populations are adjacent to USFS land (Figure XX).

A component of reference condition for all Ecological Sites is that invasive species are not present. By this metric the plots which would be meeting their benchmarks is quite low (181), however we feel another consideration is whether an invasive species was detected on the Line-Point Intercepts (Section XX). As reference condition benchmarks still have considerable association with pre-Columbian times ecology, and much has changed in the interim, we believe each of these plots deserves another more liberal consideration. Of the plots which had invasive species, they were detected via line-point intercept at 142 of them.

There were 35 species which were detected on LPI lines at 142 plots, and of these species **r** were of immediate management concern. The proportion of all vegetation cover which was invasive at plots varied from 0%

to 100% (Mdn = 3.66, mean = 14.37). In general, the few plots with exceptionally high cover of invasive species were small parcels of BLM land adjacent to roads, and private lands, in low elevation areas near Delta (figure XX).

A caveat with detecting invasive species is that, in the study area, most of them are annuals (##/##). Given the exceptional drought conditions (SECTION XX) under which field work was conducted, it is quite likely that the estimated covers of them by plot are notably lower than are during more normal conditions (Mack & Pyke (1984), Bowers (1987)). While it seems that their presence at plots is unlikely to be altered, their abundances were likely underestimated in most years of sampling (cite).




Figure 1: Uncommon Invasives in the Northern Field Office

side plantings, and is worth eradication efforts (Pringle (1993+)). *Cynoglossum officinale* is generally limited to higher elevation forested areas, and in the absence of forest fire it's spread may be slow on the portions of BLM land which it inhabits. A couple of these (Prairie Pepperweed (*Lepidium densiflorum*), prostrate knotweed (*Polygonum aviculare*)) are generally associated with heavily compacted soils, and tend to not spread aggressively outside of these areas.




Figure 2: Distribution of Invasive Species Across the Field Office

A handful of the invasive species detected (Whitetops (*Cardaria chalapensis*, *Cardaria draba*, alternatively *Lepidium draba*), Canada thistle (*Cirsium arvense*), field bindweed (*Convolvulus arvensis*), Timothy (*Phleum pratense*)), especially in the Northern Portion of the field office are generally associated with slightly more mesic conditions than occur at most BLM land, especially adjacent to irrigated pastures. Unless these populations are entrenched near streams, their spread is likely curtailed by the general aridity of BLM land, but their successful extirpation, in the face of continual re-colonization from adjacent pasture lands, or if they have invaded wetlands, is unlikely. Baby's Breath (*Gypsophila elegans*) may have been introduced for roadside plantings, and is worth eradication efforts (Pringle (1993+)). *Cynoglossum officinale* is generally limited to higher elevation forested areas, and in the absence of forest fire it's spread may be slow on the portions of BLM land which it inhabits. A couple of these (Prairie Pepperweed (*Lepidium densiflorum*), prostrate knotweed (*Polygonum aviculare*)) are generally associated with heavily compacted soils, and tend to not spread aggressively outside of these areas.

A wide range of invasive species were present throughout the field office in a variety of habitats, and at varying abundances.

rainfall variability. *Journal of Vegetation Science*, 16(2), 237–248.




Figure 3: Invasive Index across the Field Office

<https://CRAN.R-project.org/package=dunn.test>

Duncan, C. A., Jachetta, J. J., Brown, M. L., Carrithers, V. F., Clark, J. K., DiTOMASO, J. M., Lym, R. G., McDANIEL, K. C., Renz, M. J., & Rice, P. M. (2004). Assessing the economic, environmental, and societal losses from invasive plants on rangeland and wildlands. *Weed Technology*, 1411–1416.

Dunn, O. J. (1964). Multiple comparisons using rank sums. *Technometrics*, 6(3), 241–252.

Ehrenfeld, J. G. (2010). Ecosystem consequences of biological invasions. *Annual Review of Ecology, Evolution, and Systematics*, 59–80.

Evans, R. D., Rimer, R., Sperry, L., & Belnap, J. (2001). Exotic plant invasion alters nitrogen dynamics in an arid grassland. *Ecological Applications*, 11(5), 1301–1310.

Fantle-Lepczyk, J. E., Haubrock, P. J., Kramer, A. M., Cuthbert, R. N., Turbelin, A. J., Crystal-Ornelas, R., Diagne, C., & Courchamp, F. (2022). Economic costs of biological invasions in the united states. *Science of the Total Environment*, 806, 151318.

Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 65–70.

Keeley, J. E., & Brennan, T. J. (2012). Fire-driven alien invasion in a fire-adapted ecosystem. *Oecologia*, 169(4), 1043–1052.

Klinger, R., & Brooks, M. (2017). Alternative pathways to landscape transformation: Invasive grasses, burn severity and fire frequency in arid ecosystems. *Journal of Ecology*, 105(6), 1521–1533.

References

Bowers, M. A. (1987). Precipitation and the relative abundances of desert winter annuals: A 6-year study in the northern mohave desert. *Journal of Arid Environments*, 12(2), 141–149.

Chambers, J. C., Leger, E., & Goergen, E. (2009). Cold desert fire and invasive species management: Resources, strategies, tactics, and response. *Rangelands*, 31(3), 14–20.

Clarke, P. J., Latz, P. K., & Albrecht, D. E. (2005). Long-term changes in semi-arid vegetation: Invasion of an exotic perennial grass has larger effects than

Crystal-Ornelas, R., Hudgins, E. J., Cuthbert, R. N., Haubrock, P. J., Fantle-Lepczyk, J., Angulo, E., Kramer, A. M., Ballesteros-Mejia, L., Leroy, B., Leung, B., et al. (2021). Economic costs of biological invasions within north america. *NeoBiota*, 67, 485–510.

D’Antonio, C. M., & Vitousek, P. M. (1992). Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics*, 63–87.

Davies, K. W. (2011). Plant community diversity and native plant abundance decline with increasing abundance of an exotic annual grass. *Oecologia*, 167(2), 481–491.

Dinno, A. (2017). *Dunn.test: Dunn’s test of multiple comparisons using rank sums*.

- Kruskal, W. H., & Wallis, W. A. (1952). Use of ranks in one-criterion variance analysis. *Journal of the American Statistical Association*, 47(260), 583–621.
- Land Management, B. of. (n.d.). *About weeds and invasive species*. <https://www.blm.gov/programs/natural-resources/weeds-and-invasives/about>
- Lopez, B. E., Allen, J. M., Dukes, J. S., Lenoir, J., Vila, M., Blumenthal, D. M., Beaury, E. M., Fusco, E. J., Laginhas, B. B., Morelli, T. L., et al. (2022). Global environmental changes more frequently offset than intensify detrimental effects of biological invasions. *Proceedings of the National Academy of Sciences*, 119(22), e2117389119.
- Mack, R. N., & Pyke, D. A. (1984). The demography of *bromus tectorum*: The role of microclimate, grazing and disease. *The Journal of Ecology*, 731–748.
- Morse, L. E., Randall, J. M., Renton, N., Hiebart, R., & Lu, S. (2004). An invasive species assessment protocol: Evaluating non-native plants for their impact on biodiversity. *Methods in Ecology and Evolution*.
- Nesom, G. L. (2000). Which non-native plants are included in floristic accounts? *Sida, Contributions to Botany*, 189–193.
- Ogle, D. H., Doll, J. C., Wheeler, P., & Dinno, A. (2022). *FSA: Fisheries stock analysis*. <https://github.com/fishR-Core-Team/FSA>
- Pringle, J. S. (1993+). *Flora of north america north of mexico [online], volume 5*. Flora of North America Editorial Committee. http://www.efloras.org/florataxon.aspx?flora_id=1&taxon_id=242324275
- Pysek, P., Jarosik, V., Hulme, P. E., Pergl, J., Hejda, M., Schaffner, U., & Vila, M. (2012). A global assessment of invasive plant impacts on resident species, communities and ecosystems: The interaction of impact measures, invading species’ traits and environment. *Global Change Biology*, 18(5), 1725–1737.
- Pysek, P., & Richardson, D. M. (2010). *Invasive species, environmental change and management, and health*.
- Quinn, L. D., Barney, J. N., McCubbins, J. S., & Endres, A. B. (2013). Navigating the “noxious” and “invasive” regulatory landscape: Suggestions for improved regulation. *BioScience*, 63(2), 124–131.
- Smith, J. T., Allred, B. W., Boyd, C. S., Davies, K. W., Jones, M. O., Kleinhesselink, A. R., Maestas, J. D., Morford, S. L., & Naugle, D. E. (2022). The elevational ascent and spread of exotic annual grass dominance in the great basin, USA. *Diversity and Distributions*, 28(1), 83–96.
- Smith, P., Doyle, Georgia, & Lemly, J. (2020). *Revision of colorado’s floristic quality assessment indices*. Colorado Natural Heritage Program. https://cnhp.colostate.edu/download/documents/2020/CO_FQA_2020_Final_Report.pdf