

## Review

## Biodiversity and Resilience of Ecosystem Functions

Tom H. Oliver,<sup>1,2,\*</sup> Matthew S. Heard,<sup>2</sup> Nick J.B. Isaac,<sup>2</sup>  
David B. Roy,<sup>2</sup> Deborah Procter,<sup>3</sup> Felix Eigenbrod,<sup>4</sup>  
Rob Freckleton,<sup>5</sup> Andy Hector,<sup>6</sup> C. David L. Orme,<sup>7</sup>  
Owen L. Petchey,<sup>8</sup> Vânia Proença,<sup>9</sup> David Raffaelli,<sup>10</sup>  
K. Blake Suttle,<sup>11</sup> Georgina M. Mace,<sup>12</sup>  
Berta Martín-López,<sup>13,14</sup> Ben A. Woodcock,<sup>2</sup> and  
James M. Bullock<sup>2</sup>

Accelerating rates of environmental change and the continued loss of global biodiversity threaten functions and services delivered by ecosystems. Much ecosystem monitoring and management is focused on the provision of ecosystem functions and services under current environmental conditions, yet this could lead to inappropriate management guidance and undervaluation of the importance of biodiversity. The maintenance of ecosystem functions and services under substantial predicted future environmental change (i.e., their ‘resilience’) is crucial. Here we identify a range of mechanisms underpinning the resilience of ecosystem functions across three ecological scales. Although potentially less important in the short term, biodiversity, encompassing variation from within species to across landscapes, may be crucial for the longer-term resilience of ecosystem functions and the services that they underpin.

## The Importance of Resilience

Across the globe, conservation efforts have not managed to alleviate biodiversity loss [1], and this will ultimately impact many functions delivered by ecosystems [2,3]. To aid environmental management in the face of conflicting land-use pressures, there is an urgent need to quantify and predict the spatial and temporal distribution of **ecosystem functions** and **services** (see Glossary) [4–6]. Progress is being made in this area, but a serious issue is that monitoring and modeling the delivery of ecosystem functions has been largely based on the current set of environmental conditions (e.g., current climate, land use, habitat quality). This ignores the need to ensure that essential ecosystem functions will be provided under a range of environmental perturbations that could occur in the near future (i.e., the provision of **resilient ecosystem functions**). The objective of this review is to identify the range of mechanisms that underpin the provision of resilient ecosystem functions to inform better environmental monitoring and management.

A focus on current environmental conditions is problematic because future conditions might be markedly different from current ones (e.g., increased frequency of extreme weather events [7] and pollution [8]) and might therefore lead to rapid, nonlinear shifts in ecosystem function provision that are not predicted by current models. Reactive management might be too slow to avert consequent deficits in function, with impacts for societal well-being [9]. An analogy of this situation is the difference between monitoring whether a bridge is either standing (i.e., providing

<sup>1</sup>School of Biological Sciences, University of Reading, Reading RG6 6AH, UK

<sup>2</sup>NERC Centre for Ecology and Hydrology, Wallingford, UK

<sup>3</sup>Joint Nature Conservation Committee, UK

<sup>4</sup>University of Southampton, Southampton, UK

<sup>5</sup>University of Sheffield, Sheffield, UK

<sup>6</sup>Department of Plant Sciences, University of Oxford, Oxford, UK

<sup>7</sup>Imperial College London, London, UK

<sup>8</sup>Institute of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland

<sup>9</sup>Instituto Superior Técnico, University of Lisbon, Lisbon, Portugal

<sup>10</sup>University of York, York, UK

<sup>11</sup>Department of Earth and Planetary Science, University of California, Berkeley, CA, USA

<sup>12</sup>University College London, London, UK

<sup>13</sup>Social-Ecological Systems Laboratory, Universidad Autónoma de Madrid, Madrid, Spain

<sup>14</sup>Environmental Change Institute, Oxford University, Oxford, UK

\*Correspondence: [t.oliver@reading.ac.uk](mailto:t.oliver@reading.ac.uk) (T.H. Oliver).

its function) or collapsed, prompting need for a rebuild, as opposed to monitoring and repairing damage to prevent the collapse from ever happening. In environmental science, attempts have been made to identify this ‘safe operating space’ at a global level to ensure that boundaries are not crossed that could lead to rapid losses in ecosystem functions [10,11]. However, there is a danger that current regional and local assessments of ecosystem functions and management advice do not incorporate such risk assessments. This could result in poor management advice and undervaluation of the importance of biodiversity, because while relatively low levels of biodiversity can be adequate to provide current function [12], higher levels might be needed to support similar levels of function under environmental change [2,13–18]. Therefore, there is a need to identify the characteristics of resilient ecosystem functions and capture these in both predictive models and management guidance.

### Defining and Applying the Resilience Concept

Resilience is a concept with numerous definitions in ecological [19], social [20], and other sciences [21]. In ecology, an initial focus on the stability of ecosystem processes and the speed with which they return to an equilibrium state following disturbance (**recovery** or ‘engineering resilience’ [22]) has gradually been replaced by a broader concept of ‘ecological resilience’ recognizing multiple stable states and the ability for systems to **resist** regime shifts and maintain functions, potentially through internal reorganization (i.e., their ‘adaptive capacity’ [23]). Recent definitions of resilience encompass aspects of both recovery and resistance, although different mechanisms can underpin these and in some cases there might be trade-offs between them [24]. However, some mechanisms can promote both resistance and recovery depending on the timeframe in which a system is observed (e.g., very rapid recovery can look like resistance). Therefore, we treat resistance and recovery here as two related complementary aspects of resilience [25].

There has been much semantic and theoretical treatment of the resilience concept, but here we are concerned with identifying metrics for real-world applications. An ecological system can be defined by the species composition at any point in time [26] and there is a rich ecological literature, both theoretical and experimental, that focuses on the stability of communities [16, 27–29] with potential relevance to resilience. Of course, the species in a community are essential to the provision of many ecosystem functions that are the biological foundation of ecosystem services [3]. However, the stability of species composition itself is not a necessary prerequisite for the resilience of ecosystem functions. Turnover in species communities might be the very thing that allows resilient functions. For example, in communities subjected to climatic warming, cold-adapted species are expected to decline while warm-adapted species increase [30]. The decline of cold-adapted species can be limited through management [31], but in many cases their local loss might be inevitable [32]. If these species have important functional roles, ecosystem functions can suffer unless other species with similar functional roles replace them. Indeed, similar sets of functions might be achieved by very different community structures [33]. Therefore, while the species composition of an ecosystem is typically the target of conservation, it is ecosystem functions, rather than species composition *per se*, that need to be resilient if ecosystem services are to be maintained (Figure 1). In this case the most relevant definition of resilience is the degree to which an ecosystem function can resist or recover rapidly from environmental perturbations, thereby maintaining function above a socially acceptable level. This can be thought of as the ecosystem functions-related meaning of resilience [19], or alternatively as the inverse of ecological ‘vulnerability’ [34]. Resilience in this context is related to the stability of an ecosystem function as defined by its constancy over time [35], but the approach of using a minimum threshold more explicitly measures deficits of ecological function that impact on human well-being (e.g., [14]). Note that here we focus on the resilience of individual ecosystem functions, which might be appropriate for policy formulation (e.g., pollination resilience), although ecosystem managers will ultimately want to consider the suite of ecosystem functions supporting essential services in a given location.

### Glossary

**Alternative stable states:** when an ecosystem has more than one stable state (e.g., community structure) for a particular set of environmental conditions. These states can differ in the levels of specific ecosystem functions.

**Beta diversity:** variation in the composition of species communities across locations.

**(Demographic) Allee effects:** where small populations exhibit very slow or negative growth contrary to the rapid growth usually expected. Explanations range from an inability to find mates or avoid predators or herbivores to limited ability to engage in cooperative behaviors.

**Ecosystem functions:** the biological underpinning of ecosystem services. While ecosystem services are governed by both ecological and social factors (e.g., business demand–supply chains), in this review we focus on the proximate biological processes – such as productivity, pest control, and pollination – that determine the supply of ecosystem services.

**Ecosystem services:** outputs of ecosystem processes that provide benefits to humans (e.g., crop and timber production).


**Effect traits:** attributes of the individuals of a species that underlie its impacts on ecosystem functions and services.

**Functional redundancy:** the tendency for species to perform similar functions, such that they can compensate for changes in each other's contribution to ecosystem processes. Functional redundancy arises when multiple species share similar effect traits but differ in response traits.

**Phenotypic plasticity:** gene-by-environment interactions that lead to the same genotypes expressing changed behavior or physiology under different environmental conditions.

**Resilient ecosystem function:** see main text for a history of the term resilience. The definition used here is the degree to which an ecosystem function can resist or recover rapidly from environmental perturbations, thereby maintaining function above a socially acceptable level.

**Resistance/recovery:** in the context used here, these refer to the tendency of ecosystem function




**Figure 1. Schematic Showing Varying Resilience Levels of an Ecosystem Function ( $\Psi$ ) to Environmental Perturbations (Red Arrows).** Panel (A) shows a system with high resistance but slow recovery; panel (B) shows a system with low resistance but rapid recovery; panel (C) shows a system with both low resistance and slow recovery. Lack of resilience (vulnerability) could be quantified as the length of time that ecosystem functions are provided below some minimum threshold set by resource managers (this threshold shown with the symbol  $\Psi_1$ ) or the total deficit of ecosystem function (i.e., the total shaded-red area). Note that, in the short term, mean function is similar in all systems but in the longer term mean function is lower and the extent of functional deficit is higher in the least resilient system (C).

### Threats to Ecosystem Functions

Environmental change is not unusual (ecosystems have always faced periodic and persistent changes), but anthropogenic activity (e.g., land conversion, carbon emissions, nitrogen cycle disruption, species introductions) is now increasing both the rate and the intensity of environmental change to previously unprecedented levels [36–38]. Rapid changes to the abiotic environment might alter local and regional species pools through environmental filtering and disrupting biotic interactions, leading to changes in the suites of traits and interactions that affect ecosystem functioning [39]. The timescales involved tend to be measured with respect to relevant human interventions; that is, usually over years to decades. The environmental changes may be rapid-onset (e.g., disease), chronic (e.g., habitat loss), or transitory (e.g., drought; Figure 2A) perturbations. Some environmental pressures can show complex temporal patterns. For example, climate change includes transitory perturbations due to climatic extremes overlaid on a background of long-term warming, with the potential for rapid-onset changes if tipping points are reached [40].

The impacts of environmental perturbations on ecosystem functions will depend on the presence of ecosystem characteristics that confer resilience, involving interacting mechanisms at multiple ecological scales (see next section). These processes govern the form of functional



**Figure 2. Different Possible Relationships between Environmental Change ( $\epsilon$ ), time ( $t$ ), and level of ecosystem function provided ( $\Psi$ ).** Panel (A) shows three types of environmental change: rapid onset (A), chronic (B), and transitory perturbation (C). Panel (B) shows that ecosystem function might be relatively resistant to increasing levels of environmental change (D), less resistant (E), or demonstrate hysteresis (F). Panel (C) shows the four qualitatively different outcomes for how ecosystem function varies over time, whether the system is fully resistant to an environmental change (H), shows limited resistance but full recovery (I), or shows limited (J) or low resistance (K) with no recovery of function. The horizontal line at  $\Psi_1$  indicates some minimum threshold for ecosystem function that is set by resource managers. In both panels (A) and (C), short-term stochasticity about trends is omitted for clarity.

response to environmental change (Figure 2B) and their rates relative to the environmental change driver will govern the resilience and ultimate temporal trends in ecosystem function (Figure 2C).

Mechanisms Underpinning Resilient Ecosystem Functions

Previous studies have attempted to identify the characteristics of resilient systems from a broad socioeconomic perspective [20,21], but here we focus on the biological underpinnings of the resilience of ecosystem functions, to inform targeted environmental management practices. The resilience of ecosystem functions to environmental change is likely to be determined by multiple factors acting at various levels of biological organization; namely, species, communities, and landscapes (Table 1). These ecological levels are interconnected so that changes at a particular level can cascade to other levels in the same system. For instance, individual species' responses to environmental change mediate changes in the population abundance and resulting interactions with other species, thus affecting community structure and composition as well as the distribution of effect and **response traits** [39]. These changes can extend to the level of whole ecosystems but are mediated by the ecosystem context, such as landscape level heterogeneity or habitat connectivity, to determine the resilience of ecosystem function.

Here we provide a new assessment of evidence for the mechanisms underpinning the resilience of ecosystem functions across these ecological levels (Table 1). Our assessment is focused on promoting general resilience to a range of different primary threats to ecosystem function.

Species-Level Mechanisms

Species rarely experience identical impacts of environmental change due to interactions between traits, landscape composition, and the scale at which they experience environmental drivers [41,42]. This variation in response within and between individual species determines both the short-term provision and the long-term resilience of ecosystem functions. Below we list five key mechanisms operating at the species level and provide hypotheses for their effects on the resilience of ecosystem functions.

Sensitivity to Environmental Change

Species vary in their capacity to persist in the face of environmental perturbations, mediated by a range of behavioral and physiological adaptations (response traits) [43]. Such traits show both interspecific and intraspecific variation. Individuals with traits conferring reduced sensitivity to environmental change will confer higher resistance on ecosystem functions [44]. For example,

Table 1. Mechanisms Underpinning the Resistance and Recovery of Ecosystem Functions to Environmental Perturbation

Species (Intraspecific)	Community (Interspecific)	Landscape (Ecosystem Context)
Sensitivity to environmental change (RES)	Correlation between response and effect traits (RES)	Local environmental heterogeneity (RES)
Intrinsic rate of population increase (RES/REC)	Functional redundancy (RES/REC)	Landscape-level functional connectivity (RES/REC)
Adaptive phenotypic plasticity (RES/REC)	Network interaction structure (RES)	Potential for alternative stable states (RES/REC)
Genetic variability (RES/REC)	–	Area of natural habitat cover at the landscape scale (RES/REC)
Allee effects (RES/REC)	–	–

The abbreviations RES, REC, and RES/REC indicate the importance of each mechanism for resistance, recovery, or both, respectively.

trees vary in their sensitivity to drought depending on nonstructural carbohydrate levels [44], which in turn might affect the resistance of the ecosystem functions that they provide. Broader suites of traits, such as the plant resource economics spectrum [45], are also likely to explain variation in sensitivity. Note, however, that there might be negative correlations between sensitivity and intrinsic growth rates, with slow-growing species providing more resistant ecosystem functions but with lower capacity to recover if perturbation does occur.

#### *Intrinsic Rate of Population Increase*

The capacity of species populations to grow rapidly from low numbers is determined by a suite of related characteristics including generation time and mortality and fecundity rates. Species with a high intrinsic rate of increase will recover more quickly from environmental perturbations [46] or show resistance if this population reinforcement occurs during the perturbation.

#### *Adaptive Phenotypic Plasticity*

Individuals have the capacity to respond to environmental changes through flexible behavioral or physiological strategies that promote their survival [43] and the resistance of ecosystem functions. For example, thermoregulatory behavior appears to be an essential survival tool in many ectotherms that operate in temperature conditions close to or beyond their physiological limits [47]. Additionally, adaptations might allow flexibility to maximize resource acquisition and growth rates in changed environmental conditions, enabling more rapid population recovery and recovery of ecosystem function.

#### *Genetic Variability*

Higher adaptive genetic variation increases the likelihood that genotypes that are tolerant to a given environmental perturbation will be present in a population [18]. This reduces the population impacts of environmental perturbations [48] and promotes resistance of ecosystem functions [49]. In addition, the persistence of tolerant genotypes locally means that population recovery rates are likely to be higher, leading to enhanced function recovery rates [48,50]. Adaptive genotypes can be present in standing genetic variation, which is more likely at higher effective population sizes. Alternatively, they can arise locally through mutation or through immigration from other populations [18]. It is also becoming increasingly apparent that epigenetic effects can provide heritable variation in ecologically relevant traits [51].

#### *Allee Effects*

**Allee effects** make populations more susceptible to environmental perturbations causing crashes from which it is difficult to recover [52,53]. Certain species are more susceptible to Allee effects through mechanisms such as an inability to find mates or avoid predators or a limited ability to engage in cooperative behaviors.

#### *Community-Level Mechanisms*

Beyond the tolerance and adaptability of individuals, the composition and structure of the biological community is of particular importance for the resilience of ecosystem functions. Below we list three key underpinning mechanisms.

#### *Correlation between Response and Effect Traits*

If the extent of species' population decline following an environmental perturbation (mediated by response traits) is positively correlated with the magnitude of species' effects on an ecosystem function (via **effect traits**), this will lead to less resistant ecosystem functions [39,54]. This might occur if the same traits mediate both response and effects or through indirect associations between different traits. Correlations and trade-offs are probably a common aspect of traits as a result of biophysical limitations in structure and function [55]. For example, traits such as body size have been linked with both sensitivity to environmental change (response traits) and the

maintenance of ecosystem functions (effects traits) such as pollination by bees [56,57], nutrient recycling by dung beetles [56], and pest control by predatory invertebrates [58,59]. By contrast, completely uncorrelated response and effects traits cause higher resistance in ecosystem function, since responses of species to environmental change are decoupled from their effects on function [54,56]. For example, Diaz *et al.* [39] summarize several studies that show no correlation between decomposability in plants (an effect trait for nutrient cycling and soil fertility) and persistence in the seed bank (a response trait to disturbance under agricultural intensification).

### Functional Redundancy

When multiple species perform similar functions (i.e., species exhibit some **redundancy** in their contributions to ecosystem processes), the resistance of an ecosystem function will be higher if those species also have differing responses to environmental perturbations [60,61]. This gives rise to the 'insurance effect' of biodiversity [62], which is well supported both empirically [14,15] and theoretically [16,28]. Underpinning mechanisms include a statistical effect, where averaging across independently fluctuating species populations results in higher resistance ('portfolio effects'), which is enhanced further where there is negative spatial and/or temporal covariance (asynchrony) between species' population sizes, driven by differing responses to environmental change or competition [14–16,28,62].

The functional roles of species can be mediated by either continuous or categorical traits (e.g., complementary effect traits such as sward- and ground-active predators for pest control [63]). Resistance is increased both by more species in total (assuming that there is variation in their response traits) and, for a given total number of species, when they are dispersed equally across effect-trait space (Figure 3). In reality, intraspecific variation in traits also occurs, and where this is substantial relative to interspecific variation it might be relevant to consider the redundancy and dispersion of individuals across effect-trait space [64].

### Network Interaction Structure


Most of the theory and empirical work discussed above concerns organisms occupying a single trophic level, but interactions between species (e.g., predation, parasitism, mutualism) can have large influences on community responses to environmental change [2,65]. Loss of highly connected species in interaction networks can cause extinction cascades and reduce network stability [66–68]. If these species are particularly sensitive to environmental change, the resistance of the ecosystem functions they provide will be low [69]. Impacts on ecosystem function will be greater when response and effect traits are correlated and patterned in networks along extinction cascades. For example, body size is linked with both extinction risk and the provision of ecosystem functions in taxa including pollinators [56] and pest-control agents [70]. In general, highly connected nested networks dominated by generalized interactions are less susceptible to cascading extinction effects and provide more resistant ecosystem functions, in contrast to networks dominated by strong specialized interactions [71,72].

An important consideration is that the impacts of species loss are likely to lead to changes in the abundance of surviving species, so that the presence or absence of density compensation following species loss can be the key predictor of ecosystem function provision [56,67,73]. For example, atmospheric deposition of nitrogen can result in species loss from some plant communities but density compensation of remaining species might support net primary productivity [74].

### Landscape-Level Mechanisms

The intraspecific- and community-level mechanisms described above are influenced by the environmental context of both the local site and the wider landscape. The landscape context





Trends in Ecology &amp; Evolution

**Figure 3. Functional Redundancy and Effects on Resilience of Ecosystem Functions.** Complementary effect-trait space occupied by all species in a community can be characterized by an  $n$ -dimensional hypervolume for continuous traits [main panels (A–C)] or as discrete functional groups for categorical traits (inset panels A–C). A high density of species spread evenly across complementary trait space [(A), shown for two of  $n$  possible traits] leads to higher resistance of ecosystem functions. This is shown in panel (D) (scenario A), which shows the hypothetical average impact on ecosystem function as species are lost from a community under increasing environmental perturbation. The same number of species less evenly dispersed across complementary effect-trait space [i.e., a more ‘clumped’ distribution, (B)] leads to less resistant ecosystem functions [(D), scenario B]. Similarly, fewer species that are evenly but thinly spread across complementary effect-trait space (C), also lead to less resistant ecosystem functions. In both cases, the communities are said to have lower ‘functional redundancy’. The exact rate of loss of ecosystem function will be context dependent (e.g., depending on initial number of species, ordering of species extinctions, and degree of species clustering in trait space).

determines the local and regional species pool and also the abiotic environment, which can modify the impacts of environmental perturbations on individuals and communities.

#### Local Environmental Heterogeneity

Spatial heterogeneity can enhance the resistance of ecosystem functions by: (i) facilitating the persistence of individual species under environmental perturbations by providing a range of resources and microclimatic refugia [75–78]; and (ii) increasing overall species richness [79] and, therefore, functional redundancy. These heterogeneity effects can operate at: the fine scale, for example through vegetation structural diversity [75]; the medium scale, for example through topographic diversity [76]; or the larger scale, for example, through diversity of land cover types [77,78]. Additionally, environmental heterogeneity across locations (promoting **beta diversity**) has been shown to increase the stability of ecosystem functions [27].

#### Landscape-Level Functional Connectivity

Metapopulation theory suggests that populations in well-connected landscapes will persist better or recolonize more rapidly following environmental perturbation (the ‘rescue effect’). Empirical studies confirming this hypothesis range from mesocosm experiments [80,81] to

landscape-level field studies [82,83]. This prediction extends to metacommunities and experiments have shown that connectivity enhances community recovery after local perturbations [81,84]. In a few cases, this recovery of community structure through dispersal has been shown to lead to recovery of ecosystem functions such as productivity and carbon sequestration to pre-perturbation levels, a process termed 'spatial insurance' [85,86].

#### *Area of Natural Habitat Cover at the Landscape Scale*

In addition to improving functional connectivity for particular species, larger areas of natural or seminatural habitat tend to provide a greater range and amount of resources, which promotes higher species richness and larger population sizes for each species [87,88]. This, in turn, is likely to mean greater genetic diversity and functional redundancy, both of which promote the resistance of ecosystem functions [18,60,61].

#### *Potential for Alternative Stable States*

**Alternative stable states** are associated with abrupt shifts in ecosystems, tipping points, and hysteresis, all of which challenge traditional approaches to ecosystem management [17,89]. Ecosystem states maintain their stability through internal feedback mechanisms, which confer resistance on ecosystem functions. However, environmental perturbations can increase the likelihood of regime shift leading to a fundamental change in the assemblages of species providing functions [17]. Systems can be more susceptible to environmental stochasticity and transient perturbations close to these critical tipping points leading to sudden changes to a new equilibrium [53]. Some alternative stable states might be unfavorable in terms of ecosystem functions with return to previous states possible only through large and costly management interventions (hysteresis), thereby limiting the recovery capacity of ecosystem function. Alternative states are documented in a wide variety of ecosystems from local to global scales, although how stable and persistent these are remains uncertain [89–91].

### **Managing for Resilience**

#### *Applied Ecosystem Management*

Ecosystem services are beginning to be integrated within major land-management programs (e.g., the EU Common Agricultural Policy, REDD+). However, the measurement, monitoring, and direct management of ecosystem function resilience in these programs is lacking [92]. The ecological theory and empirical evidence discussed above suggest that multiple factors will determine ecosystem resilience. However, we do not yet know which will be the most important in determining resilience in particular functions or ecosystems. It is clear that some factors will be more amenable to management (e.g., population-level genetic variability, landscape structure [18,31]) than others (e.g., environmental sensitivity of individual species, presence of alternative stable states). Additionally, there can be trade-offs and synergies between resilience and the short-term performance of ecosystem functions [49,93].

#### *Synergies and Trade-Offs with Short-Term Performance*

In some cases there are synergies between the short-term performance of ecosystem functions and their longer-term resilience; for example, if species richness is associated with higher levels of function under current conditions due to complementarity [13], and with higher resilience of function due to higher functional redundancy [39,54]. In these cases, management targeted toward short-term performance will also enhance resilience. In other cases, however, trade-offs can occur. For example, maintaining genetic diversity for the resilience of ecosystem functions may conflict with the aim to produce the 'best locally adapted phenotype' [49]. Much intensive agricultural management currently focuses on such low-diversity systems that produce high levels of provisioning services but which might have low resilience [93]. Furthermore, while habitat heterogeneity can promote the persistence of species through climatic extremes [77,78], it can, in the shorter term, reduce the availability of specific habitats required by key species. In



### Box 1. Indicators of Short-Term Ecosystem Function Flows versus Resilience

The development of indicators for ecosystem functions is hampered by a lack of primary data and there is strong reliance on proxy measures such as habitat extent [94,95]. These proxy measures are currently used to inform on spatial and temporal trends in ecosystem function for the reporting and management of biodiversity change [4–6]. Such models use abiotic variables such as land cover, topography, and climate data as explanatory variables in spatially explicit statistical correlative models [96,97] or process models [98,99] to predict the provision of ecosystem functions and services. However, because models are parameterized and validated (where undertaken) on the current set of environmental conditions they are often suitable only for producing indicators of short-term ecosystem function flows rather than resilience under environmental perturbations (Figure 4).


Attempts at developing resilience indicators for ecological functions have been limited mostly to ‘early-warning systems’ [53,92]. These focus on emergent properties of systems that might precede impending critical state transitions (e.g., ‘critical slowing down’ [53]). However, these properties occur before critical transitions in only a subset of cases and thus are likely to be poor general predictive indicators of resilience [91]. A focus on emergent properties of systems also ignores the mechanisms that underpin resilience and therefore has limited ability to inform management advice.

Therefore, assessments of the resilience of ecosystem functions and services are currently severely lacking. The development of robust yet cost-effective indicators is likely to be dependent on proxy measures that can be both derived from existing monitoring [4] and shown to covary with resilience. For example, an attempt to assess the importance and feasibility of resilience indicators based on expert opinion for coral-reef systems is provided by McClanahan *et al.* [100]. Validation of practicable proxy measures is then important to ensure they are reliable.

these cases, short-term management for higher levels of ecosystem function might hinder resilience.

### Measuring and Monitoring Resilience

Reporting on ecosystem services has focused on the short term [6], despite the acknowledgement of long-term resilience in Earth systems management [10,92]. Therefore, a challenge is the development of robust yet cost-effective indicators of the resilience of ecosystem functions and services (Box 1). To develop indicators, research is needed into current data availability, the feasibility of data collection, and the validation of indicator metrics. The subsequent implementation of resilience indicators to inform environmental management will also require significant interdisciplinary research with the socioeconomic sciences; for example, to ascertain target suites of ecosystem functions in different areas and to set socially acceptable minimum thresholds for functions. An additional challenge will be to identify and balance trade-offs between the resilience of multiple functions. Such research, however, is essential to safeguard the provision of ecosystem functions under the significant environmental perturbations expected within the next century (see Outstanding Questions).



**Figure 4.** Hypothetical Example of Indicator Values for an Ecosystem Function Flow (e.g., Estimates of Pollen Delivery to Crops) or Resilience of that Function (e.g., Pollination under Environmental Perturbations as Measured by Some Combination of the Mechanisms Highlighted in this Review) as an Ecosystem is Degraded over Time. The thresholds to initiate management action (red dotted lines) differ depending on which indicator is used (A for the resilience indicator, B for the ecosystem function flow indicator). Given that remedial management takes time to put in place and become effective, unacceptable losses of ecosystem function might occur if ecosystem function flow indicators are solely relied on. These losses can be costly for society and difficult to reverse.

## Concluding Remarks

In this review we have highlighted mechanisms by which biodiversity, at different hierarchical scales, can influence the resilience of ecosystem functions. We hope that a focus on resilience rather than short-term delivery of ecosystem functions and services, and the consideration of specific underpinning mechanisms, will help to join the research areas of biodiversity–ecosystem function and ecological resilience and ultimately aid the development of evidence-based yet flexible ecosystem management. Further work will also need to draw significantly on other disciplines to develop appropriate indicators for the simultaneous resilience of multiple ecosystem functions.

## Acknowledgments

The authors thank two anonymous reviewers and Volker Grimm for comments and discussion that helped to improve the manuscript. The review forms part of the output from the Tansley Working Groups initiative sponsored by the UK Natural Environment Research Council (NERC) (<http://www.nerc.ac.uk/>). A series of workshops leading to the review were held at Imperial College London. T.H.O. was supported by the Wessex Biodiversity Ecosystem Services Sustainability (BESS) project within the NERC BESS program. V.P. was supported by Fundação para a Ciência e a Tecnologia (BPD/80726/2011). Contributions of OLP were supported by the University of Zurich Research Priority Program on 'Global Change and Biodiversity' and SNSF Project 31003A\_137921 'Predicting community responses to environmental change'.

## References

- Butchart, S.H.M. *et al.* (2010) Global biodiversity: indicators of recent declines. *Science* 328, 1164–1168
- Cardinale, B.J. *et al.* (2012) Biodiversity loss and its impact on humanity. *Nature* 486, 59–67
- Mace, G.M. *et al.* (2012) Biodiversity and ecosystem services: a multilayered relationship. *Trends Ecol. Evol.* 27, 19–26
- Geizendorfer, I.R. *et al.* (2015) Bridging the gap between biodiversity data and policy reporting needs: an Essential Biodiversity Variables perspective. *J. Appl. Ecol.* Published online March 24, 2015. <http://dx.doi.org/10.1111/1365-2664.12417>
- Heltenstein, J. and Kienast, F. (2014) Ecosystem service state and trends at the regional to national level: a rapid assessment. *Ecol. Ind.* 36, 11–18
- Maes J. *et al.* (2013) Mapping and Assessment of Ecosystems and their Services. An Analytical Framework for Ecosystem Assessments under Action 5 of the EU Biodiversity Strategy to 2020 ([http://ec.europa.eu/environment/nature/knowledge/ecosystem\\_assessment/pdf/MAESWorkingPaper2013.pdf](http://ec.europa.eu/environment/nature/knowledge/ecosystem_assessment/pdf/MAESWorkingPaper2013.pdf)).
- Mora, C. *et al.* (2013) The projected timing of climate departure from recent variability. *Nature* 502, 183–187
- Fowler, D. *et al.* (2013) *The Global Nitrogen Cycle in the Twenty-First Century*, The Royal Society
- Oliver, T.H. and Roy, D.B. (2015) The pitfalls of ecological forecasting. *Biol. J. Linn. Soc.* 115, 767–778
- Steffen, W. *et al.* (2015) Planetary boundaries: guiding human development on a changing planet. *Science* Published online February 13, 2015. <http://dx.doi.org/10.1126/science.1259855>
- Scheffer, M. *et al.* (2015) Creating a safe operating space for iconic ecosystems. *Science* 347, 1317–1319
- Winfree, R. *et al.* (2015) Abundance of common species, not species richness, drives delivery of a real-world ecosystem service. *Ecol. Lett.* 18, 626–635
- Balvanera, P. *et al.* (2013) Linking biodiversity and ecosystem services: current uncertainties and the necessary next steps. *Bioscience* Published online December 5, 2013. <http://dx.doi.org/10.1093/biosci/bit003>
- Allan, E. *et al.* (2011) More diverse plant communities have higher functioning over time due to turnover in complementary dominant species. *Proc. Natl. Acad. Sci. U.S.A.* 108, 17034–17039
- Downing, A.L. *et al.* (2014) Multiple diversity–stability mechanisms enhance population and community stability in aquatic food webs. *Ecology* 95, 173–184
- Loreau, M. and de Mazancourt, C. (2013) Biodiversity and ecosystem stability: a synthesis of underlying mechanisms. *Ecol. Lett.* 16, 106–115
- Folke, C. *et al.* (2004) Regime shifts, resilience, and biodiversity in ecosystem management. *Annu. Rev. Ecol. Syst.* 35, 557–581
- Sgrò, C.M. *et al.* (2011) Building evolutionary resilience for conserving biodiversity under climate change. *Evol. Appl.* 4, 326–337
- Brand, F. (2009) Critical natural capital revisited: ecological resilience and sustainable development. *Ecol. Econ.* 68, 605–612
- Biggs, R. *et al.* (2012) Toward principles for enhancing the resilience of ecosystem services. *Annu. Rev. Environ. Res.* 37, 421–448
- Zolli, A. and Healy, A.M. (2013) *Resilience: Why Things Bounce Back*, Headline
- Pimm, S.L. (1984) The complexity and stability of ecosystems. *Nature* 307, 321–326
- Gunderson, L. *et al.* (2010) *Foundations of Ecological Resilience*, Island Press
- MacGillivray, C.W. *et al.* (1995) Testing predictions of the resistance and resilience of vegetation subjected to extreme events. *Funct. Ecol.* 9, 640–649
- Hodgson, D. *et al.* (2015) What do you mean, 'resilient'? *Trends Ecol. Evol.* 30, 503–506
- Standish, R.J. *et al.* (2014) Resilience in ecology: abstraction, distraction, or where the action is? *Biol. Conserv.* 177, 43–51
- Wang, S. and Loreau, M. (2014) Ecosystem stability in space:  $\alpha$ ,  $\beta$  and  $\gamma$  variability. *Ecol. Lett.* 17, 891–901
- Morin, X. *et al.* (2014) Temporal stability in forest productivity increases with tree diversity due to asynchrony in species dynamics. *Ecol. Lett.* 17, 1526–1535
- Karp, D.S. *et al.* (2011) Resilience and stability in bird guilds across tropical countryside. *Proc. Natl. Acad. Sci. U.S.A.* 108, 21134–21139
- Devictor, V. *et al.* (2012) Differences in the climatic debts of birds and butterflies at a continental scale. *Nat. Clim. Change* 2, 121–124
- Oliver, T.H. *et al.* (2012) A decision framework for considering climate change adaptation in biodiversity conservation. *J. Appl. Ecol.* 49, 1247–1255
- Thomas, C.D. *et al.* (2006) Range retractions and extinction in the face of climate warming. *Trends Ecol. Evol.* 21, 415–416
- Gallagher, R.V. *et al.* (2013) Species loss and gain in communities under future climate change: consequences for functional diversity. *Ecography* 36, 531–540

## Outstanding Questions

The following research questions have particular priority for advancing research into the management of resilient ecosystem functions.

Are there thresholds that should be avoided to prevent sudden collapse of ecosystem functions? If so, how quickly are systems moving toward these thresholds and do the thresholds themselves move?

How exactly can each of the mechanisms identified in this review and any others be used to inform applied management to enhance the resilience of ecosystem functions?

How can the relevance and feasibility of these mechanisms be assessed to develop robust indicators for the measurement and monitoring of resilience?

Given that the values people give to ecosystem services are likely to be context dependent over space and time, how do we decide which services and the underpinning functions are priorities in a given area and what the minimum thresholds are?

Given that ecosystem services are the products of both natural capital (i.e., ecosystem functions) and other socioeconomic capitals, what is the relative contribution of resilient ecosystem functions to the maintenance of different ecosystem services over time?

How can the measures to promote resilience be justified when, under stable environmental conditions and in many decision-making-relevant time-scales, they lead to apparent redundancy?

34. Mumby, P.J. *et al.* (2014) Ecological resilience, robustness and vulnerability: how do these concepts benefit ecosystem management? *Curr. Opin. Environ. Sust.* 7, 22–27
35. Hautier, Y. *et al.* (2015) Anthropogenic environmental changes affect ecosystem stability via biodiversity. *Science* 348, 336–340
36. Steffen, W. *et al.* (2015) The trajectory of the Anthropocene: The Great Acceleration. *Anthropocene Rev.* 2, 81–98
37. Krausmann, F. *et al.* (2013) Global human appropriation of net primary production doubled in the 20th century. *Proc. Natl. Acad. Sci. U.S.A.* 110, 10324–10329
38. Simberloff, D. *et al.* (2013) Impacts of biological invasions: what's what and the way forward. *Trends Ecol. Evol.* 28, 58–66
39. Díaz, S. *et al.* (2013) Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecol. Evol.* 3, 2958–2975
40. McNeall, D. *et al.* (2011) Analyzing abrupt and nonlinear climate changes and their impacts. *WIREs Clim. Change* 2, 663–686
41. Swift, T.L. and Hannon, S.J. (2010) Critical thresholds associated with habitat loss: a review of the concepts, evidence, and applications. *Biol. Rev.* 85, 35–53
42. Chase, J.M. and Bengtsson, J. (2010) Increasing spatio-temporal scales: metacommunity ecology. In *Community Ecology: Processes, Models, and Applications* (Verhoef, H.A. and Morin, P.J., eds), pp. 57–68, Oxford University Press
43. Hofmann, G.E. and Todgham, A.E. (2010) Living in the now: physiological mechanisms to tolerate a rapidly changing environment. *Annu. Rev. Physiol.* 72, 127–145
44. O'Brien, M.J. *et al.* (2014) Drought survival of tropical tree seedlings enhanced by non-structural carbohydrate levels. *Nat. Clim. Change* 4, 710–714
45. Reich, P.B. (2014) The world-wide 'fast-slow' plant economics spectrum: a traits manifesto. *J. Ecol.* 102, 275–301
46. Grman, E. *et al.* (2010) Mechanisms contributing to stability in ecosystem function depend on the environmental context. *Ecol. Lett.* 13, 1400–1410
47. Sunday, J.M. *et al.* (2014) Thermal-safety margins and the necessity of thermoregulatory behavior across latitude and elevation. *Proc. Natl. Acad. Sci. U.S.A.* 111, 5610–5615
48. Forsman, A. and Wennersten, L. (2015) Inter-individual variation promotes ecological success of populations and species: evidence from experimental and comparative studies. *Ecography* Published online July 31, 2015. <http://dx.doi.org/10.1111/ecog.01357>
49. Kettenring, K.M. *et al.* (2014) Application of genetic diversity–ecosystem function research to ecological restoration. *J. Appl. Ecol.* 51, 339–348
50. Reusch, T.B.H. *et al.* (2005) Ecosystem recovery after climatic extremes enhanced by genotypic diversity. *Proc. Natl. Acad. Sci. U.S.A.* 102, 2826–2831
51. Bosse, O. *et al.* (2008) Epigenetics for ecologists. *Ecol. Lett.* 11, 106–115
52. Berec, L. *et al.* (2007) Multiple Allee effects and population management. *Trends Ecol. Evol.* 22, 185–191
53. Dai, L. *et al.* (2012) Generic indicators for loss of resilience before a tipping point leading to population collapse. *Science* 336, 1175–1177
54. Suding, K.N. *et al.* (2008) Scaling environmental change through the community-level: a trait-based response-and-effect framework for plants. *Global Change Biol.* 14, 1125–1140
55. Weiher, E. *et al.* (1999) Challenging Theophrastus: a common core list of plant traits for functional ecology. *J. Veg. Sci.* 10, 609–620
56. Larsen, T.H. *et al.* (2005) Extinction order and altered community structure rapidly disrupt ecosystem functioning. *Ecol. Lett.* 8, 538–547
57. Williams, N.M. *et al.* (2010) Ecological and life-history traits predict bee species responses to environmental disturbances. *Biol. Conserv.* 143, 2280–2291
58. Kotze, D.J. and O'Hara, R.B. (2003) Species decline – but why? Explanations of carabid beetle (Coleoptera, Carabidae) declines in Europe. *Oecologia* 135, 138–148
59. Dixon, A.F.G. and Hemptinne, J.L. (2001) Body size distribution in predatory ladybird beetles reflects that of their prey. *Ecology* 82, 1847–1856
60. Moullot, D. *et al.* (2013) A functional approach reveals community responses to disturbances. *Trends Ecol. Evol.* 28, 167–177
61. Moullot, D. *et al.* (2014) Functional over-redundancy and high functional vulnerability in global fish faunas on tropical reefs. *Proc. Natl. Acad. Sci. U.S.A.* 111, 13757–13762
62. Yachi, S. and Loreau, M. (1999) Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *Proc. Natl. Acad. Sci. U.S.A.* 96, 1463–1468
63. Losey, J.E. and Denno, R.F. (1998) Interspecific variation in the escape responses of aphids: effect on risk of predation from foliar-foraging and ground-foraging predators. *Oecologia* 115, 245–252
64. Schindler, D.E. *et al.* (2010) Population diversity and the portfolio effect in an exploited species. *Nature* 465, 609–612
65. Duffy, J.E. (2002) Biodiversity and ecosystem function: the consumer connection. *Oikos* 99, 201–219
66. Dunne, J.A. *et al.* (2002) Network structure and biodiversity loss in food webs: robustness increases with connectance. *Ecol. Lett.* 5, 558–567
67. Fung, T. *et al.* (2015) Impact of biodiversity loss on production in complex marine food webs mitigated by prey-release. *Nat. Commun.* 6, 6657
68. Memmott, J. *et al.* (2004) Tolerance of pollination networks to species extinctions. *Proc. Biol. Sci.* 271, 2605–2611
69. Schleuning, M. *et al.* (2014) Predicting ecosystem functions from biodiversity and mutualistic networks: an extension of trait-based concepts to plant–animal interactions. *Ecography* Published online October 24, 2014. <http://dx.doi.org/10.1111/ecog.00983>
70. Isikber, A.A. and Copland, M.J.W. (2001) Food consumption and utilisation by larvae of two coccinellid predators, *Scymnus levalantii* and *Cycloneda sanguinea*, on cotton aphid, *Aphis gossypii*. *Biocontrol* 46, 455–467
71. Rooney, N. and McCann, K.S. (2012) Integrating food web diversity, structure and stability. *Trends Ecol. Evol.* 27, 40–46
72. Lever, J.J. *et al.* (2014) The sudden collapse of pollinator communities. *Ecol. Lett.* 17, 350–359
73. Berg, S. *et al.* (2014) Ecological communities are vulnerable to realistic extinction sequences. *Oikos* 124, 486–496
74. Stevens, C.J. *et al.* (2011) Addressing the impact of atmospheric nitrogen deposition on Western European grasslands. *Environ. Manage.* 48, 885–894
75. Kindvall, O. (1996) Habitat heterogeneity and survival in a bush cricket metapopulation. *Ecology* 77, 207–214
76. Godfree, R. *et al.* (2011) Multiscale topodaphic heterogeneity increases resilience and resistance of a dominant grassland species to extreme drought and climate change. *Global Change Biol.* 17, 943–958
77. Piha, H. *et al.* (2007) Anuran abundance and persistence in agricultural landscapes during a climatic extreme. *Global Change Biol.* 13, 300–311
78. Oliver, T. *et al.* (2010) Heterogeneous landscapes promote population stability. *Ecol. Lett.* 13, 473–484
79. Stein, A. *et al.* (2014) Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecol. Lett.* 17, 866–880
80. Mora, C. *et al.* (2007) Experimental simulations about the effects of overexploitation and habitat fragmentation on populations facing environmental warming. *Proc. Biol. Sci.* 274, 1023–1028
81. Altermatt, F. *et al.* (2011) Effects of connectivity and recurrent local disturbances on community structure and population density in experimental metacommunities. *PLoS ONE* 6, e19525
82. Oliver, T.H. *et al.* (2013) Population resilience to an extreme drought is influenced by habitat area and fragmentation in the local landscape. *Ecography* 36, 579–586
83. Piessens, K. *et al.* (2009) Synergistic effects of an extreme weather event and habitat fragmentation on a specialised insect herbivore. *Oecologia* 159, 117–126
84. Perdomo, G. *et al.* (2012) The role of temperature and dispersal in moss–microarthropod community assembly after a catastrophic event. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 367, 3042–3049

85. Zitter, C. *et al.* (2013) Functional diversity and management mediate aboveground carbon stocks in small forest fragments. *Ecosphere* 4, art85
86. Symons, C.C. and Arnott, S.E. (2013) Regional zooplankton dispersal provides spatial insurance for ecosystem function. *Global Change Biol.* 19, 1610–1619
87. Hodgson, J.A. *et al.* (2011) Habitat area, quality and connectivity: striking the balance for efficient conservation. *J. Appl. Ecol.* 48, 148–152
88. Fahrig, L. *et al.* (2011) Functional landscape heterogeneity and animal biodiversity in agricultural landscapes. *Ecol. Lett.* 14, 101–112
89. Petraitis, P. (2013) *Multiple Stable States in Natural Ecosystems*, Oxford University Press
90. Schröder, A. *et al.* (2005) Direct experimental evidence for alternative stable states: a review. *Oikos* 110, 3–19
91. Dakos, V. *et al.* (2015) *Resilience Indicators: Prospects and Limitations for Early Warnings of Regime Shifts*, The Royal Society
92. Spears, B.M. *et al.* (2015) Effective management of ecological resilience – are we there yet? *J. Appl. Ecol.* Published online July 31, 2015. <http://dx.doi.org/10.1111/1365-2664.12497>
93. Foley, J.A. (2005) Global consequences of land use. *Science* 309, 570–574
94. Eigenbrod, F. *et al.* (2010) The impact of proxy-based methods on mapping the distribution of ecosystem services. *J. Appl. Ecol.* 47, 377–385
95. Stephens, P.A. *et al.* (2015) Management by proxy? The use of indices in applied ecology. *J. Appl. Ecol.* 52, 1–6
96. Lavorel, S. *et al.* (2011) Using plant functional traits to understand the landscape distribution of multiple ecosystem services. *J. Ecol.* 99, 135–147
97. Henrys, P.A. *et al.* (2014) Mapping natural capital: optimising the use of national scale datasets. *Ecography* 38, 632–638
98. Nelson, E. *et al.* (2009) Modeling multiple ecosystem services, biodiversity conservation, commodity production, and tradeoffs at landscape scales. *Front. Ecol. Environ.* 7, 4–11
99. Villa, F. *et al.* (2014) A methodology for adaptable and robust ecosystem services assessment. *PLoS ONE* 9, e91001
100. McClanahan, T.R. *et al.* (2012) Prioritizing key resilience indicators to support coral reef management in a changing climate. *PLoS ONE* 7, e42884