

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281754117>

Climate change and physical disturbance cause similar community shifts in biological soil crusts

Article in *Proceedings of the National Academy of Sciences* · September 2015

DOI: 10.1073/pnas.1509150112

CITATIONS

197

READS

561

3 authors, including:



[Scott Ferrenberg](#)

University of Montana

74 PUBLICATIONS 3,469 CITATIONS

[SEE PROFILE](#)



[Sasha C Reed](#)

United States Geological Survey

184 PUBLICATIONS 9,689 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Master's Thesis [View project](#)



Alpine-Treeline Warming Experiment [View project](#)

Climate change and physical disturbance cause similar community shifts in biological soil crusts

Scott Ferrenberg¹, Sasha C. Reed, and Jayne Belnap

Southwest Biological Science Center, US Geological Survey, Moab, UT 84532

Edited by William H. Schlesinger, Cary Institute of Ecosystem Studies, Millbrook, NY, and approved August 25, 2015 (received for review May 9, 2015)

Biological soil crusts (biocrusts)—communities of mosses, lichens, cyanobacteria, and heterotrophs living at the soil surface—are fundamental components of drylands worldwide, and destruction of biocrusts dramatically alters biogeochemical processes, hydrology, surface energy balance, and vegetation cover. Although there has been long-standing concern over impacts of physical disturbances on biocrusts (e.g., trampling by livestock, damage from vehicles), there is increasing concern over the potential for climate change to alter biocrust community structure. Using long-term data from the Colorado Plateau, we examined the effects of 10 y of experimental warming and altered precipitation (in full-factorial design) on biocrust communities and compared the effects of altered climate with those of long-term physical disturbance (>10 y of replicated human trampling). Surprisingly, altered climate and physical disturbance treatments had similar effects on biocrust community structure. Warming, altered precipitation frequency [an increase of small (1.2 mm) summer rainfall events], and physical disturbance from trampling all promoted early successional community states marked by dramatic declines in moss cover and increases in cyanobacteria cover, with more variable effects on lichens. Although the pace of community change varied significantly among treatments, our results suggest that multiple aspects of climate change will affect biocrusts to the same degree as physical disturbance. This is particularly disconcerting in the context of warming, as temperatures for drylands are projected to increase beyond those imposed as treatments in our study.

alternate states | biocrusts | community structure | secondary succession | warming

The potential for ecological state transitions in response to global change, particularly transitions that promote feedbacks to terrestrial biogeochemical cycling, is a growing concern (1–6). Anticipating state transitions in terrestrial ecosystems largely hinges on understanding the response of primary producers to warming temperatures, altered precipitation patterns, and novel disturbance regimes (4–7). In arid and semiarid ecosystems (drylands), a substantial portion of primary production can take place in biological soil crusts (biocrusts; Fig. 1) (8, 9), which are communities of mosses, lichens, cyanobacteria, and heterotrophs living at the soil surface that can constitute up to 70% of dryland ground cover (10–12).

Biocrust organisms are adapted to limited moisture and low nutrient conditions and respond rapidly to pulsed, dynamic environmental conditions (13–16). Due to their extensive cover and rapid responses to even small inputs of moisture and nutrients, biocrusts often locally regulate soil hydrology and the cycling of soil carbon (9, 12, 17–21) and nitrogen (9, 22–27). However, the same traits that adapt biocrust organisms to pulsed environmental conditions also make them potentially vulnerable to anticipated changes in climate (28–30). Given their significant influence on ecosystem processes, understanding how biocrust communities will respond to changing climate and disturbance regimes is essential for predicting ecological state changes in drylands—which cover roughly 40% of the Earth's terrestrial surface and hold an estimated 25% of global organic soil carbon (31).

Biocrusts are bound together by filamentous strands of cyanobacteria, which glue soil particles together to form the characteristic soil shields for which the communities are renowned (10). Although this structure results in resistance to wind shear stress, it does not provide much resistance to physical disturbance. Not surprisingly, both the physical and biotic structures of biocrust are highly sensitive to a range of disturbances, such as vehicle traffic and trampling by humans and livestock (32). Disturbance-induced changes in biocrust community structure and subsequent variation across successional states are well-characterized and are remarkably similar globally: Physical disturbance typically transforms later-successional communities of lichens and mosses to early-successional communities dominated by cyanobacteria (10, 32). This successional resetting significantly impacts ecosystem processes, including large changes in carbon and nitrogen cycling (11, 24, 25, 33). Alarming, some experimental work suggests biocrust communities may also be highly sensitive to changing climate. Specifically, increased temperatures have been reported to reduce lichen cover in semiarid Spain (12, 34), and altered precipitation patterns promoted rapid moss mortality (35) and greater variability in cyanobacterial species composition and abundance (36, 37) in a cool desert of the Colorado Plateau. These changes in biocrusts due to climate manipulations also impact ecosystem processes (12, 35, 38).

Despite concern over potential climate change-induced shifts in biocrust community structure and the impacts on ecosystem processes, available reports of biocrust responses to climate manipulations are based on relatively short experimental time spans, often less than 3 y in duration (e.g., 12, 34–36, 39). In addition to the lack of long-term data on biocrust responses to climate change, it is not yet known how climate change impacts on biocrust communities will compare with large, continued

Significance

In drylands worldwide, where plant cover is sparse, large amounts of the ground surface are covered by specialized organisms that form biological soil crusts (biocrusts). Biocrusts fix carbon and nitrogen, stabilize soils, and influence hydrology. Extensive physical disturbance from livestock/human trampling and off-road vehicles is known to destroy biocrusts and alter ecosystem function. More recent work also indicates that climate change can affect biocrust communities. Contrary to our expectations, experimental climate change and physical disturbance had strikingly similar impacts on biocrust communities, with both promoting a shift to degraded, early successional states. These results herald ecological state transitions in drylands as temperatures rise, calling for management strategies that consider risks from both physical disturbances and climate change.

Author contributions: S.C.R. and J.B. designed research; S.C.R. performed research; S.F. contributed new reagents/analytic tools; S.F. analyzed data; and S.F., S.C.R., and J.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. Email: sferrenberg@usgs.gov.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1509150112/-DCSupplemental.




Fig. 1. Biocrusts can locally regulate ecosystem processes and cover large portions of dryland ecosystems as in A (photo by Bill Bowman). Biocrusts are sensitive to physical disturbances from vehicles and trampling by livestock or people as depicted in B, which shows an experimentally trampled plot (foreground) bordered by undisturbed biocrust (background).

threats from physical disturbances due to development, agriculture, and other human activities (32, 40, 41). Understanding the magnitude of threats to biocrusts from both climate change scenarios and novel disturbance regimes is thus a necessary step for predicting future ecological states and developing comprehensive management plans in drylands.

We compared the effects of warming temperatures and altered precipitation patterns on biocrust community structure using data from 10 y (autumn 2005 to autumn 2014) of biocrust community surveys, completed in a full-factorial climate manipulation experiment (control, warming, watering, warming + watering) on the Colorado Plateau, Utah. We simultaneously examined the impact of 15 y annual physical disturbance on biocrust community structure within a replicated human-trampling experiment in a site with vegetation and biocrust communities similar to those in our climate manipulation study (Fig. 1). Finally, we compared the effects of climate manipulation treatments and physical disturbance from trampling on the relative cover of three biotic groups that typically dominate biocrusts of our study region: cyanobacteria, mosses, and lichens. Our goals were, first, to compare the responses of biocrust communities to increased temperature versus increased frequency of small precipitation events, both of which are forecast by climate models for the Colorado Plateau (28, 29, 42) and, second, to compare the effects of climate manipulations to the effects of physical disturbance on biocrust community structure, with a focus on understanding potential variation in responses across different fractions of the community.

Results

Climate manipulation treatments had significant effects on biocrust community structure ($P < 0.001$; Table 1), particularly when the interaction of treatment and time and the interaction of treatment and experimental blocks were considered ($P < 0.001$ for both; Table 1 and Fig. 2). Variance partitioning from principle response curves (PRCs) (43, 44) indicated that interannual variation accounted for 17% of the changes seen in biocrust community structure, whereas treatment \times time interactions accounted for 45% of the variation in community structure. Dunnett's comparison of treatments to the control [completed on principle component analysis (PCA) axis scores] also indicated that watering and warming + watering treatments (intended to represent model predictions of more frequent, small-volume rain events) had rapid effects on community structure, with both treatments leading to a

biocrust community structure significantly different from control plots within 1 y of the start of the experiment ($P < 0.05$; Fig. 2). Warming alone also altered biocrust community structure, but at a slower rate than watering treatments, with warmed and control communities first differing significantly nearly 6 y after the start of the experiment and roughly 3 y after warming treatments were increased from $+2^\circ\text{C}$ to $+4^\circ\text{C}$ ($P < 0.05$; Fig. 2).

Climate manipulations had significant yet variable effects across different fractions of the biocrust community. In particular, warming, watering, and warming + watering all led to a dramatic decrease in relative cover of mosses over time ($P = 0.016$; Table 2 and Fig. 3) and a dramatic increase in the relative cover of cyanobacteria, which filled areas previously occupied by mosses and lichens ($P < 0.0001$; Table 2 and Fig. 3). Lichen responses to treatments varied over time, often mirroring natural fluctuations in lichen relative cover seen in control plots (Fig. 3), but an apparent effect of warming and warming + watering treatments emerged after treatment temperatures were increased from $+2^\circ\text{C}$ to $+4^\circ\text{C}$ (Table 2 and Fig. 3). Mixed-effects modeling indicated that experimental blocks (considered as a random effect) accounted for 15.3% and 13.5% of the variation in the cover of cyanobacteria and mosses, respectively, but accounted for only 1.4% of the variation in lichen cover.


Physical disturbance from human trampling significantly altered biocrust community structure ($P = 0.001$; Table S1), with significant interaction effects among trampling treatment and sample year ($P = 0.009$; Table S1). Similar to climate manipulations, physical disturbance dramatically decreased the relative cover of mosses ($F = 30.4$, $df = 1, 12$, $P = 0.0001$) and lichens ($F = 28.4$, $df = 1, 12$, $P = 0.0002$), while increasing the cover of cyanobacteria, which filled in areas where mosses and lichens declined ($F = 31.4$, $df = 1, 12$, $P = 0.0001$). Overall, mosses and lichens comprised $\leq 0.5\%$ of cover in biocrust communities in trampled plots, compared with 18% and 1.5% of cover in biocrust communities, respectively, in undisturbed control plots. Cyanobacteria cover increased from an average of 81% of the community in control plots to an average of 99% of the community in trampled plots. Mixed-effects modeling indicated that experimental blocks accounted for $<1\%$ of the variation in the cover of cyanobacteria and mosses across the trampling experiment but accounted for 27% of the variation in lichen cover.

Comparing the long-term effects from physical disturbance (community survey data from 11 and 15 y after the start of trampling) to those from climate manipulation treatments (community survey data from 10 y after the start of warming and watering treatments) revealed similar treatment impacts on the cover of cyanobacteria, mosses, and lichens (Fig. 4). In all cases, treatments drastically reduced moss cover and dramatically increased cyanobacteria cover in soil crusts (Fig. 4). Climate manipulations and

Table 1. Two-way PERMANOVA test of the effects of climate manipulation treatments and time (year of experiment) on biocrust community structure (2005–2014)

Source*	df	SS	Pseudo-F	P
Treatment	3	64,912	11.5	0.001
Block	4	7,820	18.0	0.001
Time	18	43,097	13.6	0.001
Treatment \times block	12	22,609	17.4	0.001
Treatment \times time	54	20,396	3.5	0.001
Block \times time	72	12,693	1.6	0.001
Residual	216	23,407	—	—


*First biocrust surveys were completed in Autumn 2005, and treatments were initiated in December 2005 and included controls (no climate manipulations), warming, watering (frequent, small-volume water additions), and warming + watering. SS, sums of squares from PERMANOVA.



We found evidence of substantial block (landscape) effects on both community composition and the responses of specific biotic groups to treatments across experiments. These block effects suggest the existence of small-scale gradients in climate and disturbance impacts on biocrust communities. These gradients, possibly in soil nutrients or microclimate, may interact with interannual climatic variation such that areas prone to fluctuations in soil moisture or temperature may become more or less harsh for biocrust organisms over time (51–53). Environmental gradients can also affect the strength and direction of biotic interactions within these communities (47, 54), highlighting the need to consider spatial and temporal influences on species responses in environmental change studies.

Conclusions

We examined the effects of experimental warming and augmented precipitation on biocrust communities and then compared the impacts of climate manipulations to physical disturbance (replicated human trampling) on biocrust community structure. We found that warming, watering, warming + watering, and physical disturbance



had similar effects on biocrust communities, collectively shifting communities toward early successional states. Although physical disturbances can be mitigated (41) and future precipitation patterns are uncertain (28, 42), increasing temperatures (meeting or exceeding those imposed by warming treatments) are considered a near certainty across many dryland ecosystems (29). Thus, our results suggest that climate change will affect biocrusts to an extent similar to changes observed with physical disturbance, leading to a landscape-level shift to early successional states with seemingly limited potential for a return to late successional states. However, given the different rates at which these pressures alter biocrust community structure and the potential for longer term adaptation and species' shifts, variation in the ultimate effects of these disturbances on ecosystem processes are likely and require continued study.

Ecological state changes in response to global change factors are a growing concern. Transitions to alternate community states in drylands are a particularly poignant issue, as these ecosystems cover roughly 40% of the Earth's terrestrial surfaces and may hold up to 25% of global soil organic carbon (31). Biocrust communities can represent a substantial portion of primary production in drylands and can locally stabilize and regulate ecosystem processes including carbon and nitrogen cycles as well as short-term hydrologic cycles and surface energy balance (8, 9, 45, 62–64). The role biocrusts play in soil stability is also a topic of significant public interest, due to the human health and safety issues associated with increasing dust storms. Because different components of the biocrust community (i.e., mosses versus lichens) play different roles in the functioning of drylands (8, 10, 56, 65, 66), understanding the factors that control the composition of biocrust communities will be essential to predictions of future dryland function. In addition, because drylands may dominate the interannual variability in global atmospheric CO₂ concentrations (67, 68), state transitions in dryland structure could have effects beyond arid and semiarid systems.

Methods

Climate manipulation and physical disturbance experiments were located on the Upper Colorado Plateau, Utah. The region is classified as cool desert with a majority of precipitation occurring in winter and spring. The sites have extensive cover of biocrusts held together primarily by filaments of the cyanobacteria *Microcoleus vaginatus* and covered by a mosaic of the mosses *Syntrichia caninervis* and *Syntrichia ruralis* and various lichens (mostly *Collema tenax* and *Collema coccophorum*). Climate manipulations representing model predictions of both increasing temperature and frequency of small (<2 mm) rainfall events for the study region (28, 29, 42) were completed near Castle Valley, Utah (38.6748 N, –109.4163 W, 1,310 m above sea level). The soil of this area is Lithic Torriorthent, a sandy-loam ranging in depth from 17 to 122 cm. Human trampling plots, mimicking disturbances from humans and livestock, were located in Arches National Park, Utah, on soil of the Mido-Sazi complex, a loamy sand with depths to 100 cm.

Climate manipulations began in autumn 2005 in a randomized, complete block design consisting of twenty 2 m × 2.5 m plots divided among four treatments: control (no manipulations), warmed, watered, and warmed + watered. Warming was completed with infrared lamps heating the topsoil (temperature at 1–2 cm depth) by +2 °C above ambient for the first 3 y of

the experiment and then by +4 °C above ambient from June 30, 2008 until present. In addition to ambient precipitation, watering treatments consisted of frequent, 1.2 mm “rainfall events” applied by hand sprayers. Watered plots received on average 35 such events (roughly four times the average natural frequency) throughout the summer months starting in 2006 (35–37, 39). Physical disturbance by human trampling started in May 1996 in ten 2 m × 5 m plots, divided as five trampled and five undisturbed controls. Plots were trampled annually in May of successive years until 2011 (69). Biocrust communities were assessed as the relative cover of all visible species present within standardized 40 cm × 40 cm frames. Climate manipulation plots were surveyed twice per year (May and September) from Autumn 2005 to present, with survey frames placed in four repeated locations per plot (6,400 cm² surveyed in each plot in a repeated-measures design). Trampled–control plot pairs were surveyed in 2007 and 2011, with survey frames placed in 10 random locations within each plot (16,000 cm² surveyed in each plot).

Biocrust community responses to climate manipulations and trampling were assessed via permutational multivariate analysis of variance (PERMANOVA). PERMANOVAs were performed on Bray–Curtis distance matrices of log-transformed species' abundances derived from cover measures. PERMANOVAs on climate manipulation data included treatment, block, and time since the start of the experiment as factors, and PERMANOVAs on physical disturbance data included treatment and biocrust survey year (2007 and 2011) as factors, with block effects ruled out by stepwise fitting procedures. Treatment effects of climate manipulations were visualized via PRCs, which measure the differences between the species composition and cover in each treatment compared with controls over time (43, 70). To determine the date when biocrust communities in climate manipulation treatments first differed from controls, the community structure of treatment and the control plots were compared via Dunnett's tests on PCA axis scores. PRC and PCA were completed on log-transformed species abundance data. To determine effects of climate and disturbance treatments on different fractions of the biocrust community, we calculated relative abundances for three biotic groups common to biocrusts of our sites (mosses, lichens, and cyanobacteria). To account for the repeated-measures design of biocrust surveys, relative cover of the three biotic groups were analyzed with linear mixed-effect models with restricted maximum likelihood estimators. Treatment and time (or sample years in the case of the trampling experiment) were considered fixed effects, and field experimental blocks were considered as random effects. Finally, we compared the long-term effects of climate manipulations (data from the 10th year of the study) and physical disturbance (data from the 11th and 15th year of the study when biocrusts were surveyed) on the relative cover of mosses, lichens, and cyanobacteria by relativizing treatment responses to controls in a manner accounting for cover values in both treatments and controls: Relative Cover Change Index, or RCCI = $\frac{\text{cover}_{(\text{treatment})} - \text{cover}_{(\text{control})}}{\text{cover}_{(\text{treatment})} + \text{cover}_{(\text{control})}}$ (71). RCCI values within biotic groups were then compared via Kruskal–Wallis one-way analysis of variance followed by Steel–Dwass nonparametric pairwise comparisons (72). Analyses were completed in R (72).

ACKNOWLEDGMENTS. We thank Hilda Smith, Ed Grote, and many technicians for data collection and maintaining experiments. We are grateful to the National Park Service Southeast Utah Group and the Bureau of Land Management Moab Field Office for help with permits and logistics. Our manuscript was improved by comments from Peter Vitousek, Jeffery Mitton, and three anonymous reviewers. This work was supported by the US Department of Energy Office of Science, Office of Biological and Environmental Research Terrestrial Ecosystem Sciences Program, under Award DE-SC-0008168, and the USGS Climate and Land Use, and Ecosystems programs. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

- Friedlingstein P, et al. (2001) Positive feedback between future climate change and the carbon cycle. *Geophys Res Lett* 28(8):1543–1546.
- Scheffer M, Carpenter S, Foley JA, Folke C, Walker B (2001) Catastrophic shifts in ecosystems. *Nature* 413(6856):591–596.
- Field CB, Lobell DB, Peters HA, Chiariello NR (2007) Feedbacks of terrestrial ecosystems to climate change. *Annu Rev Environ Resour* 32:1–29.
- Luo Y (2007) Terrestrial carbon-cycle feedback to climate warming. *Annu Rev Ecol Syst* 38:683–712.
- Turner MG (2010) Disturbance and landscape dynamics in a changing world. *Ecology* 91(10):2833–2849.
- Zhou J, et al. (2012) Microbial mediation of carbon-cycle feedbacks to climate warming. *Nat Clim Chang* 2(2):106–110.
- Lindenmayer DB, Likens GE, Krebs CJ, Hobbs RJ (2010) Improved probability of detection of ecological “surprises”. *Proc Natl Acad Sci USA* 107(51):21957–21962.
- Yoshitake S, Uchida M, Koizumi H, Kanda H, Nakatsubo T (2010) Production of biological soil crusts in the early stage of primary succession on a high Arctic glacier foreland. *New Phytol* 186(2):451–460.
- Elbert W, et al. (2012) Contribution of cryptogamic covers to the global cycles of carbon and nitrogen. *Nat Geosci* 5(7):459–462.
- Belnap J, Büdel B, Lange OL (2003a) Biological soil crusts: Characteristics and distribution. *Biological Soil Crusts: Structure, Function, and Management*, eds Belnap J, Lange OL (Springer-Verlag, Berlin), pp 3–30.
- Housman DC, Powers HH, Collins AD, Belnap J (2006) Carbon and nitrogen fixation differ between successional stages of biological soil crusts in the Colorado Plateau and Chihuahuan Desert. *J Arid Environ* 66(4):620–634.
- Maestre FT, et al. (2013) Changes in biocrust cover drive carbon cycle responses to climate change in drylands. *Glob Change Biol* 19(12):3835–3847.
- Austin AT, et al. (2004) Water pulses and biogeochemical cycles in arid and semiarid ecosystems. *Oecologia* 141(2):221–235.

14. Cable JM, Huxman TE (2004) Precipitation pulse size effects on Sonoran Desert soil microbial crusts. *Oecologia* 141(2):317–324.
15. Huxman TE, et al. (2004) Precipitation pulses and carbon fluxes in semiarid and arid ecosystems. *Oecologia* 141(2):254–268.
16. Schwinning S, Sala OE (2004) Hierarchy of responses to resource pulses in arid and semi-arid ecosystems. *Oecologia* 141(2):211–220.
17. Wilske B, et al. (2008) The CO₂ exchange of biological soil crusts in a semiarid grass-shrubland at the northern transition zone of the Negev desert, Israel. *Biogeosciences* 5(3):1411–1423.
18. Wilske B, et al. (2009) Modeling the variability in annual carbon fluxes related to biological soil crusts in a Mediterranean shrubland. *Biogeosciences Discuss* 6(4):7295–7324.
19. Bowker MA, Mau RL, Maestre FT, Escobar C, Castillo-Monroy AP (2011) Functional profiles reveal unique ecological roles of various biological soil crust organisms. *Funct Ecol* 25(4):787–795.
20. Castillo-Monroy AP, et al. (2011) Relationships between biological soil crusts, bacterial diversity and abundance, and ecosystem functioning: Insights from a semi-arid Mediterranean environment. *J Veg Sci* 22(1):165–174.
21. Miralles I, Trasac-Cepeda C, Leirós MC, Gil-Sotres F (2013) Labile carbon in biological soil crusts in the Tabernas desert, SE Spain. *Soil Biol Biochem* 58:1–8.
22. Belnap J (2002) Impacts of off-road vehicles on nitrogen cycles in biological soil crusts: Resistance in different U.S. deserts. *J Arid Environ* 52(2):155–165.
23. Hawkes CV (2003) Nitrogen cycling mediated by biological soil crusts and arbuscular mycorrhizal fungi. *Ecology* 84(6):1553–1562.
24. Barger NN, Belnap J, Ojima DS, Mosier A (2005) NO gas loss from biologically crusted soils in Canyonlands National Park, Utah. *Biogeochemistry* 75(3):373–391.
25. Barger NN, Herrick JE, Van Zee JW, Belnap J (2006) Impacts of biological soil crust disturbance and composition on C and N loss from water erosion. *Biogeochemistry* 77(2):247–263.
26. Castillo-Monroy AP, Maestre FT, Delgado-Baquerizo M, Gallardo A (2010) Biological soil crusts modulate nitrogen availability in semi-arid ecosystems: Insights from a Mediterranean grassland. *Plant Soil* 333(1–2):21–34.
27. Delgado-Baquerizo M, Castillo-Monroy AP, Maestre FT, Gallardo A (2010) Plants and biological soil crusts modulate the dominance of N forms in a semi-arid grassland. *Soil Biol Biochem* 42(2):376–378.
28. Christensen NS, Wood AW, Voisin N, Lettenmaier DP, Palmer RN (2004) The effects of climate change on the hydrology and water resources of the Colorado River basin. *Clim Change* 62(1–3):337–363.
29. Solomon S, et al., eds (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge Univ Press, New York).
30. Cayan DR, et al. (2010) Future dryness in the southwest US and the hydrology of the early 21st century drought. *Proc Natl Acad Sci USA* 107(50):21271–21276.
31. Safriel U, Adeel Z (2005) Dryland systems. *Ecosystems and Human Well-Being, Current State and Trends*, eds Hassan R, Scholes R, Ash N (Island Press, Washington, DC), Vol 1, pp 625–658.
32. Belnap J, Eldridge D (2003) Disturbance and recovery of biological soil crusts. *Biological Soil Crusts: Structure, Function, and Management*, eds Belnap J, Lange OL (Springer-Verlag, Berlin), pp 363–383.
33. Zaady E, Kuhn U, Wilske B, Sandoval-Soto L, Kesselmeier J (2000) Patterns of CO₂ exchange in biological soil crusts of successional age. *Soil Biol Biochem* 32(7):959–966.
34. Escobar C, Martínez I, Bowker MA, Maestre FT (2012) Warming reduces the growth and diversity of biological soil crusts in a semi-arid environment: Implications for ecosystem structure and functioning. *Philos Trans R Soc Lond B Biol Sci* 367(1606):3087–3099.
35. Reed SC, et al. (2012) Changes to dryland rainfall result in rapid moss mortality and altered soil fertility. *Nat Clim Chang* 2(10):752–755.
36. Johnson SL, et al. (2012) Increased temperature and altered summer precipitation have differential effects on biological soil crusts in a dryland ecosystem. *Glob Change Biol* 18(8):2583–2593.
37. Yeager CM, et al. (2012) Response of biological soil crust diazotrophs to season, altered summer precipitation, and year-round increased temperature in an arid grassland of the Colorado plateau, USA. *Front Microbiol* 3:358.
38. Escobar C, Maestre FT, Rey A (2015) Biocrusts modulate warming and rainfall exclusion effects on soil respiration in a semi-arid grassland. *Soil Biol Biochem* 80:9–17.
39. Zelikova TJ, Housman DC, Grote EE, Neher DA, Belnap J (2012) Warming and increased precipitation frequency on the Colorado Plateau: Implications for biological soil crusts and soil processes. *Plant Soil* 355(1–2):265–282.
40. Reynolds JF, et al. (2007) Global desertification: Building a science for dryland development. *Science* 316(5826):847–851.
41. Bestelmeyer BT, et al. (2015) Desertification, land use, and the transformation of global drylands. *Front Ecol Environ* 13(1):28–36.
42. Weltzin JF, et al. (2003) Assessing the response of terrestrial ecosystems to potential changes in precipitation. *Bioscience* 53(10):941–952.
43. Van den Brink PJ, Ter Braak CJF (1999) Principal response curves: Analysis of time-dependent multivariate responses of biological community to stress. *Environ Toxicol Chem* 18(2):138–148.
44. Legendre P, Legendre L (1998) *Numerical Ecology* (Elsevier, Amsterdam), 2nd English Ed.
45. Warren SD, Eldridge DJ (2003) Biological soil crusts and livestock in arid ecosystems: Are they compatible? *Biological Soil Crusts: Structure, Function, and Management*, eds Belnap J, Lange OL (Springer-Verlag, Berlin), pp 401–415.
46. Garfin G, Jardine A, Merideth R, Black M, LeRoy S (2013) *Assessment of Climate Change in the Southwest United States: A Report Prepared for the National Climate Assessment. A Report by the Southwest Climate Alliance* (Island Press, Washington, DC).
47. Bowker MA, Soliveres S, Maestre FT (2010) Competition increases with abiotic stress and regulates the diversity of biological soil crusts. *J Ecol* 98(3):551–560.
48. de Vries FT, Shade A (2013) Controls on soil microbial community stability under climate change. *Front Microbiol* 4:265.
49. Nemergut DR, et al. (2013) Patterns and processes of microbial community assembly. *Microbiol Mol Biol Rev* 77(3):342–356.
50. Wertin TM, et al. (2012) Elevated CO₂ did not mitigate the effect of a short-term drought on biological soil crusts. *Biol Fertil Soils* 48(7):797–805.
51. Bowker MA, Belnap J (2008) A simple classification of soil types as habitats of biological soil crusts on the Colorado Plateau, USA. *J Veg Sci* 19(6):831–840.
52. Bowker MA, Belnap J, Davidson DW, Phillips SL (2005) Evidence for micronutrient limitation of biological soil crusts: Importance to arid-lands restoration. *Ecol Appl* 15(6):1941–1951.
53. Cable JM, Ogle K, Williams DG, Weltzin JF, Huxman TE (2008) Soil texture drives responses of soil respiration to precipitation pulses in the Sonoran Desert: Implications for climate change. *Ecosystems (N Y)* 11(6):961–979.
54. Maestre FT, et al. (2010) Do biotic interactions modulate ecosystem functioning along stress gradients? Insights from semi-arid plant and biological soil crust communities. *Philos Trans R Soc Lond B Biol Sci* 365(1549):2057–2070.
55. Evans RD, Belnap J (1999) Long-term consequences of disturbance on nitrogen dynamics in an arid ecosystem. *Ecology* 80(1):150–160.
56. Belnap J, Hawkes CV, Firestone MK (2003b) Boundaries in miniature: Two examples from soil. *Bioscience* 53(8):739–749.
57. McCalley CK, Sparks JP (2009) Abiotic gas formation drives nitrogen loss from a desert ecosystem. *Science* 326(5954):837–840.
58. Jackson LE, Schimel JP, Firestone MK (1989) Short-term partitioning of ammonium and nitrate between plants and microbes in an annual grassland. *Soil Biol Biochem* 21(3):409–415.
59. Austin AT, Sala OE, Jackson RB (2006) Inhibition of nitrification alters carbon turnover in the Patagonian steppe. *Ecosystems* 9(8):1257–1265.
60. Maestre FT, Huesca M, Zaady E, Bautista S, Cortina J (2002) Infiltration, penetration resistance and microphytic crust composition in contrasted microsites within a Mediterranean semi-arid steppe. *Soil Biol Biochem* 34(6):895–898.
61. Chamizo S, Canton Y, Lázaro R, Solé-Benet A, Domingo F (2012) Crust composition and disturbance drive infiltration through biological soil crusts in semiarid ecosystems. *Ecosystems* 15(1):148–161.
62. Evans RD, Lange OL (2003) Biological soil crusts and ecosystem nitrogen and carbon dynamics. *Biological Soil Crusts: Structure, Function, and Management*, eds Belnap J, Lange OL (Springer-Verlag, Berlin), pp 401–415.
63. Delgado-Baquerizo M, Maestre FT, Gallardo A (2013) Biological soil crusts increase the resistance of soil nitrogen dynamics to changes in temperatures in a semi-arid ecosystem. *Plant Soil* 366(1–2):35–47.
64. Wei W, Yu Y, Chen L (2015) Response of surface soil hydrology to the micro-pattern of bio-crust in a dry-land Loess environment, China. *PLoS One* 10(7):e0133565.
65. Holst J, et al. (2009) Dinitrogen fixation by biological soil crusts in an Inner Mongolian steppe. *Biol Fertil Soils* 45(7):679–690.
66. Zhang P, et al. (2012) [Nitrogen fixation potential of biological soil crusts in southeast edge of Tengger Desert, Northwest China]. *Ying Yong Sheng Tai Xue Bao* 23(8):2157–2164.
67. Poulter B, et al. (2014) Contribution of semi-arid ecosystems to interannual variability of the global carbon cycle. *Nature* 509(7502):600–603.
68. Ahlström A, et al. (2015) Carbon cycle. The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink. *Science* 348(6237):895–899.
69. Kuske CR, Yeager CM, Johnson S, Ticknor LO, Belnap J (2012) Response and resilience of soil biocrust bacterial communities to chronic physical disturbance in arid shrublands. *ISME J* 6(4):886–897.
70. Frampton GK, Van den Brink PJ, Gould PJL (2000) Effects of spring precipitation on a temperate arable collembolan community analysed using Principal Response Curves. *Appl Soil Ecol* 14(3):231–248.
71. Armas C, Ordiales R, Pugnaire FI (2004) Measuring plant interactions: A new comparative index. *Ecology* 85(10):2682–2686.
72. R Development Core Team (2011) *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria), www.r-project.org.