

Soil Erosion Potential

Introduction

AIM uses the soil stability of (*macro*)aggregates as an indicator of the potential of soil to erosion (soil erodibility) (J. E. Herrick et al. (2021), J. Herrick et al. (2001)). Soil aggregates are groups of soil particles which clump together to form individual strongly connected (micro)aggregates, which may then continually clump together with each other and organic matter to form larger less weakly connected (macro)aggregates (Totsche et al. (2018)). High numbers of macroaggregates at the soil surface have been shown to correlate strongly to the effects of rain, and wind, on the loss of soil from areas (Barthes & Roose (2002), J. Herrick et al. (2001)). While the relationships between soil erosion, plant cover and functional type (Cerdeira (1998), Greene et al. (1994), Torre-Robles et al. (2023)), landform position (Swanson et al. (1988), Torre-Robles et al. (2023)), slope shape (e.g. concave, convex) (Canton et al. (2009), Torre-Robles et al. (2023)), and the cover of biocrusts are oftentimes complex (Leys & Eldridge (1998)), the quantitative observation that soils with low macroaggregate stability have greater rates of erodibility are always evident (J. Herrick et al. (2018)).

For this report, soil aggregates refer to relatively large coherent portions of soils ($> 2\text{mm}$ diameter) (J. E. Herrick et al. (2021)). Soil microaggregates are continually being slowly created by processes, such as the initial attraction of negatively charged clay and positively charged salts on silt particles, followed by cementation (Totsche et al. (2018)). Cementation is often achieved through organic matter and/or calcium carbonates and oxides, which then leads to biological processes. These involve the creation of numerous long organic (Carbon containing) molecules (generally polysaccharides), by organisms such as bacteria especially filamentous cyanobacteria, fungal hyphae, and plant roots, which act as ‘glue’ between these particles and will create macroaggregates (Six et al. (2004), Totsche et al. (2018), Moonilall (n.d.)). Soil macroaggregates in non agricultural lands are continually quickly formed, and subsequently broken back into modified microaggregates by: certain wildland fire conditions (Urbanek (2013)), rapid drying and wetting, freeze-thaw cycles, some chemical interactions with water, and compaction (Le Bissonnais (1996)). When many more soil aggregates are being broken apart than are created areas become susceptible to erosion from water, or wind (Leys & Eldridge (1998), Six et al. (2004)).

Soil erosion decreases water infiltration to soil and becomes available to plants, reduces soil nutrients available to plants and microorganisms, removes soil carbon which foster soil microorganisms, and decrease root depth and space for plants; all leading to decreases in plant diversity, abundance and production (reviewed in Pimentel & Kounang (1998)). Accordingly, soil erosion may lead to the inability of an Ecological Site to support certain plant species essential to the maintenance of the site Bestelmeyer et al. (2015). In most instances this will tend to lead a different *state* and *phase*, generally with fewer ecosystem occurrences, to occur on the site (Bestelmeyer et al. (2015)). However, in severe instances soil erosion will lead to conversion of a site into a state from which land management agencies are unlikely to be capable to restore it to an area providing even few of it’s potential ecosystem services (Bestelmeyer et al. (2015)). Realistically in nearly all semi-arid lands utilized as rangelands, this equates to desertification.

... Current concerns regarding soil stability are to be compounded with climate change (Munson et al. (2011)), soil crusts, perhaps with the exception of ‘light’ cyanobacteria, are slow to regenerate. More episodic, and intense rainfalls, badddd. (Chen et al. (2018))

Soil stability will be one of the only core-indicators in this report that is treated as a *categorical* variable. The way that crews collect soil stability means that it is an ordinal categorical variable, i.e. an object with discrete categories which are ordered. Soil stability measures are on a scale of 1-6, where ‘1’ indicates little to no stability and ‘6’ indicates very high stability. While it is tempting to treat these values as *continuous*, it

is generally inappropriate to do so. These values represent data which are *ordinal categorical variables*, and while these types of information *can* readily be, and are oftentimes, treated as continuous values they should meet a few conditions first. One condition where ordinal variables can be transformed is when the values they represent are at equal distances from each other, *e.g.* if we estimate the number of number of jelly beans in a jar from 1-10, where 1 encompasses 1-10 beans, and 2 21-30 beans etc., However, our rankings of 0-6 all show different ‘distances’ between them. For example:

“Stability class 4: 10–25% of soil remains on sieve after five dipping cycles;
 Stability class 5: 25–75% of soil remains on sieve after five dipping cycles” — AIM 2021, V. 1
 p. 51

As can readily be seen, from these two classes which are the most similar, breaks are of wildly different sizes (15% and 50%), and clearly violate this assumption. Another condition where ordinal values can be treated as continuous is when they represent a great range. E.g. if you had a quiz of 100 questions, and you can get each answer either ‘Correct’, or ‘Wrong’, we could turn your score into a continuous value, and combine it with other scores to get an aggregate. Lenient opinions differ on whether you should have 10+ or 5+ classes to do this (CITE). Finally, we have few replicates per site. Soil stability is measured at only 18 locations per plot, roughly only half of the recommended observations for using parametric statistics. While non-parametric statistics are often applied to numeric data, they perform very well with small samples sizes. Accordingly, we end up in relatively the same place statistically by treating these values as ordinal categorical variables.

Methods

The first step in assessing whether the field office was achieving benchmark conditions regarding soil stability was to impute the measurements for these values at Ecological Sites which were lacking Descriptions, or which had incomplete descriptions. Similar to section (X.X), these values were imputed by *feature engineering*, however since they were ordinal categorical variables the *median* of the values were used.

A relatively high amount of Ecological Sites Descriptions (33 of 52, 63.5%) contained soil stability reference benchmarks under two conditions as well as a site ‘average’:

- 1) Interspaces (the distance between plant canopies)
- 2) Under Canopy (space beneath plant cover)
- 3) Site ‘average’ (hereafter: median)

A hand-full of sites ($n = 7$) contained values for both under canopies and interspaces, but lacked a site wide median; a dozen sites had only one value ($n = 9$ ‘site’, and $n = 3$ ‘interspaces’). To calculate these estimates for ESD’s which were missing them, the median for each category was gathered using each observation (Figure 1, Table 1). The missing values were then imputed in each ESD which was missing a value, as well as for each ES which was missing all values.




Figure 1: Specified benchmarks at all ESDs which included all three metrics and derived imputed value

Results

References

- Barthes, B., & Roose, E. (2002). Aggregate stability as an indicator of soil susceptibility to runoff and erosion; validation at several levels. *Catena*, 47(2), 133–149.
- Bestelmeyer, B. T., Okin, G. S., Duniway, M. C., Archer, S. R., Sayre, N. F., Williamson, J. C., & Herrick, J. E. (2015). Desertification, land use, and the transformation of global drylands. *Frontiers in Ecology and the Environment*, 13(1), 28–36.
- Canton, Y., Sole-Benet, A., Asensio, C., Chamizo, S., & Puigdefabregas, J. (2009). Aggregate stability in range sandy loam soils relationships with runoff and erosion. *Catena*, 77(3), 192–199.
- Cerda, A. (1998). Soil aggregate stability under different mediterranean vegetation types. *Catena*, 32(2), 73–86.
- Chen, H., Zhang, X., Abia, M., Lu, D., Yan, R., Ren, Q., Ren, Z., Yang, Y., Zhao, W., Lin, P., et al. (2018). Effects of vegetation and rainfall types on surface runoff and soil erosion on steep slopes on the loess plateau, china. *Catena*, 170, 141–149.
- Greene, R., Kinnell, P., & Wood, J. T. (1994). Role of plant cover and stock trampling on runoff and soil-erosion from semi-arid wooded rangelands. *Soil Research*, 32(5), 953–973.
- Herrick, J. E., Van Zee, J. W., McCord, C., Sarah E. and, Karl, J. W., & Burkett, L. M. (2021). *Monitoring manual for grassland, shrubland, and savanna ecosystems* (2nd ed., Vol. 1). United States Department of Agriculture, Agricultural Research Services, Jornada Experimental Range.
- Herrick, J., Weltz, M., Reeder, J., Schuman, C., & Simanton, J. (2018). Rangeland soil erosion and soil quality: Role of soil resistance, resilience, and disturbance regime. In *Soil quality and soil erosion* (pp. 209–233). CRC press.
- Herrick, J., Whitford, W., De Soyza, A., Van Zee, J., Havstad, K., Seybold, C., & Walton, M. (2001). Field soil aggregate stability kit for soil quality and rangeland health evaluations. *Catena*, 44(1), 27–35.
- Le Bissonnais, Y. le. (1996). Aggregate stability and assessment of soil crustability and erodibility: I. Theory and methodology. *European Journal of Soil Science*, 47(4), 425–437.
- Leys, J. F., & Eldridge, D. J. (1998). Influence of cryptogamic crust disturbance to wind erosion on sand and loam rangeland soils. *Earth Surface Processes and Landforms: The Journal of the British Geomorphological Group*, 23(11), 963–974.
- Moonilall, N. I. (n.d.). *What are soil aggregates?* wordpress. <https://soilsmatter.wordpress.com/2019/07/15/what-are-soil-aggregates/>
- Munson, S. M., Belnap, J., & Okin, G. S. (2011). Responses of wind erosion to climate-induced vegetation changes on the colorado plateau. *Proceedings of the National Academy of Sciences*, 108(10), 3854–3859.
- Pimentel, D., & Kounang, N. (1998). Ecology of soil erosion in ecosystems. *Ecosystems*, 1(5), 416–426.
- Six, J., Bossuyt, H., Degryze, S., & Denef, K. (2004). A history of research on the link between (micro) aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research*, 79(1), 7–31.
- Swanson, F., Kratz, T., Caine, N., & Woodmansee, R. (1988). Landform effects on ecosystem patterns and processes. *BioScience*, 38(2), 92–98.
- Torre-Robles, L. de la, Munoz-Robles, C., Huber-Sannwald, E., & Reyes-Aguero, J. A. (2023). Functional stability: From soil aggregates to landscape scale in a region severely affected by gully erosion in semi-arid central mexico. *CATENA*, 222, 106864.
- Totsche, K. U., Amelung, W., Gerzabek, M. H., Guggenberger, G., Klumpp, E., Knief, C., Lehndorff, E., Mikutta, R., Peth, S., Prechtel, A., et al. (2018). Microaggregates in soils. *Journal of Plant Nutrition and Soil Science*, 181(1), 104–136.
- Urbanek, E. (2013). Why are aggregates destroyed in low intensity fire? *Plant and Soil*, 362(1), 33–36.