

1 Inequality Practice

- (a) X is a random variable such that $X > -5$ and $\mathbb{E}[X] = -3$. Find an upper bound for the probability of X being greater than or equal to -1 .
- (b) Y is a random variable such that $Y < 10$ and $\mathbb{E}[Y] = 1$. Find an upper bound for the probability of Y being less than or equal to -1 .
- (c) You roll a die 100 times. Let Z be the sum of the numbers that appear on the die throughout the 100 rolls. Compute $\text{Var}(Z)$. Then use Chebyshev's inequality to bound the probability of the sum Z being greater than 400 or less than 300.

2 Vegas

On the planet Vegas, everyone carries a coin. Many people are honest and carry a fair coin (heads on one side and tails on the other), but a fraction p of them cheat and carry a trick coin with heads on both sides. You want to estimate p with the following experiment: you pick a random sample of n people and ask each one to flip his or her coin. Assume that each person is independently likely to carry a fair or a trick coin.

1. Given the results of your experiment, how should you estimate p ?
(*Hint:* Construct an (unbiased) estimator for p such that $E[\hat{p}] = p$.)

2. How many people do you need to ask to be 95% sure that your answer is off by at most 0.05?

3 Working with the Law of Large Numbers

- (a) A fair coin is tossed multiple times and you win a prize if there are more than 60% heads. Which number of tosses would you prefer: 10 tosses or 100 tosses? Explain.

- (b) A fair coin is tossed multiple times and you win a prize if there are more than 40% heads. Which number of tosses would you prefer: 10 tosses or 100 tosses? Explain.

- (c) A fair coin is tossed multiple times and you win a prize if there are between 40% and 60% heads. Which number of tosses would you prefer: 10 tosses or 100 tosses? Explain.

- (d) A fair coin is tossed multiple times and you win a prize if there are exactly 50% heads. Which number of tosses would you prefer: 10 tosses or 100 tosses? Explain.