

Discrete Time Crystal: The Vanilla Unitary as a Brickwall Circuit

Sagnik Ghosh

1. Model

The model we will be concerned with is the brick-wall circuit from Ipolitti et al [1, 2]. The General Floquet Model considered here, that exhibits a Discrete Time Crystal in 1D is an disordered Ising model with periodic $\pi/2$ Kicks about the x -axis. The system is probed only at stroboscopic times, $t = n\tau$, and $\tau = 1$

$$(1) \quad U_F = e^{-ig \sum_i X_i} e^{-i\tau(\sum_i J_i Z_i Z_{i+1} + H_{int})} e^{-i\tau(\sum_i h_i Z_i)}$$

Here we have adopted the standard information theoretic notation (i.e have denoted Pauli operator $\sigma_i^Z = Z_i$ etc.) The H_{int} denotes some generic interaction such as fields or coupling,

$$H_{int} = \frac{\theta}{2} \sum_i [X_i X_{i+1} + Y_i Y_{i+1}] \text{(XY Coupling)}$$




FIGURE 1. Schematic representation of the circuit and the associated phase diagram. The circuit exhibit three phases. For smaller values of θ and ϵ close to 1, that is g close to 0, it can support a MBL phase which is paramagnetic in nature. For small θ and ϵ close to 0, that is g close to 1, it supports a DTC MBL phase with a period multiplicity of 2. In the middle region the system tend to scramble. The phase boundary was extracted from results of [1], and was obtained from finite size scaling of the disorder averaged level spacing ratio.

Among these parameters J_i, h_i are disordered and are uniformly sampled from $[0, \frac{\pi}{2}]$ to ensure localisation and thereby to prevent the system from heating up to triviality from the kicks [3]. In absence of interaction and with perfect kicks it is trivial to see that operators of the form $\langle Z(0), Z(n) \rangle$ breaks the discrete time symmetry and exhibits a period doubling. The non-triviality of the DTC as a *phase* comes from the fact that this feature persists strongly even in the presence of interactions as well as with imperfect kicks (i.e $g = \frac{1}{2} = (\pi - \epsilon)$ with finite but small ϵ).

In a quantum device this is realised as a brick-wall circuit.

We also add further small uncertainties in the realisations of the two body gates

$$(2) \quad \theta = [\bar{\theta} - \frac{\Delta\theta}{2}, \bar{\theta} + \frac{\Delta\theta}{2}]$$

where $\Delta\theta = \frac{\pi}{50}$ and random single body Z gates, $RZ(\Delta h)$, to both the legs before and after the application of the two body gates. $\Delta h = \frac{\pi}{50}$.

References

- ¹M. Ippoliti, K. Kechedzhi, R. Moessner, S. Sondhi, and V. Khemani, “Many-body physics in the nisq era: quantum programming a discrete time crystal”, PRX Quantum **2**, 030346 (2021) 10.1103/PRXQuantum.2.030346.
- ²X. Mi, M. Ippoliti, C. Quintana, A. Greene, Z. Chen, J. Gross, F. Arute, K. Arya, J. Atalaya, R. Babbush, et al., “Time-crystalline eigenstate order on a quantum processor”, Nature **601**, 531–536 (2022).
- ³V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, “Phase structure of driven quantum systems”, Phys. Rev. Lett. **116**, 250401 (2016) 10.1103/PhysRevLett.116.250401.

PHYSIKALISCHES INSTITUT DER UNIVERSITÄT BONN, NUSSALLEE 12, 53115 BONN, GERMANY