

Closed Subsets

1 Closed Subsets

Let X be a metric space. A subset E of X is *closed* if its complement $X \setminus E$ is open.

Example 1.1. In any metric space X , the sets \emptyset and X are always closed, because their complements are X and \emptyset , respectively, and we have already noted that these are always open. In particular, both the sets \emptyset and X are both open *and* closed!

Example 1.2. Let $E := \{1/n : n = 1, 2, \dots\}$ as a subset of $X := \mathbb{R}$ with the euclidean metric. This set is not closed. Indeed, note that its complement includes the point 0, and 0 is not an interior point of the complement. Indeed, for every open ball $B(0, r)$, there exists some positive integer n such that $1/n \leq r$ using the archimedean property of the reals, which means that $1/n$ is a point of $B(0, r)$, so $B(0, r)$ is not entirely contained in the complement of E .

Lemma 1.3. *Let X be a metric space.*

- (a) *If a set I indexes a collection $(E_i)_{i \in I}$ of closed subsets of X , then $E := \bigcap_{i \in I} E_i$ is also closed.*
- (b) *If E_1, \dots, E_n are all closed subsets of X , then $E_1 \cup \dots \cup E_n$ is also closed.*

Proof. This follows immediately from the analogous result for open sets and De Morgan's Laws. I leave it as an exercise for you to fill in the details. \square

Closely related to the notion of closedness is the notion of limit points. Let X be a metric space. A point $a \in X$ is a *limit point* of E if, for every open set U containing a , there exists some point $x \in E \cap U$ which is distinct from a . Note that a limit point of E need not be a point of E itself.

Example 1.4. Let $X := \mathbb{R}$ and $E := \{1/n : n = 1, 2, \dots\}$. The only limit point of E is 0. To see that 0 is a limit point, we basically repeat the same argument as above. Given any open set U containing 0, there exists an open ball $B(0, r)$ contained entirely inside U since U is open. Then we can find some positive integer n such that $1/n \leq r$ using the archimedean property, which means that $1/n \in E \cap B(0, r) \subseteq E \cap U$ and $1/n$ is distinct from 0. I will leave it as an exercise for you to convince yourself that 0 is the only limit point.

Example 1.5. Inside $X := \mathbb{R}$ with the euclidean metric, consider the set $E := (-\infty, 0]$. This set is closed, because its complement $(0, \infty)$ is open as we saw last time. Also, the set of limit points of E is precisely equal to E itself. Indeed, suppose first that $a \in E$. Then for any open set U

containing a , there exists an open ball $B(a, r) \subseteq U$. Notice that $a - (r/2) \in E \cap B(a, r) \subseteq E \cap U$ and $a - (r/2)$ is distinct from a itself, so a is a limit point. Conversely, suppose that $a \notin E$. Then a is positive, and the open ball $B(a, a)$ is entirely disjoint from E , so a cannot be a limit point of E .

Lemma 1.6. *Let X be a metric space. Then a subset E of X is closed if and only if every limit point of E is contained in E .*

Proof. Suppose E is closed. Let us show that if $a \notin E$, then a is not a limit point of E . Since $a \notin E$, a is a point in the open set $X \setminus E$. But $X \setminus E$ is disjoint from E , so it contains no points of E , so a is not a limit point of E .

Conversely, suppose every limit point of E is contained in E . We want to show that E is closed, or, in other words, that $X \setminus E$ is open. Suppose $a \in X \setminus E$. Since a is not in E , in particular it is not a limit point of E . That means that there exists an open set U containing a which contains no points of E . In other words, $U \subseteq X \setminus E$. But then since U is open, there exists an open ball $B(a, r) \subseteq U$, so $B(a, r) \subseteq X \setminus E$ also. Thus $X \setminus E$ is open, so E is closed. \square

Example 1.7. It need not be that every point of a closed set is a limit point of that set. For example, consider $E := \{0\}$ inside $X := \mathbb{R}$. Then $X \setminus E = (-\infty, 0) \cup (0, \infty)$ is open, but 0 is not a limit point of E since no open subset of \mathbb{R} containing 0 also contains a point of E which is distinct from E .

2 Closure

If X is a metric space and E is any subset, we define the *closure* of E , denoted \bar{E} , to be the intersection of all closed subsets of X which contain E . Lemma 1.3(a) guarantees immediately that \bar{E} is closed. Thus, \bar{E} is the smallest closed subset of X containing E .

Proposition 2.1. *Let X be a metric space and let E be a subset. If E' is the set of limit points of E , then $\bar{E} = E \cup E'$.*

Proof. Suppose F is any closed subset of X containing E . If $a \in E'$, then a is also a limit point of F clearly, but F is closed and contains all of its limit points by lemma 1.6, so $a \in F$. Thus $E' \subseteq F$, so $E \cup E' \subseteq F$. Thus every closed set containing E contains $E \cup E'$, so $E \cup E' \subseteq \bar{E}$.

To see that $E \cup E' = \bar{E}$, it suffices to show that $E \cup E'$ is itself closed. We do this using lemma 1.6 by showing that $E \cup E'$ contains all of its limit points. Suppose that a is a limit point of $E \cup E'$ and suppose that $a \notin E$. We want to show that then a must be a limit point of E also. Suppose that it is not a limit point of E , and let U be an open set containing a which contains no points of E . Since a is a limit point of $E \cup E'$, we know that U contains some point $b \in E \cup E'$ distinct from a . But U contains no points of E by assumption, so actually we must have that b is a point of E' distinct from a . But then U is an open set containing b , so since b is a limit point of E , U must contain some point of E . This is a contradiction, since U was supposed to contain no points of E . \square

3 Density

Let X be a metric space. Then a subset $E \subseteq X$ is *dense* inside X if $\bar{E} = X$.

Lemma 3.1. *Let X be a metric space and E a subset. Then E is dense inside X if and only if $E \cap U$ is nonempty for every nonempty open subset U of X .*

Proof. Suppose E is dense inside X and suppose for a contradiction that there exists some nonempty open subset U of X such that $E \cap U = \emptyset$. Then $F := X \setminus U$ is a closed set containing E , so $\bar{E} \subseteq F$. In particular, \bar{E} does not contain any points of U , but this is a contradiction since U is nonempty and we were supposed to have $\bar{E} = X$.

Conversely, suppose $E \cap U$ is nonempty whenever U is a nonempty open subset of X . Let F be a closed set containing E . Then $U := X \setminus F$ is an open set and clearly $E \cap U = \emptyset$ since E is contained inside F . This means that U must already be empty, since if it was nonempty it would have nonempty intersection with E by assumption. Now U being empty means that $F = X$. Since any closed set containing E is equal to X , we conclude that $\bar{E} = X$. \square

4 Sample Problems

Problem 1. Let E be a subset of a metric space X and let a be a limit point of E . Show that every open subset of X containing a also contains infinitely many points of E .

Solution. Let U be an open subset of X containing a and suppose for that U contains only finitely many points of E . In other words, we are assuming that $E \cap U$ is finite. Let x_1, \dots, x_n be the elements of $E \cap U$ distinct from a . Since U is open, there exists an open ball $B(a, r) \subseteq U$. Let

$$\varepsilon := \min\{r, d(a, x_1), \dots, d(a, x_n)\}.$$

Then $\varepsilon > 0$ since all of the elements of the set are also positive, so $B(a, \varepsilon)$ is an open ball. It is contained inside U since $\varepsilon \leq r$, so $B(a, \varepsilon) \subseteq B(a, r) \subseteq U$, and moreover $x_i \notin B(a, \varepsilon)$ for all i since $d(a, x_i) \geq \varepsilon$ for all i . Thus the open ball $B(a, \varepsilon)$ contains no points of E other than possibly a itself, so a is not a limit point for E . \square

Problem 2. Let $X := \mathbb{R}$ and $E := (0, \infty)$. What is the closure of E when \mathbb{R} is given the euclidean metric? What is the closure of E when \mathbb{R} is given the discrete metric?

Solution. The closure of E when \mathbb{R} has the euclidean metric is $[0, \infty)$. Indeed, 0 is the only limit point of E which is not already contained in E . To see this, notice that for any $a < 0$, the set $(-\infty, 0)$ is an open subset of \mathbb{R} containing a but containing no points of E , so a cannot be a limit point of E . Next, when \mathbb{R} is given the discrete metric, all subsets are open, so all subsets are also closed. Thus $(0, \infty)$ is already closed, so it is its own closure. \square

Problem 3. Let E be a nonempty subset of \mathbb{R} which is bounded above, and let $\alpha := \sup E$. Show that $\alpha \in \bar{E}$.

Solution. If $\alpha \in E$, then we are done immediately, so suppose $\alpha \notin E$. Since $\bar{E} = E \cup E'$ where E' is the set of limit points of E , in order to have $\alpha \in \bar{E}$ even though $\alpha \notin E$, we must have $\alpha \in E'$, so let us show this. Let U be any open subset containing α . Then there exists some open ball $B(\alpha, r) \subseteq U$. Since $\alpha - r \not\leq \alpha$, we know that $\alpha - r$ is not an upper bound for α , so there exists some $x \in E$ such that $\alpha - r \leq x \leq \alpha$. In other words, we have $|\alpha - x| \leq r$, so $x \in B(\alpha, r) \cap E \subseteq U \cap E$. Thus α is a limit point of E . \square

Problem 4. Show that \mathbb{Q} is dense inside \mathbb{R} with the euclidean metric.

Proof. Let U be a nonempty open subset of \mathbb{R} . Then there exists some $a \in U$ and since U is open there exists some open ball $B(a, r)$ inside U . But by the result we've been calling "density of the rationals," we know that there exists a rational number x such that $a \leq x \leq a + r$, which means that $x \in B(a, r) \subseteq U$, so $\mathbb{Q} \cap U$ is nonempty. Thus \mathbb{Q} is dense using lemma 3.1. \square

Problem 5. Consider \mathbb{Z} with the discrete metric. Describe all dense subsets of \mathbb{Z} , and prove that you have described all of them.

Solution. Suppose E is a dense subset of \mathbb{Z} . Pick any $a \in \mathbb{Z}$. Then $B(a, 1/2) = \{a\}$ is an open ball, so $E \cap B(a, 1/2)$ must be nonempty by lemma 3.1. But the only element of $B(a, 1/2)$ is a itself, so $a \in E$. Thus $E = \mathbb{Z}$. Thus the only dense subset of \mathbb{Z} is \mathbb{Z} itself. \square