

Skills

Blood Pressure: Orthostatic - CE/NCPD

Checklist

Checklist

S = Satisfactory U = Unsatisfactory NP = Not Performed

Step	S	U	NP	Comments
Complete all pre-Skill patient education, assessment, and Skill preparation per individual scope of practice and role.	☐	☐	☐	
Be sure that the patient is lying in a flat, supine position for a minimum of 2 minutes before taking the initial blood pressure and heart rate measurements.	☐	☐	☐	
Measure the patient's blood pressure and heart rate after the patient has been in the supine position for the necessary amount of time. Use the appropriate-size blood pressure cuff.	☐	☐	☐	
Assist the patient to a standing position. 1. Assist the patient, as needed, with moving from a supine to a standing position. When moving the patient from supine to standing, enlist assistance from another health care team member to ensure patient safety. 2. If the patient is unable to stand for a blood pressure measurement, help the patient to a sitting position. Do not take measurements if the patient becomes extremely dizzy and needs to lie down or experiences syncope.	☐	☐	☐	
Measure the patient's blood pressure and heart rate and assess for symptoms upon standing or sitting. 1. Take the measurements in the same arm as the initial reading. 2. Be sure that the patient's arm, relative to the patient's trunk, is consistent with the arm's position when the patient was supine. 3. Take the measurements again after 1 to 3 minutes of standing or sitting. 4. Ask the patient about symptoms of weakness, dizziness, or visual dimming associated with a change of position. Observe for pallor or diaphoresis.	☐	☐	☐	
Compare the blood pressure and heart rate results.	☐	☐	☐	
Assist the patient as needed with returning to a supine or sitting position.	☐	☐	☐	
Complete all post-Skill patient education, documentation, and reassessment per individual scope of practice and role.	☐	☐	☐	

Learner: _____
Evaluator: _____
Date: _____

Signature: _____
Signature: _____