

Skills

Inserting an Indwelling Urinary Catheter in a Male Patient

Checklist

Checklist

S = Satisfactory U = Unsatisfactory NP = Not Performed

Step	S	U	NP	Comments
Verified the health care provider's orders.	☐	☐	☐	
Reviewed the medical record for any issues that may hinder the procedure.	☐	☐	☐	
Provided for the patient's privacy.	☐	☐	☐	
Performed hand hygiene.	☐	☐	☐	
Introduced self to the patient and family.	☐	☐	☐	
Identified the patient using two identifiers.	☐	☐	☐	
Explained the provider's order and the procedure to the patient and ensured that he agreed to treatment.	☐	☐	☐	
Examined the patient by using a bladder scanner or by palpating the patient's bladder for distention.	☐	☐	☐	
Gathered the necessary equipment and supplies. Checked the patient's plan of care for the size and type of catheter to use if for a reinsertion. Used the smallest-size catheter possible.	☐	☐	☐	
Performed hand hygiene and applied clean gloves. Donned appropriate personal protective equipment (PPE) based on the patient's need for isolation precautions or risk of exposure to bodily fluids.	☐	☐	☐	
Raised the bed to a comfortable working height. Adjusted side rails accordingly.	☐	☐	☐	
Positioned and draped the patient: 1. Positioned the patient supine. 2. Draped the patient's upper and lower body so that only the perineum was exposed.	☐	☐	☐	

3. Made sure there was a waterproof absorbent pad under the patient.

Washed the perineal area with soap and water, rinsed, and dried.
Examined the patient and identified the urinary meatus. Retracted the foreskin if needed. Removed and discarded used gloves.

☐☐☐

Performed hand hygiene.

☐☐☐

Positioned a trash can on the floor nearby or placed a bag between the nurse and the patient.

☐☐☐

Obtained an indwelling catheterization kit. Opened the outer wrapping of the kit by tearing the designated edge of the plastic wrap. Placed the empty outer plastic wrap near the end of the bed and used it for waste disposal. Placed the inner catheter kit tray on a clean surface.

☐☐☐

Opened the inner sterile wrap using sterile technique:

☐☐☐

1. For an open system, began with the separate package containing the drainage bag. Made sure the clamp on the drainage port was closed. Placed the drainage bag and tubing in an easily accessible sterile location.

Applied sterile gloves.

☐☐☐

Draped the patient's perineum, keeping gloves sterile:

☐☐☐

1. Picked up the square drape, and allowed it to unfold without touching unsterile surfaces. Formed a cuff by letting the top edge of the drape fall over both hands.
2. Placed the drape over the patient's thighs. Centered the opening of the fenestrated drape over the penis.

Placed the sterile tray on the sterile drape:

☐☐☐

1. Opened the package of sterile antiseptic solution. Poured the solution over the sterile cotton balls. Opened the end of the package for easy access.
2. Opened the packet of lubricant, and squeezed it out onto the sterile field.
3. Opened the inner sterile wrapper of the catheter.
4. Lubricated the catheter.

Cleansed the patient's urethral meatus:

☐☐☐

1. With nondominant hand, gently grasped the shaft of the penis below the glans; if the patient was uncircumcised, retracted the foreskin. Held the shaft of the penis at a right angle to the body.
2. Using uncontaminated, dominant hand, cleansed the meatus with cotton balls or swab sticks, using circular strokes, beginning at the meatus and working outward in a spiral motion.
3. Repeated process three times, using a clean cotton ball or swab stick each time. Appropriately discarded the cotton balls without contaminating sterile field.

Picked up catheter 7.5 cm to 10 cm from the catheter tip, with the catheter loosely coiled in palm.

☐☐☐

Inserted the catheter:

☐☐☐

1. Gently applied upward traction to the penis while holding it at a 90-degree angle to the patient's body.
2. Asked the patient to push down as if to void, and slowly inserted the catheter through the urethral meatus. If the catheter was not connected to a collection device, made sure the urine would drain into a sterile container.
3. Advanced the catheter 17 cm to 22.5 cm or until urine flowed out of the end of the catheter. Continued to advance the catheter to the bifurcation of the tubing. Stopped advancing catheter if there was resistance or patient reported pain.

Lowered the penis and held the catheter securely with nondominant hand.

☐☐☐

Inflated the catheter balloon with the amount of fluid designated by the manufacturer. Did not inflate the balloon with a 0.9% sodium chloride solution as the fluid in the balloon could crystalize making it difficult to remove.

☐☐☐

1. With dominant hand, connected the prefilled syringe to the injection port at the end of the catheter.
2. Slowly injected the total amount of solution.
3. After inflating the catheter balloon, released the catheter. Gently pulled the catheter until resistance was felt.

Removed sterile gloves, performed hand hygiene, and applied clean gloves.

☐☐☐

Secured the indwelling catheter with a catheter strap or other securement device just at or above the catheter bifurcation. Left enough slack to allow leg movement. Attached the securement device. Secured the catheter tubing to the patient's upper medial aspect of the inner thigh or lower abdomen, with the penis directed toward the patient's chest. Allowed sufficient slack in the catheter tubing to prevent tension.

☐☐☐

Positioned the drainage bag lower than the bladder by attaching it to the bed frame.

☐☐☐

Coiled any excess tubing on the bed, and fastened it to the bottom sheet with a clip or other securement device. Ensured that the tubing had no obstructions or kinks.

☐☐☐

When the patient's foreskin was retracted, cleaned off the iodine preparation and replaced the foreskin over the glans penis.

☐☐☐

Provided hygiene as needed. Helped the patient into a comfortable position, and placed toiletries and personal items within reach.

☐☐☐

Provided patient teaching with Teach Back to promote patient understanding.

☐☐☐

Raised the appropriate number of side rails and lowered the bed to the lowest position.

☐☐☐

Disposed of supplies in the appropriate receptacles.

☐☐☐

Removed and disposed of gloves. Performed hand hygiene.

☐☐☐

Measured and recorded urine output and characteristics of the urine.

☐☐☐

Documented the amount of urine and its characteristics.
Documented and reported the patient's response and expected or unexpected outcomes.

☐☐☐

Learner: _____
Evaluator: _____
Date: _____

Signature: _____
Signature: _____