Evaluation of the unbinding kinetics of Mineralocorticoid receptor (MR) steroid agonist aldosterone, cortisol, and progesteron ligands using molecular simulations: MD and MC

Sebastian Aguilera Novoa

April 14, 2023

Abstract

asdasdasd

I Introduction

Is wildly known the important role of the mineralocorticoid hormone aldosterone (MR/Aldo) in the cardiovascular system, in particular the effects that have in the kidney which have been intensively studied. Nevertheless, several researches have demonstrated the importance of this hormone in the circulatory system. In fact, this hormone is related to several number of common diseases of high mortality and health cost such as hypertension, stroke, heart failure, chronic kidney diseases, obesity, and diabetes disorders which appears when aldosterone is strongly present. Because the main function of MR/Aldo is the regulation of renal sodium balance as well as biological effects several researches have reported its presence in organs like kidney, heart, blood vessels, adipose tissue, brain, eyes, and skin, Figure 1. [1, 2]

The MR/aldo affinity have been already established, but a discovery at 1987 [3], first studies with recombinant human MR (hMR), showed that hMR has strong binding affinities for several corticosteroids: aldosterone (aldo), cortisol (col), corticosterone, and 11-deoxycorticosterone; and for progesterone (str). [1] Nowadays is not well known why MR is more sensitive to aldolsterone than to cortisol, this have been showed by ex vivo experiments which also demonstrated that the glucocorticoid cortisol hormone binds to the hMR with the same affinity as aldo, but is less efficient than aldo in stimulating the hMR transactivation. [4]

Several studies of MR and its mutations have revealed that the substitution of leucine for serine at codon 810 (S810L) in hMR (MR_{S810L}) is responsible of early-onset hypertension and pregnancy-related hypertension raising the risk of pre-eclampsia and early hypertension. Clinical studies on hypertensive patients and their families showed that all MR_{S810L} carriers had developed hypertension before they were 20 yr old, whereas who do not present this mutant had normal blood pressure. [5, 6]




Figure 1: Mineralocorticoid/aldosterone receptor impact on health. *Image taken and adapted from* [1].

Currently, the crystal structure of MR is known and have been obtained using different methods such as X-Ray Diffraction or Cryogenic Electron Microscopy (CryoEM); even, resent articles have showed the asymmetric dimer form of MR [7]. Nevertheless, these methods give just a static view of the protein behavior and do not allow to us to properly describe its dynamical: how MR interacts with a ligand in an un/binding process?. A confident and explored approach to answer this question is the molecular modeling. The huge progress in computational resources has led us to created robust and efficient methods to simulate proteins dynamics, such as Molecular Dynamics (MD) and Monte Carlo (MC) simulations, permitting to the researches explore the dynamical contents.

ics of protein-ligand interactions and given strong explanations to their behavior and are they interacting. [8] orphaz'n question

II State of Art

Selection of the best method to simulate the system, description of the simulation methods and how are we analyzing the date, since the MD simulations have a sense of time while the MC simulations does not, pyEMMA.

Describe how LiGaMD and LiBELa works and the difference between them.

II.I Molecular Dynamics (MD)

Assisted Model Building with Energy Refinement (Amber)

Ligand Gaussian Accelerated Molecular Dynamics 2 (LiGaMD)

[10, 11, 12]

Other option is the making it rain project based on LiG-aMD and google colab. [14]

II.II Monte Carlo (MC)

Ligand Binding Energy Landscape (LiBELa)

[9]

Python API for Emma's Markov Model Algorithms (PyEMMA)

[13]

III Methods

IV Results and Discussion

Select images of interest and explain them

V Conclusion

• LiBELa is able to reproduce unbinding events in the MR/Aldo-Col-Str systems, even when it simulates

just the dynamics of the ligand and consider the protein static.

- Using LiGaMD is not enough to reproduce un/binding process, no matter how long is the simulation since the electrical potential is strong.
- LiGaMD2 is good enough to reproduce un/binding process?

References

- [1] Brian Harvey and Frederic Jaisser. Aldosterone-Mineralocorticoid Receptor. IntechOpen, Rijeka, Sep 2019.
- [2] Leonardo Liccardo Daniela Elia Andrea Marzano Federica Gambino Giuseppina D'Amico Maria Loreta Perna Claudia Ferrara Nicola Rengo Giuseppe Paolocci Nazareno Cannavo, Alessandro. Bencivenga. Aldosterone and mineralocorticoid receptor system in cardiovascular physiology and pathophysiology. Oxidative Medicine and Cellular Longevity, 2018:1204598, Sep 2018.
- [3] Jeffrey L. Arriza, Cary Weinberger, Gail Cerelli, Tom M. Glaser, Barbara L. Handelin, David E. Housman, and Ronald M. Evans. Cloning of human mineralocorticoid receptor complementary dna: Structural and functional kinship with the glucocorticoid receptor. *Science*, 237(4812):268–275, 1987.
- [4] Chantal Hellal-Levy, Jérôme Fagart, Anny Souque, and Marie-Edith Rafestin-Oblin. Mechanistic aspects of mineralocorticoid receptor activation. *Kidney International*, 57(4):1250–1255, 2000.
- [5] Marie-Edith Rafestin-Oblin, Anny Souque, Brigitte Bocchi, Gregory Pinon, Jerome Fagart, and Alain Vandewalle. The Severe Form of Hypertension Caused by the Activating S810L Mutation in the Mineralocorticoid Receptor Is Cortisone Related. *Endocrinology*, 144(2):528–533, 02 2003.
- [6] David S. Geller, Anita Farhi, Nikki Pinkerton, Michael Fradley, Michael Moritz, Adrian Spitzer, Gretchen Meinke, Francis T. F. Tsai, Paul B. Sigler, and Richard P. Lifton. Activating mineralocorticoid receptor mutation in hypertension exacerbated by pregnancy. *Science*, 289(5476):119–123, 2000.
- [7] Carina A. Johansson Anders Gunnarsson Euan Gordon Barry Collins Marie Castaldo Christian Köhler David Öling Patrik Johansson Linda Fröderberg Roth Brice Beinsteiner Ian Dainty Stephen Delaney Bruno P. Klaholz Isabelle M. L. Billas Sandra Postel, Lisa Wissler and Karl Edman. Quaternary glucocorticoid receptor structure highlights allosteric interdomain communication. *Nature Structural & Molecular Biology*, 30(3):286–295, Mar 2023.
- [8] Milton T. Sonoda, Leandro Martínez, Paul Webb, Munir S. Skaf, and Igor Polikarpov. Ligand dissociation from estrogen receptor is mediated by receptor dimerization: Evidence from molecular dynamics simulations. *Molecular Endocrinology*, 22(7):1565–1578, 07 2008.
- [9] Heloisa dos Santos Muniz and Alessandro S. Nascimento. Ligand- and receptor-based docking with libela. Journal of Computer-Aided Molecular Design, 29(8):713–723, Aug 2015.
- [10] K. Belfon I.Y. Ben-Shalom J.T. Berryman S.R. Brozell D.S. Cerutti T.E. Cheatham III G.A. Cisneros V.W.D. Cruzeiro T.A. Darden R.E. Duke G. Giambasu M.K. Gilson H. Gohlke A.W. Goetz R. Harris S. Izadi S.A. Izmailov K. Kasavajhala M.C. Kaymak E. King A. Ko-valenko T. Kurtzman T.S. Lee S. LeGrand P. Li C. Lin J. Liu T. Luchko R. Luo M. Machado V. Man M. Manathunga K.M. Merz Y. Miao O. Mikhailovskii G. Monard H. Nguyen K.A. O'Hearn A. Onufriev F. Pan S. Pantano R. Qi A. Rahnamoun D.R. Roe A. Roitberg C. Sagui S. Schott-Verdugo A. Shajan J. Shen C.L. Simmerling N.R. Skrynnikov J. Smith J. Swails R.C. Walker J. Wang J. Wang H. Wei R.M. Wolf X. Wu Y. Xiong Y. Xue D.M. York S. Zhao D.A. Case, H.M. Aktulga and P.A. Kollman (2022). Amber 2022. University of California, San Francisco.
- [11] Yinglong Miao, Apurba Bhattarai, and Jinan Wang. Ligand gaussian accelerated molecular dynamics (ligand): Characterization of ligand binding thermodynamics and kinetics. bioRxiv, 2020.
- [12] Yinglong Miao, Apurba Bhattarai, and Jinan Wang. Ligand gaussian accelerated molecular dynamics (ligand): Characterization of ligand binding thermodynamics and kinetics. *Journal of Chemical Theory and Computation*, 16(9):5526–5547, 2020. PMID: 32692556.
- [13] Martin K. Scherer, Benjamin Trendelkamp-Schroer, Fabian Paul, Guillermo Pérez-Hernández, Moritz Hoffmann, Nuria Plattner, Christoph Wehmeyer, Jan-Hendrik Prinz, and Frank Noé. PyEMMA 2: A Software Package for Estimation, Validation, and Analysis of Markov Models. *Journal of Chemical Theory and Computation*, 11:5525–5542, October 2015.
- [14] Pablo R. Arantes, Marcelo D. PolÃato, Conrado Pedebos, and Rodrigo Ligabue-Braun. Making-it-rain, August 2021.