


Lecture Series 5B: Diagnostic tests CM2018


Adam S. Darwich

darwich@kth.se


Diagnostic tests



Roberts *et al*. 2021



Epner *et al*. 2013



WHO 2016


Epidemiology

- **Prevalence** = how big proportion of the population has the disease at a given point in time?


4/10 = 40%

- **Incidence** = how big proportion of the population acquires the disease during a given period?

2/10 = 20%



Confusion matrix



Karl Pearson

	Pos. test	Neg. test	Sum
Diseased	TP	FN	$N_{1,1:2}$
Healthy	FP	TN	$N_{2,1:2}$
Sum	$N_{1:2,1}$	$N_{1:2,2}$	$N_{1:2,1:2}$

Real condition

Predicted condition

```
library(caret)

confusionMatrix(data, reference, positive = "1"…)
```

**

TP: true positive
TN: true negative
FP: false positive
FN: false negative

Confusion matrix

		Predicted condition			
	Total population = P + N	**Predicted positive**	**Predicted negative**	Informedness, bookmaker informedness (BM) = TPR + TNR − 1	Prevalence threshold (PT) = $\frac{\sqrt{TPR \times FPR} - FPR}{TPR - FPR}$
Actual condition	**Real Positive (P)** [a]	**True positive** (TP), hit[b]	**False negative** (FN), miss, underestimation	True positive rate (TPR), recall, sensitivity (SEN), probability of detection, hit rate, power = $\frac{TP}{P} = 1 - FNR$	False negative rate (FNR), miss rate type II error [c] = $\frac{FN}{P} = 1 - TPR$
	Real Negative (N) [d]	**False positive** (FP), false alarm, overestimation	**True negative** (TN), correct rejection[e]	False positive rate (FPR), probability of false alarm, fall-out type I error [f] = $\frac{FP}{N} = 1 - TNR$	True negative rate (TNR), specificity (SPC), selectivity = $\frac{TN}{N} = 1 - FPR$
	Prevalence = $\frac{P}{P+N}$	Positive predictive value (PPV), precision = $\frac{TP}{TP+FP} = 1 - FDR$	False omission rate (FOR) = $\frac{FN}{TN+FN} = 1 - NPV$	Positive likelihood ratio (LR+) = $\frac{TPR}{FPR}$	Negative likelihood ratio (LR−) = $\frac{FNR}{TNR}$
	Accuracy (ACC) = $\frac{TP+TN}{P+N}$	False discovery rate (FDR) = $\frac{FP}{TP+FP} = 1 - PPV$	Negative predictive value (NPV) = $\frac{TN}{TN+FN} = 1 - FOR$	Markedness (MK), deltaP (Δp) = PPV + NPV − 1	Diagnostic odds ratio (DOR) = $\frac{LR+}{LR-}$
	Balanced accuracy (BA) = $\frac{TPR+TNR}{2}$	F_1 score = $\frac{2\,PPV \times TPR}{PPV+TPR} = \frac{2\,TP}{2\,TP+FP+FN}$	Fowlkes–Mallows index (FM) = $\sqrt{PPV \times TPR}$	*phi* or Matthews correlation coefficient (MCC) = $\sqrt{TPR \times TNR \times PPV \times NPV} - \sqrt{FNR \times FPR \times FOR \times FDR}$	Threat score (TS), critical success index (CSI), Jaccard index = $\frac{TP}{TP+FN+FP}$

Sources: [12][13][14][15][16][17][18][19] view · talk · edit

a. ^ the number of real positive cases in the data
b. ^ A test result that correctly indicates the presence of a condition or characteristic
c. ^ Type II error: A test result which wrongly indicates that a particular condition or attribute is absent
d. ^ the number of real negative cases in the data
e. ^ A test result that correctly indicates the absence of a condition or characteristic
f. ^ Type I error: A test result which wrongly indicates that a particular condition or attribute is present

Source: https://en.wikipedia.org/wiki/Confusion_matrix

```
library(caret)
library(ROCit)
library(pROC)
library(SIMMS)
...
```

A diagnostic test

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

- Naïve question: How often is the test correct?
- Accuracy = Pr(Test=Truth)
- (25+105)/150 = 87%

A diagnostic test

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

- How big is the chance for a diseased person to be correctly classified?
- Sensitivity = P(Test=pos | Truth=pos)
- 25/30 = 83%

A diagnostic test


	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

- How big is the chance for a healthy person to be correctly classified?
- Specificity = P(Test=neg | Truth=neg)
- 105/120 = 88%

Bayes' theorem

- Thomas Bayes 1763. An Essay Towards Solving a Problem in the Doctrine of Chances.
- A method for calculating the probability of a cause given an effect.

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$



Positive Predictive Value

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

- How big is the probability that a person with a positive test is actually diseased?
- Positive predictive value:
  PPV = P(Truth=pos | Test=pos)

Using Bayes' theorem

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

$$\text{PPV} = P(\text{Truth=pos} \mid \text{Test=pos}) =$$

$$= \frac{P((\text{Truth=pos}) \cap (\text{Test=pos}))}{P(\text{Test=pos})} =$$

$$= \frac{P(\text{Test=pos} \mid \text{Truth=pos}) \times P(\text{Truth=pos})}{P(\text{Test=pos})} =$$

$$= \frac{\text{Sensitivity} \times \text{Prevalence}}{\text{Sensitivity} \times \text{Prevalence} + (1 - \text{Specificity}) \times (1 - \text{Prevalence})} =$$

$$= \frac{(25/30) \times (30/150)}{(40/150)} = \frac{0.167}{0.267} = 63\%$$

25/40 = 63%

Negative Predictive Value

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

- How big is the probability that a person with a negative test is actually healthy?
- Negative predictive value:
  NPV = Pr(Truth=neg | Test=neg)

Using Bayes' theorem

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

$$\text{NPV} = P(\text{Truth=neg} \mid \text{Test=neg}) =$$

$$= \frac{P((\text{Truth=neg}) \cap (\text{Test=neg}))}{P(\text{Test=neg})} =$$

$$= \frac{P(\text{Test=neg} \mid \text{Truth=neg}) \times P(\text{Truth=neg})}{P(\text{Test=neg})} =$$

$$= \frac{\text{Specificity} \times (1 - \text{Prevalence})}{\text{Specificity} \times (1 - \text{Prevalence}) + (1 - \text{Sensitivity}) \times \text{Prevalence}} =$$

$$= \frac{(105/120) \times (120/150)}{(110/150)} = \frac{0.7}{0.733} = 95\%$$

105/110 = 95%

A diagnostic test

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	15	105	**120**
Sum	**40**	**110**	**150**

- How big is the probability that a person with a positive test is actually diseased?
- Positive predictive value:
- 25/40 = 63%

Generate data and a confusion matrix in R

Required Information

- Sensitivity: TP / (TP + FN)
- Specificity: TN / (TN + FP)
- N healthy & N diseased, or N total and prevalence of disease.

```
sens_1 <- 0.98
spec_1 <- 0.95

n_diseased <- 1000
n_healthy <- 1000
n_tot <- n_diseased + n_healthy

Prevalence <- n_diseased / n_tot
TP <- sens_1 * Prevalence*n_tot
TN <- spec_1 * (1 - Prevalence)*n_tot
FP <- (1 - spec_1)*(1 - Prevalence)*n_tot
FN <- (1 - sens_1)*Prevalence*n_tot

y1_obs <- as.factor(c(rep(1, TP), rep(1, FN), rep(0, TN), rep(0, FP)))
y2_pred <- as.factor(c(rep(1, TP), rep(0, FN), rep(0, TN), rep(1, FP)))

library(caret)

confusionMatrix(data = y2_pred, reference = y1_obs, positive = "1")
```

Generate data and a confusion matrix in R

```
Confusion Matrix and Statistics

          Reference
Prediction   0   1
         0 950  20
         1  50 980

               Accuracy : 0.965
                 95% CI : (0.956, 0.9726)
    No Information Rate : 0.5
    P-Value [Acc > NIR] : < 2.2e-16

                  Kappa : 0.93

 Mcnemar's Test P-Value : 0.0005279

            Sensitivity : 0.9800
            Specificity : 0.9500
         Pos Pred Value : 0.9515
         Neg Pred Value : 0.9794
             Prevalence : 0.5000
         Detection Rate : 0.4900
   Detection Prevalence : 0.5150
      Balanced Accuracy : 0.9650

       'Positive' Class : 1
```

A diagnostic test

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	150	1050	**1200**
Sum	**175**	**1055**	**1230**

- How big is the probability that a person with a positive test is actually diseased?
- Positive predictive value:
- 25/175 = 14%

A diagnostic test

	Pos test	Neg test	Sum
Diseased	25	5	**30**
Healthy	150	1050	**1200**
Sum	**175**	**1055**	**1230**

- How big is the probability that a person with a negative test is actually healthy?
- Negative predictive value:
- 1050/1055 = 99.5%

Predictive values depend on the prevalence of the disease. Sensitivity and specificity do not.

Other measures

	Pos test	Neg test	Sum
Diseased	TP	FN	**TP+FN**
Healthy	FP	TN	**FP+TN**
Sum	**TP+FP**	**FN+TN**	**Total**


- True positive rate: $TPR = sensitivity = \Pr(Test = pos|Truth = pos)$
- False positive rate: $FPR = 1 - spec. = \Pr(Test = pos|Truth = neg)$
- True negative rate: $TNR = specificity = \Pr(Test = neg|Truth = neg)$
- False negative rate: $FNR = 1 - sens. = \Pr(Test = neg|Truth = pos)$

Likelihood ratio

	Pos test	Neg test	Sum
Diseased	TP	FN	**TP+FN**
Healthy	FP	TN	**FP+TN**
Sum	**TP+FP**	**FN+TN**	**Total**

- Positive likelihood ratio:
- $LR+ = \frac{TPR}{FPR} = \frac{TP}{TP+FN} / \frac{FP}{FP+TN} = \frac{\Pr(Test = pos|Truth = pos)}{\Pr(Test=pos|Truth=neg)}$
- $= \frac{sensitivity}{1-specificity}$
- Negative likelihood ratio:
- $LR- = \frac{TNR}{FNR} = \frac{TN}{FP+TN} / \frac{FN}{TP+FN} = \frac{\Pr(Test=neg|Truth=neg)}{\Pr(Test=neg|Truth=pos)}$
- $= \frac{specificity}{1-sensitivity}$


Probabilities



How the probabilities are affected by a positive or a negative test can be computed with ”likelihood ratios” (LR+ och LR–)
(See Wikipedia article on ”Likelihood ratios in diagnostic testing”)


Studying odds instead of probability

Regardless of the pre-test value, the odds (=$p/(1–p)$) is multiplied by one constant (>1) for a positive test and by a different constant (<1) for a negative test.



Working with log odds


Regardless of the pre-test value, the log odds is increased by one constant for a positive test and decreased by a different constant for a negative test.



This is similar to how logistic regression works. Every term in the regression equation moves a certain distance on the log odds axis.

Clinical thresholds

Thresholds for treatment vs. expectancy depend on the consequences of each decision




Source: McPhee SJ, Papadakis MA: *Current Medical Diagnosis and Treatment 2011*, 50th Edition: http://www.accessmedicine.com

Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

ROC curve

Sensitivity



1 - Specificity

Source: https://en.wikipedia.org/wiki/Receiver_operating_characteristic

Summary measures and alternative terms

- Accuracy $Acc = \frac{TP+TN}{TP+FP+FN+TN}$
- Youden's index $J = sensitivity + specificity - 1$
- Recall = sensitivity (*in machine learning*)
- Precision = PPV (*in machine learning*)
- F_1 statistic = Dice similarity coefficient =

$$DSC = \frac{|X \cap Y|}{(|X| + |Y|)/2} = \frac{2\,TP}{2\,TP + FP + FN}$$

More terms at: https://en.wikipedia.org/wiki/Sensitivity_and_specificity