

Term presentation

Problem 6

Satvik Saha, 19MS154

November 19, 2020

MA2102: Linear Algebra I

Indian Institute of Science Education and Research, Kolkata

Problem statement

Find a basis of the quotient space $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$, where $\text{Sym}_n(\mathbb{R})$ is the subspace of symmetric matrices.

The subspace of symmetric matrices is such that for any $A \in \text{Sym}_n(\mathbb{R})$,

$$A = A^T \quad \Leftrightarrow \quad a_{ij} = a_{ji}.$$

It can be shown that the skew-symmetric matrices form a subspace such that for any $B \in \text{Skew}_n(\mathbb{R})$,

$$B = -B^T \quad \Leftrightarrow \quad b_{ij} = -b_{ji}.$$

The subspace of symmetric matrices is such that for any $A \in \text{Sym}_n(\mathbb{R})$,

$$A = A^{\top} \quad \Leftrightarrow \quad a_{ij} = a_{ji}.$$

It can be shown that the skew-symmetric matrices form a subspace such that for any $B \in \text{Skew}_n(\mathbb{R})$,

$$B = -B^{\top} \quad \Leftrightarrow \quad b_{ij} = -b_{ji}.$$

Preliminaries

Any matrix $X \in M_n(\mathbb{R})$ can be written uniquely as the sum of a symmetric and a skew-symmetric matrix.

$$X = \frac{1}{2}(X + X^T) + \frac{1}{2}(X - X^T).$$

Furthermore, $\text{Sym}_n(\mathbb{R}) \cap \text{Skew}_n(\mathbb{R}) = \{\mathbf{0}\}$, because if X is both symmetric and skew-symmetric,

$$X^T = X = -X^T \quad \Rightarrow \quad X = \mathbf{0}.$$

Thus, we can write

$$M_n(\mathbb{R}) = \text{Sym}_n(\mathbb{R}) \oplus \text{Skew}_n(\mathbb{R}).$$

Preliminaries

Any matrix $X \in M_n(\mathbb{R})$ can be written uniquely as the sum of a symmetric and a skew-symmetric matrix.

$$X = \frac{1}{2}(X + X^T) + \frac{1}{2}(X - X^T).$$

Furthermore, $\text{Sym}_n(\mathbb{R}) \cap \text{Skew}_n(\mathbb{R}) = \{\mathbf{0}\}$, because if X is both symmetric and skew-symmetric,

$$X^T = X = -X^T \quad \Rightarrow \quad X = \mathbf{0}.$$

Thus, we can write

$$M_n(\mathbb{R}) = \text{Sym}_n(\mathbb{R}) \oplus \text{Skew}_n(\mathbb{R}).$$

Preliminaries

Any matrix $X \in M_n(\mathbb{R})$ can be written uniquely as the sum of a symmetric and a skew-symmetric matrix.

$$X = \frac{1}{2}(X + X^T) + \frac{1}{2}(X - X^T).$$

Furthermore, $\text{Sym}_n(\mathbb{R}) \cap \text{Skew}_n(\mathbb{R}) = \{\mathbf{0}\}$, because if X is both symmetric and skew-symmetric,

$$X^T = X = -X^T \quad \Rightarrow \quad X = \mathbf{0}.$$

Thus, we can write

$$M_n(\mathbb{R}) = \text{Sym}_n(\mathbb{R}) \oplus \text{Skew}_n(\mathbb{R}).$$

Basis of $\text{Skew}_n(\mathbb{R})$

Let $E_{ij} \in M_n(\mathbb{R})$ be the matrix whose i, j^{th} element is 1, and the remaining elements are 0. The set of β of all such E_{ij} comprises the standard basis of $M_n(\mathbb{R})$. For any $X \in M_n(\mathbb{R})$,

$$X = \sum_{i=1}^n \sum_{j=1}^n x_{ij} E_{ij}.$$

Set

$$B_{ij} = E_{ij} - E_{ji}.$$

Thus, B_{ij} is a skew-symmetric matrix with 1 in the i, j^{th} position, and -1 in the j, i^{th} position. Let

$$\gamma = \{B_{ij} : 1 \leq i < j \leq n\}.$$

Basis of $\text{Skew}_n(\mathbb{R})$

Let $E_{ij} \in M_n(\mathbb{R})$ be the matrix whose i, j^{th} element is 1, and the remaining elements are 0. The set of β of all such E_{ij} comprises the standard basis of $M_n(\mathbb{R})$. For any $X \in M_n(\mathbb{R})$,

$$X = \sum_{i=1}^n \sum_{j=1}^n x_{ij} E_{ij}.$$

Set

$$B_{ij} = E_{ij} - E_{ji}.$$

Thus, B_{ij} is a skew-symmetric matrix with 1 in the i, j^{th} position, and -1 in the j, i^{th} position. Let

$$\gamma = \{B_{ij} : 1 \leq i < j \leq n\}.$$

Basis of $\text{Skew}_n(\mathbb{R})$

The linear independence of γ follows from the linear independence of $\beta = \{E_{ij}\}$.

$$\sum_{i < j} c_{ij} B_{ij} = \sum_{i < j} c_{ij} E_{ij} - c_{ij} E_{ji} = \mathbf{0}.$$

Suppose $B \in \text{Skew}_n(\mathbb{R})$. Then, $b_{ij} = -b_{ji}$ and $b_{ii} = 0$.

$$\begin{aligned} B &= \sum_{i=1}^n \sum_{j=1}^n b_{ij} E_{ij} = \sum_{i < j} b_{ij} E_{ij} + \sum_{i=j} b_{ij} E_{ij} + \sum_{i > j} b_{ij} E_{ij} \\ &= \sum_{i < j} b_{ij} E_{ij} + \mathbf{0} + \sum_{i < j} b_{ji} E_{ji} \\ &= \sum_{i < j} b_{ij} E_{ij} - b_{ij} E_{ji} = \sum_{i < j} b_{ij} B_{ij}. \end{aligned}$$

Basis of $\text{Skew}_n(\mathbb{R})$

The linear independence of γ follows from the linear independence of $\beta = \{E_{ij}\}$.

$$\sum_{i < j} c_{ij} B_{ij} = \sum_{i < j} c_{ij} E_{ij} - c_{ij} E_{ji} = \mathbf{0}.$$

Suppose $B \in \text{Skew}_n(\mathbb{R})$. Then, $b_{ij} = -b_{ji}$ and $b_{ii} = 0$.

$$\begin{aligned} B &= \sum_{i=1}^n \sum_{j=1}^n b_{ij} E_{ij} = \sum_{i < j} b_{ij} E_{ij} + \sum_{i=j} b_{ij} E_{ij} + \sum_{i > j} b_{ij} E_{ij} \\ &= \sum_{i < j} b_{ij} E_{ij} + \mathbf{0} + \sum_{i < j} b_{ji} E_{ji} \\ &= \sum_{i < j} b_{ij} E_{ij} - b_{ij} E_{ji} = \sum_{i < j} b_{ij} B_{ij}. \end{aligned}$$

Quotient spaces

Let V be vector space over F and let $W \subseteq V$ be a subspace. The quotient space V/W consists of equivalence classes $[v]$, where $v \in V$ and

$$[v] = v + W = \{v + w : w \in W\}.$$

Equivalently, $u \in [v]$ if and only if $u - v \in W$.

We define

$$[u] + [v] = [u + v], \quad \lambda[v] = [\lambda v].$$

With this, V/W is a vector space over F .

Quotient spaces

Let V be vector space over F and let $W \subseteq V$ be a subspace. The quotient space V/W consists of equivalence classes $[v]$, where $v \in V$ and

$$[v] = v + W = \{v + w : w \in W\}.$$

Equivalently, $u \in [v]$ if and only if $u - v \in W$.

We define

$$[u] + [v] = [u + v], \quad \lambda[v] = [\lambda v].$$

With this, V/W is a vector space over F .

Quotient spaces

Let $v \in V$ and $w_1, w_2 \in W$. Then

$$[v + w_1] = [v + w_2].$$

Pick $u \in [v + w_1]$. Then $u = v + w_1 + w'_1$ for some $w'_1 \in W$. Now, $w_1 + w'_1 \in W$, so $(w_1 + w'_1) - w_2 \in W$. This means that

$$u = v + w_1 + w'_1 = v + w_2 + (w_1 + w'_1 - w_2) \in [v + w_2].$$

The reverse inclusion follows by symmetry.

Alternatively, note that since addition is well-defined,

$$[v + w_1] = [v] + [w_1] = [v] + [0] = [v] + [w_2] = [v + w_2].$$

Quotient spaces

Let $v \in V$ and $w_1, w_2 \in W$. Then

$$[v + w_1] = [v + w_2].$$

Pick $u \in [v + w_1]$. Then $u = v + w_1 + w'_1$ for some $w'_1 \in W$. Now, $w_1 + w'_1 \in W$, so $(w_1 + w'_1) - w_2 \in W$. This means that

$$u = v + w_1 + w'_1 = v + w_2 + (w_1 + w'_1 - w_2) \in [v + w_2].$$

The reverse inclusion follows by symmetry.

Alternatively, note that since addition is well-defined,

$$[v + w_1] = [v] + [w_1] = [v] + [0] = [v] + [w_2] = [v + w_2].$$

Basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$

We claim that the set

$$\gamma' = \{[B_{ij}]: B_{ij} \in \gamma\} = \{[B_{ij}]: 1 \leq i < j \leq n\}$$

is a basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$.

Consider the linear combination

$$\sum_{i < j} c_{ij} [B_{ij}] = [0]$$

$$[B] = [0]$$

This means that the skew-symmetric matrix $B \in [0]$, i.e.

$B = 0 + A = A$ for some $A \in \text{Sym}_n(\mathbb{R})$. This forces $B = 0$, whence $c_{ij} = 0$. Thus, γ' is linearly independent.

Basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$

We claim that the set

$$\gamma' = \{[B_{ij}]: B_{ij} \in \gamma\} = \{[B_{ij}]: 1 \leq i < j \leq n\}$$

is a basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$.

Consider the linear combination

$$\sum_{i < j} c_{ij} [B_{ij}] = [0]$$

$$[B] = [0]$$

This means that the skew-symmetric matrix $B \in [0]$, i.e.

$B = 0 + A = A$ for some $A \in \text{Sym}_n(\mathbb{R})$. This forces $B = 0$, whence $c_{ij} = 0$. Thus, γ' is linearly independent.

Basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$

Pick $[X] \in M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$. Since $X \in M_n(\mathbb{R})$, write $X = A + B$ where $A \in \text{Sym}_n(\mathbb{R})$ and $B \in \text{Skew}_n(\mathbb{R})$. Note that

$$[X] = [A + B] = [B].$$

Now, expand B in the basis γ .

$$B = \sum_{i < j} b_{ij} B_{ij}.$$

Then,

$$[X] = [B] = \sum_{i < j} b_{ij} [B_{ij}].$$

Basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$

Pick $[X] \in M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$. Since $X \in M_n(\mathbb{R})$, write $X = A + B$ where $A \in \text{Sym}_n(\mathbb{R})$ and $B \in \text{Skew}_n(\mathbb{R})$. Note that

$$[X] = [A + B] = [B].$$

Now, expand B in the basis γ .

$$B = \sum_{i < j} b_{ij} B_{ij}.$$

Then,

$$[X] = [B] = \sum_{i < j} b_{ij} [B_{ij}].$$

Thus, γ' is linearly independent and spans $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$. This proves that γ' is a basis of $M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R})$.

Moreover, γ and γ' contain $1 + 2 + \cdots + (n - 1) = n(n - 1)/2$ elements, so

$$\dim \text{Skew}_n(\mathbb{R}) = \dim M_n(\mathbb{R}) / \text{Sym}_n(\mathbb{R}) = \frac{1}{2}n(n - 1).$$

Appendix

For a linear map $T: V \rightarrow W$, the map

$$\mathcal{T}: V / \ker T \rightarrow \operatorname{im} T, \quad [v] \mapsto T(v)$$

is a linear isomorphism. By setting

$$T: M_n(\mathbb{R}) \rightarrow M_n(\mathbb{R}), \quad X \mapsto \frac{1}{2}(X - X^T),$$

note that $\ker T = \operatorname{Sym}_n(\mathbb{R})$ and $\operatorname{im} T = \operatorname{Skew}_n(\mathbb{R})$.

If $T(X) = B \in \operatorname{Skew}_n(\mathbb{R})$, then $X = [B]$. Since a linear isomorphism sends a basis to a basis, and the inverse of a linear isomorphism is a linear isomorphism, the set $\mathcal{T}^{-1}(\gamma) = \gamma'$ is a basis of $M_n(\mathbb{R}) / \operatorname{Sym}_n(\mathbb{R})$.