

MA3101

Analysis III

Autumn 2021

Satvik Saha
19MS154

*Indian Institute of Science Education and Research, Kolkata,
Mohanpur, West Bengal, 741246, India.*

Contents

1 Euclidean spaces

1

1 Euclidean spaces

We are familiar with the vector space \mathbb{R}^n , with the standard inner product

$$\langle \mathbf{x}, \mathbf{y} \rangle = x_1 y_1 + \cdots + x_n y_n.$$

The standard norm is defined as

$$\|\mathbf{x} - \mathbf{y}\|^2 = \langle \mathbf{x} - \mathbf{y}, \mathbf{x} - \mathbf{y} \rangle = \sum_{k=1}^n (x_k - y_k)^2.$$

Exercise 1.1. What are all possible inner products on \mathbb{R}^n ?

Solution. Note that an inner product is a bilinear, symmetric map such that $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$, and $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ if and only if $\mathbf{x} = \mathbf{0}$. Thus, an inner product map on \mathbb{R}^n is completely and uniquely determined by the values $\langle \mathbf{e}_i, \mathbf{e}_j \rangle = a_{ij}$. Let A be the $n \times n$ matrix with entries a_{ij} . Note that A is a real symmetric matrix with positive entries. Now,

$$\langle \mathbf{x}, \mathbf{e}_j \rangle = x_1 a_{1j} + \cdots + x_n a_{nj} = \mathbf{x}^\top \mathbf{a}_j,$$

where \mathbf{a}_j is the j^{th} column of A . Thus,

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{a}_1 y_1 + \cdots + \mathbf{x}^\top \mathbf{a}_n y_n = \mathbf{x}^\top A \mathbf{y}.$$

Furthermore, any choice of real symmetric A with positive entries produces an inner product.

Theorem 1.1 (Cauchy-Schwarz). *Given two vectors $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$, we have*

$$|\langle \mathbf{v}, \mathbf{w} \rangle| \leq \|\mathbf{v}\| \|\mathbf{w}\|.$$

Proof. This is trivial when $\mathbf{w} = \mathbf{0}$. When $\mathbf{w} \neq \mathbf{0}$, set $\lambda = \langle \mathbf{v}, \mathbf{w} \rangle / \|\mathbf{w}\|^2$. Thus,

$$0 \leq \|\mathbf{v} - \lambda \mathbf{w}\|^2 = \|\mathbf{v}\|^2 - 2\lambda \langle \mathbf{v}, \mathbf{w} \rangle + \lambda^2 \|\mathbf{w}\|^2.$$

Simplifying,

$$0 \leq \|\mathbf{v}\|^2 - \frac{|\langle \mathbf{v}, \mathbf{w} \rangle|^2}{\|\mathbf{w}\|^2}.$$

This gives the desired result. Clearly, equality holds if and only if $\mathbf{v} = \lambda \mathbf{w}$. \square

Theorem 1.2 (Triangle inequality). *Given two vectors $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$, we have*

$$\|\mathbf{v} + \mathbf{w}\| \leq \|\mathbf{v}\| + \|\mathbf{w}\|.$$

Proof. Write

$$\|\mathbf{v} + \mathbf{w}\|^2 = \|\mathbf{v}\|^2 + 2\langle \mathbf{v}, \mathbf{w} \rangle + \|\mathbf{w}\|^2 \leq \|\mathbf{v}\|^2 + 2|\langle \mathbf{v}, \mathbf{w} \rangle| + \|\mathbf{w}\|^2.$$

Applying Cauchy-Schwarz gives

$$\|\mathbf{v} + \mathbf{w}\|^2 \leq (\|\mathbf{v}\| + \|\mathbf{w}\|)^2.$$

Equality holds if and only if $\mathbf{v} = \lambda \mathbf{w}$ for $\lambda \geq 0$. \square

This allows us to define the standard metric on \mathbb{R}^n , seen as a point set.

$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|.$$

Definition 1.1. For any $\epsilon > 0$, the set

$$B_\epsilon(\mathbf{x}) = \{\mathbf{y} \in \mathbb{R}^n : d(\mathbf{x}, \mathbf{y}) < \epsilon\}$$

is called the open ball centred at $\mathbf{x} \in \mathbb{R}^n$ with radius ϵ . This is also called the ϵ neighbourhood of \mathbf{x} .

Definition 1.2. A set U is open in \mathbb{R}^n if for every $\mathbf{x} \in U$, there exists an open ball $B_\epsilon(\mathbf{x}) \subset U$.

Remark. Every open ball in \mathbb{R}^n is open.

Remark. Both \emptyset and \mathbb{R}^n are open.

Definition 1.3. A set F is closed in \mathbb{R}^n if its complement $\mathbb{R}^n \setminus F$ is open in \mathbb{R}^n .

Remark. Both \emptyset and \mathbb{R}^n are closed.