IISER Kolkata Problem Sheet IV

MA 1101: Mathematics I

Satvik Saha, 19MS154 September 12, 2019

Solution 1.

(i) Let P(n) be the statement

$$1+2+\cdots+n = \frac{1}{2}n(n+1)$$
 for all $n \in \mathbb{N}$.

Base step We establish P(1). Clearly, $1 = \frac{1}{2}1(1+1)$. Thus, P(1) is true.

Inductive step We assume P(k) is true. We will show that P(k+1) is true.

$$1 + 2 + \dots + k + (k+1) = [1 + 2 + \dots + k] + (k+1)$$

$$= \frac{1}{2}k(k+1) + (k+1) \qquad (From $P(k)$)$$

$$= \frac{1}{2}(k+2)(k+1)$$

$$= \frac{1}{2}(k+1)((k+1)+1)$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

(ii) Let P(n) be the statement

$$1^2 + 2^2 + \dots + n^2 = \frac{1}{6}n(n+1)(2n+1)$$
 for all $n \in \mathbb{N}$.

Base step We establish P(1). Clearly, $1 = \frac{1}{6}1(1+1)(2+1)$. Thus, P(1) is true.

Inductive step We assume P(k) is true. We will show that P(k+1) is true.

$$1^{2} + 2^{2} + \dots + k^{2} + (k+1)^{2} = [1^{2} + 2^{2} + \dots + k^{2}] + (k+1)^{2}$$

$$= \frac{1}{6}k(k+1)(2k+1) + (k+1)^{2} \qquad (From P(k))$$

$$= \frac{1}{6}(k+1)(2k^{2} + k + 6k + 6)$$

$$= \frac{1}{6}(k+1)(2k^{2} + 7k + 6)$$

$$= \frac{1}{6}(k+1)(k+2)(2k+3)$$

$$= \frac{1}{6}(k+1)((k+1) + 1)(2(k+1) + 1)$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

(iii) Let P(n) be the statement

$$1^2 + 3^2 + \dots + (2n-1)^2 = \frac{1}{3}(4n^3 - n)$$
 for all $n \in \mathbb{N}$.

Base step We establish P(1). Clearly, $1 = \frac{1}{3}1(4-3)$. Thus, P(1) is true.

Inductive step We assume P(k) is true. We will show that P(k+1) is true.

$$1^{2} + 3^{2} + \dots + (2k-1)^{2} + (2k+1)^{2} = [1^{2} + 3^{2} + \dots + (2k-1)^{2}] + (2k+1)^{2}$$

$$= \frac{1}{3}(4k^{3} - k) + (2k+1)^{2} \qquad (From P(k))$$

$$= \frac{1}{3}(4k^{3} - k + 12k^{2} + 12k + 3)$$

$$= \frac{1}{3}(4(k^{3} + 3k^{2} + 3k + 1) - k - 1))$$

$$= \frac{1}{3}(4(k+1)^{3} - (k+1))$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

(iv) Let P(n) be the statement

$$1^3 + 2^3 + \dots + n^3 = \frac{1}{4}n^2(n+1)^2$$
 for all $n \in \mathbb{N}$.

Base step We establish P(1). Clearly, $1 = \frac{1}{4}1(1+1)^2$. Thus, P(1) is true.

Inductive step We assume P(k) is true. We will show that P(k+1) is true.

$$1^{3} + 2^{3} + \dots + k^{3} + (k+1)^{3} = [1^{3} + 2^{3} + \dots + k^{3}] + (k+1)^{3}$$

$$= \frac{1}{4}k^{2}(k+1)^{2} + (k+1)^{3} \qquad (From P(k))$$

$$= \frac{1}{4}(k+1)^{2}(k^{2} + 4k + 4)$$

$$= \frac{1}{4}(k+1)^{2}(k+2)^{2}$$

$$= \frac{1}{4}(k+1)^{2}((k+1)+1)^{2}$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

(v) Let P(n) be the statement

$$\sum_{r=1}^{n} r(r+1) \dots (r+9) = \frac{1}{11} n(n+1) \dots (n+10) \quad \text{for all } n \in \mathbb{N}.$$

Base step We establish P(1). Clearly,

$$1(1+1)\dots(1+9) = \frac{1}{11}1(1+1)\dots(1+9)(1+10)$$

Thus, P(1) is true.

Inductive step We assume P(k) is true. We will show that P(k+1) is true.

$$\sum_{r=1}^{k+1} r(r+1)\dots(r+9) = \left[\sum_{r=1}^{k} r(r+1)\dots(r+9)\right] + (k+1)(k+2)\dots(k+1+9)$$

$$= \frac{1}{11}k(k+1)\dots(k+10) + (k+1)(k+2)\dots(k+1+9) \quad (\text{From } P(k))$$

$$= \frac{1}{11}(k+1)\dots(k+10)(k+11)$$

$$= \frac{1}{11}(k+1)\dots((k+1)+9)((k+1)+10)$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

Solution 2.

(i) Let P(n) be the statement that for all $n \in \mathbb{N}$,

$$3^n > n^2$$

Base step We establish P(1) and P(2). Clearly, $3^1 > 1^2$. Thus, P(1) is true. Again, $3^2 = 9 > 8 = 2^2$. Thus, P(2) is true.

Inductive step We assume P(k) is true. We will show that P(k+1) is true.

$$3^{k+1} \ = \ 3 \cdot 3^k \ > \ 3 \cdot k^2$$

We must show $3k^2 > (k+1)^2 \Leftrightarrow 3k^2 - (k+1)^2 > 0$.

$$3k^2 - (k+1)^2 = 2k^2 - 2k - 1 = k^2 + (k-1)^2 - 2$$

Clearly, for $k \ge 2$, $k^2 > 2$, so $k^2 + (k-1)^2 > 2$, and we are done.

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

(ii) Let P(n) be the statement that for all $n \in \mathbb{N}$ and x > -1,

$$(1+x)^n \ge 1 + nx$$
. (Bernoulli's Inequality)

Base Step We establish P(1). Clearly, $(1+x)^1 \ge (1+1\cdot x)$, thus P(1) is true.

Inductive Step We assume P(k) is true. We will show that P(k+1) is true.

$$(1+x)^{k+1} = (1+x)^k \cdot (1+x)$$

$$\geq (1+kx) \cdot (1+x) \qquad (x+1>0)$$

$$= (1+x+kx+kx^2)$$

$$\geq (1+(k+1)x) \qquad (k>0 \text{ and } x^2 \geq 0)$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.

(iii) Let P(n) be the statement that for all $n \geq 5$, $n \in \mathbb{N}$,

$$\binom{2n}{n} < 2^{2n-2}.$$

Base Step We establish P(5). Now, $\binom{2n}{n} = 252$, while $2^{10-2} = 256$. Thus, P(5) is true.

Inductive Step We assume P(k) is true. We will show that P(k+1) is true.

$$\binom{2(k+1)}{k+1} = \frac{(2k+2)!}{(k+1)!^2}$$

$$= \frac{(2k+2)(2k+1)}{(k+1)^2} \binom{2n}{n}$$

$$< 2 \cdot \frac{2k+1}{k+1} \cdot 2^{2k-2}$$

$$< 2 \cdot \frac{2k+2}{k+1} \cdot 2^{2k-2}$$

$$< 2^{2(k+1)-2}$$

Hence, by the principle of mathematical induction, P(n) is true for all $n \in \mathbb{N}$.