

MA2202: PROBABILITY I

Conditional probability

Spring 2021

Satvik Saha
19MS154

*Indian Institute of Science Education and Research, Kolkata,
Mohanpur, West Bengal, 741246, India.*

Suppose a fair die is rolled and the outcome is even. The probability that the outcome was prime is $1/3$. This is because the only even prime is 2, and we know that we have rolled one of 2, 4 or 6, all of which are equally likely.

Definition 2.1 (Conditional probability). Suppose a random experiment is performed, and we know that an event B occurred. Let $A \in \mathcal{E}$. The probability of occurrence of A given B is called the conditional probability $P(A|B)$.

Remark. Note that the conditional probability A given B is equivalent to the occurrence of $A \cap B$ computed relative to the probability of the occurrence of B .

$$P(A|B)P(B) = P(A \cap B).$$

Theorem 2.1 (Bayes' Theorem). If $A, B \in \mathcal{E}$ with $P(B) \neq 0$, we have

$$P(A|B) = \frac{P(A)}{P(B)}P(B|A).$$

Proof. This follows directly from

$$P(A \cap B) = P(A|B)P(B) = P(B|A)P(A).$$

□

Example. Suppose a fair die is rolled and the outcome is even. The probability that the outcome is prime is $1/3$. This is because the only even prime is 2, the available even numbers are 2, 4, 6, and the available primes are 2, 3, 5. Thus,

$$P(\text{prime} | \text{even}) = \frac{P(\text{even})}{P(\text{prime})}P(\text{even} | \text{prime}) = \frac{3/6}{3/6} \cdot \frac{1}{3} = \frac{1}{3}.$$

Definition 2.2 (Independent events). If $A, B \in \mathcal{E}$ such that $P(A), P(B) \neq 0$, and we have

$$P(A|B) = P(A), \quad P(B|A) = P(B),$$

then we say that A and B are independent.

Equivalently, $P(A \cap B) = P(A)P(B)$.