

Assignment 2a

Satvik Saha

2024-09-10

Answer 1

(a)

```
print(seq(0, 20, by = 0.2))
```

```
## [1] 0.0 0.2 0.4 0.6 0.8 1.0 1.2 1.4 1.6 1.8 2.0 2.2 2.4 2.6 2.8
## [16] 3.0 3.2 3.4 3.6 3.8 4.0 4.2 4.4 4.6 4.8 5.0 5.2 5.4 5.6 5.8
## [31] 6.0 6.2 6.4 6.6 6.8 7.0 7.2 7.4 7.6 7.8 8.0 8.2 8.4 8.6 8.8
## [46] 9.0 9.2 9.4 9.6 9.8 10.0 10.2 10.4 10.6 10.8 11.0 11.2 11.4 11.6 11.8
## [61] 12.0 12.2 12.4 12.6 12.8 13.0 13.2 13.4 13.6 13.8 14.0 14.2 14.4 14.6 14.8
## [76] 15.0 15.2 15.4 15.6 15.8 16.0 16.2 16.4 16.6 16.8 17.0 17.2 17.4 17.6 17.8
## [91] 18.0 18.2 18.4 18.6 18.8 19.0 19.2 19.4 19.6 19.8 20.0
```

(b)

```
print(runif(1, min = 0, max = 10))
```

```
## [1] 1.205211
```

(c)

```
print(runif(10, min = 0, max = 10))
```

```
## [1] 0.3874994 5.3239424 2.2474365 1.2084094 0.7240667 1.8308729 0.9932816
## [8] 9.6483512 1.6905259 3.7214616
```

Answer 2

(a)

```
a <- c("Magnus Carlsen", "Hilary Hahn", "Linus Torvalds", "Parsi Diaconis")
```


(b)

```
sample(a, 1)
```

```
## [1] "Magnus Carlsen"
```


(c)

```
x <- seq(0, 10, by = 0.5)
y <- sqrt(x)
plot(x, y)
```



(d)


```
x <- seq(0, 10, length.out = 500)
y <- sqrt(x)
plot(x, y, type = "l")
```




(e)

```
x <- seq(0, 20, length.out = 500)
y <- data.frame(
  y1 = sqrt(x),
  y2 = 2.5 + sqrt(x)
```

```
)
matplot(x, y, type = "l", col = c("blue", "red"), lty = 1)
```



Answer 3



Research homework assignment

Research homework assignment

Suppose that X and Y are independent (real) random variables admitting densities f_X and f_Y , and let $Z = X/Y$. We will say that Z has Cauchy tails when $f_Z(z) \sim C/z^2$ as $|z| \rightarrow \infty$ for some $C > 0$.

Proposition 1: Suppose that $\mathbb{E}[|X|] < \infty$, and that f_Y is bounded. If f_Y is continuous at 0 with $f_Y(0) > 0$, then $Z = X/Y$ has Cauchy tails with

$$f_Z(z) \sim \frac{f_Y(0) \mathbb{E}[|X|]}{z^2} \quad \text{as } |z| \rightarrow \infty.$$

Proof: Observe that the density f_Z of Z is given by

$$f_Z(z) = \int_{\mathbb{R}} |y| f_X(zy) f_Y(y) dy = \frac{1}{z^2} \int_{\mathbb{R}} |x| f_X(x) f_Y\left(\frac{x}{z}\right) dx.$$

This can be seen by applying change of variables on (X, Y) using the map $(x, y) \mapsto (x/y, y)$, whose inverse is $(z, y) \mapsto (zy, y)$, and computing the marginal.

Let $f_Y < M$, so

$$|x| f_X(x) f_Y\left(\frac{x}{z}\right) < |x| f_X(x) M \in L^1(\mathbb{R})$$

via $\mathbb{E}[|X|] < \infty$. By the Dominated Convergence Theorem, we have

$$\lim_{z \rightarrow \infty} z^2 f_Z(z) = \lim_{z \rightarrow \infty} \int_{\mathbb{R}} |x| f_X(x) f_Y\left(\frac{x}{z}\right) dx = \int_{\mathbb{R}} |x| f_X(x) f_Y(0) dx = f_Y(0) \mathbb{E}[|X|]. \quad \square$$

Remark: In the special case that $X, Y \sim N(0, 1)$, we have $X/Y \sim \text{Cauchy}(0, 1)$.

Note that Proposition 1 does not cover situations such as $Y \sim \text{Uniform}(0, 1)$. The continuity assumption on f_Y can be relaxed slightly by dealing with the left and right tails of Z separately.

Proposition 2: Suppose that $\mathbb{E}[|X|] < \infty$, and that f_Y is bounded. If f_Y has both left and right limits at 0, then $Z = X/Y$ has a Cauchy right tail with

$$f_Z(z) \sim \frac{f_Y(0^-) \mathbb{E}[X^-] + f_Y(0^+) \mathbb{E}[X^+]}{z^2} \quad \text{as } z \rightarrow +\infty,$$

and a Cauchy left tail with

$$f_Z(z) \sim \frac{f_Y(0^+) \mathbb{E}[X^-] + f_Y(0^-) \mathbb{E}[X^+]}{z^2} \quad \text{as } z \rightarrow -\infty.$$

Proof: Following the same argument as in Proposition 1, we use the Dominated Convergence Theorem on

$$z^2 f_Z(z) = \int_{-\infty}^0 |x| f_X(x) f_Y\left(\frac{x}{z}\right) dx + \int_0^{+\infty} |x| f_X(x) f_Y\left(\frac{x}{z}\right) dx. \quad \square$$

Remark: Note that $Z = (1/Y)/(1/X)$, so Proposition 2 has an analogue which demands $\mathbb{E}[1/|Y|] < \infty$ and that the density of $1/X$ have both left and right limits at 0.