

MA3104

Linear Algebra II

Autumn 2021

Satvik Saha
19MS154

*Indian Institute of Science Education and Research, Kolkata,
Mohampur, West Bengal, 741246, India.*

Contents

1	Linear operators on a vector space	1
1.1	Preliminaries	1
1.2	Ideals in a ring	1
1.3	Eigenvalues and eigenvectors	2
1.4	Annihilating polynomials	3
1.5	Invariant subspaces	5

1 Linear operators on a vector space

1.1 Preliminaries

We discuss finite dimensional vector spaces V over some field \mathbb{F} , along with linear operators $T: V \rightarrow V$. We also assume that V has the inner product $\langle \cdot, \cdot \rangle$.

Theorem 1.1. *Let $\mathcal{L}(V)$ be the set of all linear operators on the vector space V . Then, $\mathcal{L}(V)$ is a linear algebra over the field \mathbb{F} .*

1.2 Ideals in a ring

Definition 1.1. Let $(R, +, \cdot)$ be a ring, where $(R, +)$ is its additive subgroup. A set $I \subseteq R$ is a left ideal of R if $(I, +)$ is a subgroup of $(R, +)$, and $rx \in I$ for every $r \in R$, $x \in I$.

Example. Let \mathbb{Z} be the ring of integers. For some $n \in \mathbb{N}$, the set $n\mathbb{Z}$ is an ideal. In fact, these are the only ideals (along with $\{0\}$).

Definition 1.2. The principal left ideal generated by $x \in R$ is the set

$$I_x = Rx = \{rx : r \in R\}.$$

Example. In the ring of integers \mathbb{Z} , every ideal is a principal ideal. This follows directly from the fact that $(\mathbb{Z}, +)$ is a cyclic group, thus any subgroup is cyclic and generated by a single element.

Let $I \subseteq \mathbb{Z}$ be an ideal. If $I = \{0\}$, we are done. Otherwise, let n be the smallest positive integer in I (note that if $a \in I$, then $-a \in I$ which means that I must contain positive integers). This immediately gives $I \supseteq n\mathbb{Z}$. Now for any $m \in I$, use Euclid's Division Lemma to write $m = nq + r$, where $q, r \in \mathbb{Z}$, $0 \leq r < n$. Since I is an ideal, $nq \in I$ hence $m - nq = r \in I$. The minimality of n in I forces $r = 0$, hence $m = nq$ and $I \subseteq n\mathbb{Z}$. This proves $I = n\mathbb{Z}$.

Theorem 1.2. Let \mathbb{F} be a field and let $\mathbb{F}[x]$ denote the ring of polynomials with coefficients from \mathbb{F} . Then, every ideal in $\mathbb{F}[x]$ is a principal ideal.

Remark. This is analogous to the theorem which states that every subgroup of a cyclic group is cyclic. Both lead to a precise definition of the greatest common divisor.

Corollary 1.2.1. Let I be a non-trivial ideal in $\mathbb{F}[x]$. Then, there exists a unique monic polynomial $p \in \mathbb{F}[x]$ (leading coefficient 1) such that I is precisely the principal ideal generated by p .

1.3 Eigenvalues and eigenvectors

Definition 1.3. Let $T \in \mathcal{L}(V)$ and $c \in \mathbb{F}$. We say that c is an eigenvalue or characteristic value of T if $T\mathbf{v} = c\mathbf{v}$ for some non-zero $\mathbf{v} \in V$. The vector \mathbf{v} is called an eigenvector of T .

Theorem 1.3. Let $T \in \mathcal{L}(V)$ and $c \in \mathbb{F}$. The following are equivalent.

1. c is an eigenvalue of T .
2. $T - cI$ is singular.
3. $\det(T - cI) = 0$.

Definition 1.4. The polynomial $\det(T - xI)$ is called the characteristic polynomial of T .

Definition 1.5. Two linear operators $S, T \in \mathcal{L}(V)$ are similar if there exists an invertible operator $X \in \mathcal{L}(V)$ such that $S = X^{-1}TX$.

Remark. Similarity is an equivalence relation on $\mathcal{L}(V)$, thus partitioning it into similarity classes.

Lemma 1.4. *Similar linear operators have the same characteristic polynomial.*

Proof. Let S, T be similar with $S = X^{-1}TX$. Then,

$$\begin{aligned}\det(S - xI) &= \det(X^{-1}TX - xX^{-1}X) \\ &= \det(X^{-1}) \det(T - xI) \det(X) \\ &= \det(T - xI).\end{aligned}$$

□

Definition 1.6. A linear operator $T \in \mathcal{L}(V)$ is diagonalizable if there is a basis of V consisting of eigenvectors of T .

Remark. The matrix of T with respect to such a basis is diagonal.

Theorem 1.5. *Let $T \in \mathcal{L}(V)$ where V is finite dimensional, let c_1, \dots, c_k be distinct eigenvalues of T , and let $W_i = \ker(T - c_iI)$ be the corresponding eigenspaces. The following are equivalent.*

1. T is diagonalizable.
2. The characteristic polynomial of T is of the form

$$f(x) = (x - c_1)^{d_1} \dots (x - c_k)^{d_k}$$

where each $d_i = \dim W_i$.

3. $\dim V = \dim W_1 + \dots + \dim W_k$.

1.4 Annihilating polynomials

Definition 1.7. An polynomial p such that $p(T) = 0$ for a given linear operator $T \in \mathcal{L}(V)$ is called an annihilating polynomial of T .

Lemma 1.6. *Every linear operator $T \in \mathcal{L}(V)$, where V is finite dimensional, has a non-trivial annihilating polynomial.*

Proof. Note that the operators $I, T, T^2, \dots, T^{n^2} \in \mathcal{L}(V)$, of which there are $n^2 + 1$, are linearly dependent, since $\dim \mathcal{L}(V) = n^2$. □

Lemma 1.7. *The annihilating polynomials of T form an ideal in $\mathbb{F}[x]$.*

Definition 1.8. The minimal polynomial of T is the unique monic generator of the annihilating polynomials of T .

Remark. The minimal polynomial of T divides all its annihilating polynomials.

Theorem 1.8. *The minimal polynomial and characteristic polynomial of T share the same roots, except for multiplicities.*

Proof. Let p be the minimal polynomial of T and let f be its characteristic polynomial.

First, let $c \in \mathbb{F}$ be a root of the minimal polynomial, i.e. $p(c) = 0$. The Division Algorithm guarantees

$$p(x) = (x - c)q(x)$$

for some monic polynomial q . By the minimality of the degree of p , we have $q(T) \neq 0$, hence there exists non-zero $\mathbf{v} \in V$ such that $\mathbf{w} = q(T)\mathbf{v} \neq \mathbf{0}$. Thus, $p(T)\mathbf{v} = \mathbf{0}$ gives

$$(T - cI)q(T)\mathbf{v} = \mathbf{0}, \quad T\mathbf{w} = c\mathbf{w},$$

which shows that c is an eigenvalue, i.e. a root of the characteristic polynomial f .

Next, suppose that c is a root of the characteristic polynomial, i.e. $f(c) = 0$. Thus, c is an eigenvalue of T , hence there exists non-zero $\mathbf{v} \in V$ such that $T\mathbf{v} = c\mathbf{v}$. This gives $p(T)\mathbf{v} = p(c)\mathbf{v}$, but $p(T) = 0$ identically, forcing $p(c) = 0$. \square

Theorem 1.9 (Cayley-Hamilton). *The characteristic polynomial of T annihilates T .*

Proof. Set $S = \text{adj}(T - xI)$. This is a matrix with polynomial entries, satisfying

$$(T - xI)S = \det(T - xI)I = f(x)I,$$

where f is the characteristic polynomial of T . Now, we can also collect the powers x^n from S and write

$$S = \sum_{k=0}^{n-1} x^k S_k$$

for matrices S_k . Now, calculate

$$\begin{aligned} f(x)I &= (T - xI)S \\ &= (T - xI) \sum_{k=0}^{n-1} x^k S_k \\ &= -x^k S_{k-1} + \sum_{k=1}^{n-1} x^k (TS_k - S_{k-1}) + TS_0. \end{aligned}$$

Compare coefficients with

$$f(x)I = x^n I + a_{n-1}x^{n-1} + \cdots + a_0 I$$

to get

$$S_{n-1} = -I, \quad TS_0 = a_0 I, \quad TS_k - S_{k-1} = a_k I \text{ for } 1 \leq k \leq n-1.$$

Thus,

$$\begin{aligned} f(T) &= \sum_{k=0}^n a_k T^k \\ &= -T^n S_{n-1} + \sum_{k=1}^{n-1} (TS_k - S_{k-1})T^k + TS_0 \\ &= 0. \end{aligned}$$

□

Corollary 1.9.1. *The minimal polynomial of T divides its characteristic polynomial.*

Corollary 1.9.2. *The minimal polynomial of T in a finite-dimensional vector space V is at most $\dim V$.*

Theorem 1.10. *The minimal polynomial for a diagonalizable linear operator T in a finite-dimensional vector space is*

$$p(x) = (x - c_1) \cdots (x - c_k),$$

where c_1, \dots, c_k are distinct eigenvalues of T .

Proof. The diagonalizability of T implies that V admits a basis of eigenvectors of T . Thus, for any such eigenvector \mathbf{v}_i , the operator $T - c_i I$ kills it where c_i is the corresponding eigenvalue. Thus, $p(T)\mathbf{v}_i$ vanishes for every basis vector \mathbf{v}_i □

Remark. The converse is also true, i.e. T is diagonalizable if and only if the minimal polynomial is the product of distinct linear factors.

1.5 Invariant subspaces

Definition 1.9. Let $T \in \mathcal{L}(V)$ where V is finite-dimensional, and let $W \subseteq V$ be a subspace. We say that W is invariant under T if $T(W) \subseteq W$.

If a subspace W is invariant under T , we define the linear map $T_W \in \mathcal{L}(W)$ as the restriction of T to W in the natural way, by setting $T_W(\mathbf{w}) = T(\mathbf{w})$ for all $\mathbf{w} \in W$.

Lemma 1.11. *If W is an invariant subspace under $T \in \mathcal{L}(V)$, then there is a basis of V in which T has the block triangular form*

$$[T]_\beta = \begin{bmatrix} A & B \\ 0 & C \end{bmatrix},$$

where A is an $r \times r$ matrix, $r = \dim W$.

Proof. Let $\beta_W = \{\mathbf{v}_1, \dots, \mathbf{v}_r\}$ be an ordered basis of W , and extend it to an ordered basis $\beta = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ of V . Thus, the matrix $[T]_\beta$ has coefficients a_{ij} such that

$$T\mathbf{v}_j = a_{1j}\mathbf{v}_1 + \dots + a_{rj}\mathbf{v}_r + \dots + a_{nj}\mathbf{v}_n.$$

However for all $j \leq r$, $T\mathbf{v}_j \in W$ by the invariance of W , so the coefficients of $\mathbf{v}_{i>r}$ in the expansion of $T\mathbf{v}_j$ must vanish. Thus, all $a_{ij} = 0$ where $i > r$, $j \leq r$. \square

Lemma 1.12. *If W is an invariant subspace under $T \in \mathcal{L}(V)$, the characteristic polynomial of T_W divides the characteristic polynomial of T , and the minimal polynomial of T_W divides the minimal polynomial of T .*

Proof. Choose an ordered basis β of V such that

$$[T]_\beta = \begin{bmatrix} A & B \\ 0 & C \end{bmatrix} = D.$$

Note that the matrix of T_W in the restricted basis β_W is just A . It can be shown that

$$\det(xI - D) = \det(xI - A) \det(xI - C),$$

which immediately gives the first result.

Now, it can also be shown that the powers of D are of the form

$$[T^k]_\beta = \begin{bmatrix} A^k & B_k \\ 0 & C^k \end{bmatrix} = D^k.$$

Thus, if $T^k\mathbf{v} = \mathbf{0}$ implies $T_W^k\mathbf{v} = \mathbf{0}$, hence any polynomial which annihilates T also annihilates T_W . This gives the second result. \square

Definition 1.10. Let W be an invariant subspace under $T \in \mathcal{L}(V)$, and let $\mathbf{v} \in V$. We define the T -conductor of \mathbf{v} into W as the set $S_T(\mathbf{v}; W)$ of all polynomials g such that $g(T)\mathbf{v} \in W$.

When $W = \{0\}$, $S_T(\mathbf{v}, \{0\})$ is called the T -annihilator of \mathbf{v} .

Lemma 1.13. *If W is invariant under T , then it is invariant under all polynomials of T . Thus, the conductor $S_T(\mathbf{v}, W)$ is an ideal in the ring of polynomials $\mathbb{F}[x]$.*

Definition 1.11. If W is an invariant subspace under $T \in \mathcal{L}(V)$, and $\mathbf{v} \in V$, then the unique monic generator of $S_T(\mathbf{v}, W)$ is also called the T -conductor of \mathbf{v} into W .

The unique monic generator of $S_T(\mathbf{v}, \{0\})$ is also called the T -annihilator of \mathbf{v} .

Remark. The T -annihilator of \mathbf{v} is the unique monic polynomial g of least degree such that $g(T)\mathbf{v} = \mathbf{0}$.

Remark. The minimal polynomial is a T -conductor for every $\mathbf{v} \in V$, thus every T -conductor divides the minimal polynomial of T .