

STAT6203: Theoretical Statistics III

Robust Statistics

Instructed by *Prof. Marco Avella Medina*

Transcribed by *Satvik Saha*

Department of Statistics, Columbia University

Table of Contents

1. Location estimation	2
1.1. Minimax Asymptotic Bias	3
1.2. Minimax Asymptotic Variance	4
Bibliography	6

1. Location estimation

Consider the problem of estimating an unknown location parameter ξ from a sample

$$X_1, \dots, X_n \stackrel{\text{iid}}{\sim} F(\cdot - \xi).$$

In general, we search for suitably *efficient* estimators. Huber (1964) seeks to obtain *robust* (in a certain sense) estimators which perform well when F is only being approximately known; in particular, suppose that F is a slightly ‘contaminated’ version of a known distribution F_0 .

Definition 1.1 (Tukey-Huber Contamination). Given $\varepsilon \in [0, 1]$ and a distribution F_0 , define

$$\mathcal{P}_\varepsilon(F_0) := \{(1 - \varepsilon)F_0 + \varepsilon H : \text{arbitrary distribution } H\}.$$

Remark. $\mathcal{P}_\varepsilon(F_0)$ is contained within the ε -Kolmogorov-Smirnov neighborhood of F_0 : for any $F = (1 - \varepsilon)F_0 + \varepsilon H$,

$$\|F - F_0\|_\infty = \|(1 - \varepsilon)F_0 + \varepsilon H - F_0\|_\infty = \varepsilon\|H - F_0\|_\infty \leq \varepsilon.$$

With this, we may examine how an estimator based for a parameter of F_0 suffers under the contaminated sample from $F \in \mathcal{P}_\varepsilon(F_0)$. As a first step, we may look at some simple population-level functionals.

Example 1.2. Consider the mean functional $\mu(G) := \mathbb{E}_G[X]$. The ‘bias’ of μ on $F := (1 - \varepsilon)F_0 + \varepsilon H$ against F_0 is

$$\mu(F) - \mu(F_0) = \varepsilon\mu(H) - \varepsilon\mu(F_0).$$

Note that this is completely unrestricted over $F \in \mathcal{P}_\varepsilon(F_0)$ even if we fix ε , owing to the arbitrary nature of H (which in principle may not even have a mean!).

Example 1.3. Consider the (left) median functional $m(G) := G^-(\frac{1}{2}) = \inf\{x : G(x) \geq \frac{1}{2}\}$, and let $\varepsilon < \frac{1}{2}$. For $F := (1 - \varepsilon)F_0 + \varepsilon H$, note that

$$m(F) = \inf\left\{x : (1 - \varepsilon)F_0(x) + \varepsilon H(x) \geq \frac{1}{2}\right\}.$$

Since $0 \leq H(x) \leq 1$, we can bound

$$\begin{aligned} F_0^-\left(\frac{\frac{1}{2} - \varepsilon}{1 - \varepsilon}\right) &= \inf\left\{x : (1 - \varepsilon)F_0(x) + \varepsilon \geq \frac{1}{2}\right\} \\ &\leq m(F) \\ &\leq \inf\left\{x : (1 - \varepsilon)F_0(x) \geq \frac{1}{2}\right\} = F_0^-\left(\frac{\frac{1}{2}}{1 - \varepsilon}\right). \end{aligned}$$

Unlike the previous example, the median $m(F)$ cannot deviate arbitrarily over $F \in \mathcal{P}_\varepsilon(F_0)$.

1.1. Minimax Asymptotic Bias

For the purposes of location estimation, it is natural to restrict ourselves to *translation invariant* estimators.

Definition 1.4 (Translation Invariant Estimators). We say that an estimator T is translation invariant if $T(X + a) = T(X) + a$ for all samples X and $a \in \mathbb{R}$. We denote the class of all translation invariant estimators by $\mathcal{T}$.

Definition 1.5 (Asymptotic Bias). Let $\{T_n\}$ be a sequence of estimators for $T(F_0)$. Its asymptotic bias is defined as

$$b(\{T_n\}, F) := \limsup_{n \rightarrow \infty} |\mathbb{E}_F[T_n] - T(F_0)|.$$

We can now choose our ‘best’ estimator on the basis of asymptotic bias; fix $\varepsilon > 0$ and consider the minimax problem

$$b_* := \min_{\{T_n\} \subset \mathcal{T}} \max_{F \in \mathcal{P}_\varepsilon(F_0)} b(\{T_n\}, F).$$

Theorem 1.6. Let F_0 have a density that is symmetric about zero and decreasing on $\mathbb{R}_+$, and let $T(F_0) = 0$. The sample medians $\{M_n\}$ solve the minimax asymptotic bias problem, with

$$\min_{\{T_n\} \subset \mathcal{T}} \max_{F \in \mathcal{P}_\varepsilon(F_0)} b(\{T_n\}, F) = F_0^{-1}\left(\frac{1/2}{1-\varepsilon}\right).$$

Proof. First note that the sample median estimator is consistent, and $\lim_{n \rightarrow \infty} \mathbb{E}_F[M_n] = F^{-1}(\frac{1}{2})$ under some mild conditions on F . We have already demonstrated in [Example 1.3](#) that $|F^{-1}(\frac{1}{2})| \leq F_0^{-1}(\frac{1/2}{1-\varepsilon}) =: m_*$ for all $F \in \mathcal{P}_\varepsilon(F_0)$, which gives us the upper bound $b_* \leq m_*$.

Next, we will construct two distributions $F_+, F_- \in \mathcal{P}_\varepsilon(F_0)$, and use

$$b_* \geq \min_{\{T_n\} \subset \mathcal{T}} \max(b(\{T_n\}, F_+), b(\{T_n\}, F_-)) \quad (\star)$$

to obtain the matching lower bound $b_* \geq m_*$. Define

$$F_+(x) := \begin{cases} (1-\varepsilon)F_0(x) & \text{if } x < m_*, \\ 1 - (1-\varepsilon)F_0(2m_* - x) & \text{if } x \geq m_*. \end{cases}$$

This describes a distribution which has the same shape as F_0 on $(-\infty, m_*)$, and is symmetric about m_* . Further set $F_-(x) = F_+(x + 2m_*)$, which is symmetric about $-m_*$. We can verify that F_+, F_- are indeed members of $\mathcal{P}_\varepsilon(F_0)$; solving for $F_+ = (1-\varepsilon)F_0 + \varepsilon H_+$ reveals that

$$H_+(x) = \frac{F_+(x) - (1-\varepsilon)F_0(x)}{\varepsilon} = \frac{1 - (1-\varepsilon)(F_0(x) + F_0(2m_* - x))}{\varepsilon} \mathbf{1}(x \geq m_*).$$

This is continuous with $H(-\infty) = 0$, $H(\infty) = 1$, and is nondecreasing since

$$F_0(x) + F_0(2m_* - x) = F_0(x) - F_0(x - 2m_*) + 1$$

is decreasing on $x \geq m_*$, which makes it a valid *cdf*. The argument for F_- , which is a reflection of F_+ about zero, is similar.

Now, if $b_* < m_*$, then m_* must be strictly greater than the right hand side of $(\star)$, hence there exists a sequence $\{T_n\} \subset \mathcal{T}$ such that

$$\max\left(\limsup_{n \rightarrow \infty} |\mathbb{E}_{F_+}[T_n]|, \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_-}[T_n]|\right) < m_*.$$

However, translation invariance of T_n forces $\mathbb{E}_{F_+}[T_n] = \mathbb{E}_{F_-}[T_n] + 2m_*$, hence

$$2m_* = \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_+}[T_n] - \mathbb{E}_{F_-}[T_n]| \leq \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_+}[T_n]| + \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_-}[T_n]| < 2m_*,$$

a contradiction! □

Remark. Some further technical conditions are required to ensure that $\mathbb{E}_F[M_n] \rightarrow F^{-}(\frac{1}{2})$, which we omit here.

1.2. Minimax Asymptotic Variance

Instead of relying on asymptotic bias, we will now examine the asymptotic variance of a special class of estimators.

Definition 1.7 (*M-estimator*). We say that T_n is an *M-estimator* of location if

$$T_n \in \arg \min_t \sum_{i=1}^n \rho(X_i - t)$$

for some function ρ . When ρ is differentiable, we denote $\psi := \rho'$, and T_n equivalently satisfies

$$\sum_{i=1}^n \psi(X_i - T_n) = 0.$$

Remark. *M-estimators* are translation invariant.

Lemma 1.8 (Asymptotic Normality of *M-estimators*). Let $\{T_n\}$ be the sequence of *M-estimators* associated with ψ . Define the maps $\lambda(t) := \mathbb{E}_F[\psi(X - t)]$, $\sigma^2(t) := \text{var}_F[\psi(X - t)]$, and suppose that there exists $t_0 \in \mathbb{R}$ such that the following are satisfied.

1. λ is continuous in a neighborhood of t_0 , with $\lambda(t_0) = 0$.
2. λ is differentiable at t_0 , with $\lambda'(t_0) < 0$.
3. σ^2 is continuous, finite, non-zero in a neighborhood of t_0 .

Then $\sqrt{n}(T_n - t_0) \xrightarrow{d} \mathcal{N}(0, V(\psi, F))$, where

$$V(\psi, F) := \frac{\sigma^2(t_0)}{(\lambda'(t_0))^2}$$

is the asymptotic variance of $\{T_n\}$.

Example 1.9. The M -estimator associated with $\rho(x) = x^2$ is the sample mean, with asymptotic variance $\text{var}_F(X)$.

Example 1.10. The M -estimator associated with $\rho(x) = |x|$ is the sample median, with asymptotic variance $1/4\left(f\left(F^{-}\left(\frac{1}{2}\right)\right)\right)^2$.

From now on, we will restrict ourselves to the contamination neighborhood

$$\mathcal{P}_\varepsilon^{\text{sym}}(F_0) := \{(1 - \varepsilon)F_0 + \varepsilon H : \text{symmetric } H\}$$

to ensure that $F \in \mathcal{P}_\varepsilon^{\text{sym}}(F_0)$ is symmetric (about zero), and further select $F_0 = \Phi$.

Consider the problem

$$V_* := \min_{\psi} \max_{F \in \mathcal{P}_\varepsilon^{\text{sym}}(\Phi)} V(\psi, F).$$

Theorem 1.11. *Define the Huber loss*

$$\rho_c(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } |x| < c, \\ c|x| - \frac{1}{2}c^2 & \text{if } |x| \geq c. \end{cases}$$

The sequence of M -estimators associated with ρ_c solve the minimax asymptotic variance problem when c satisfies

$$2\left(\frac{\phi(c)}{c} - \Phi(c)\right) = \frac{\varepsilon}{1 - \varepsilon}.$$

Bibliography

Huber, P. J. (1964). Robust Estimation of a Location Parameter. *The Annals of Mathematical Statistics*, 35(1), 73–101. <https://doi.org/10.1214/aoms/1177703732>