

Robust Statistics

Instructed by *Prof. Marco Avella Medina*

Transcribed by *Satvik Saha*

Department of Statistics, Columbia University

Table of Contents

1. Location Estimation	2
1.1. Minimax Asymptotic Bias	3
1.2. Minimax Asymptotic Variance	4
2. Quantification of Robustness	7
2.1. Influence Functions	7
2.2. Breakdown Points	8
3. Location and Scale Estimation	10
3.1. The Univariate Setting	10
3.2. The Multivariate Setting	11
4. Robust Regression	13
4.1. Mean-Shift Model	13
Bibliography	15

1. Location Estimation

Consider the problem of estimating an unknown location parameter ξ from a sample

$$X_1, \dots, X_n \stackrel{\text{iid}}{\sim} F(\cdot - \xi).$$

In general, we search for suitably *efficient* estimators. Huber (1964) seeks to obtain *robust* (in a certain sense) estimators which perform well when F is only being approximately known; in particular, suppose that F is a slightly ‘contaminated’ version of a known distribution F_0 .

Definition 1.1 (Tukey-Huber Contamination). Given $\varepsilon \in [0, 1]$ and a distribution F_0 , define

$$\mathcal{P}_\varepsilon(F_0) := \{(1 - \varepsilon)F_0 + \varepsilon H : \text{arbitrary distribution } H\}.$$

Remark. $\mathcal{P}_\varepsilon(F_0)$ is contained within the ε -Kolmogorov-Smirnov neighborhood of F_0 : for any $F = (1 - \varepsilon)F_0 + \varepsilon H$,

$$\|F - F_0\|_\infty = \|(1 - \varepsilon)F_0 + \varepsilon H - F_0\|_\infty = \varepsilon\|H - F_0\|_\infty \leq \varepsilon.$$

With this, we may examine how an estimator based for a parameter of F_0 suffers under the contaminated sample from $F \in \mathcal{P}_\varepsilon(F_0)$. As a first step, we may look at some simple population-level functionals.

Example 1.2. Consider the mean functional $\mu(G) := \mathbb{E}_G[X]$. The ‘bias’ of μ on $F := (1 - \varepsilon)F_0 + \varepsilon H$ against F_0 is

$$\mu(F) - \mu(F_0) = \varepsilon\mu(H) - \varepsilon\mu(F_0).$$

Note that this is completely unrestricted over $F \in \mathcal{P}_\varepsilon(F_0)$ even if we fix ε , owing to the arbitrary nature of H (which in principle may not even have a mean!).

Example 1.3. Consider the (left) median functional $m(G) := G^-(\frac{1}{2}) = \inf\{x : G(x) \geq \frac{1}{2}\}$, and let $\varepsilon < \frac{1}{2}$. For $F := (1 - \varepsilon)F_0 + \varepsilon H$, note that

$$m(F) = \inf\left\{x : (1 - \varepsilon)F_0(x) + \varepsilon H(x) \geq \frac{1}{2}\right\}.$$

Since $0 \leq H(x) \leq 1$, we can bound

$$\begin{aligned} F_0^-\left(\frac{\frac{1}{2} - \varepsilon}{1 - \varepsilon}\right) &= \inf\left\{x : (1 - \varepsilon)F_0(x) + \varepsilon \geq \frac{1}{2}\right\} \\ &\leq m(F) \\ &\leq \inf\left\{x : (1 - \varepsilon)F_0(x) \geq \frac{1}{2}\right\} = F_0^-\left(\frac{\frac{1}{2}}{1 - \varepsilon}\right). \end{aligned}$$

Unlike the previous example, the median $m(F)$ cannot deviate arbitrarily over $F \in \mathcal{P}_\varepsilon(F_0)$.

1.1. Minimax Asymptotic Bias

For the purposes of location estimation, it is natural to restrict ourselves to *translation invariant* estimators.

Definition 1.4 (Translation Invariant Estimators). We say that an estimator T is translation invariant if $T(X + a) = T(X) + a$ for all samples X and $a \in \mathbb{R}$. We denote the class of all translation invariant estimators by $\mathcal{T}$.

Definition 1.5 (Asymptotic Bias). Let $\{T_n\}$ be a sequence of estimators for $T(F_0)$. Its asymptotic bias is defined as

$$b(\{T_n\}, F) := \limsup_{n \rightarrow \infty} |\mathbb{E}_F[T_n] - T(F_0)|.$$

We can now choose our ‘best’ estimator on the basis of asymptotic bias; fix $\varepsilon > 0$ and consider the minimax problem

$$b_* := \min_{\{T_n\} \subset \mathcal{T}} \max_{F \in \mathcal{P}_\varepsilon(F_0)} b(\{T_n\}, F).$$

Theorem 1.6. Let F_0 have a density that is symmetric about zero and decreasing on $\mathbb{R}_+$, and let $T(F_0) = 0$. The sample medians $\{M_n\}$ solve the minimax asymptotic bias problem, with

$$\min_{\{T_n\} \subset \mathcal{T}} \max_{F \in \mathcal{P}_\varepsilon(F_0)} b(\{T_n\}, F) = F_0^{-1}\left(\frac{1/2}{1-\varepsilon}\right).$$

Proof. First note that the sample median estimator is consistent, and $\lim_{n \rightarrow \infty} \mathbb{E}_F[M_n] = F^{-1}(\frac{1}{2})$ under some mild conditions on F . We have already demonstrated in [Example 1.3](#) that $|F^{-1}(\frac{1}{2})| \leq F_0^{-1}(\frac{1/2}{1-\varepsilon}) =: m_*$ for all $F \in \mathcal{P}_\varepsilon(F_0)$, which gives us the upper bound $b_* \leq m_*$.

Next, we will construct two distributions $F_+, F_- \in \mathcal{P}_\varepsilon(F_0)$, and use

$$b_* \geq \min_{\{T_n\} \subset \mathcal{T}} \max(b(\{T_n\}, F_+), b(\{T_n\}, F_-)) \quad (\star)$$

to obtain the matching lower bound $b_* \geq m_*$. Define

$$F_+(x) := \begin{cases} (1-\varepsilon)F_0(x) & \text{if } x < m_*, \\ 1 - (1-\varepsilon)F_0(2m_* - x) & \text{if } x \geq m_*. \end{cases}$$

This describes a distribution which has the same shape as F_0 on $(-\infty, m_*)$, and is symmetric about m_* . Further set $F_-(x) = F_+(x + 2m_*)$, which is symmetric about $-m_*$. We can verify that F_+, F_- are indeed members of $\mathcal{P}_\varepsilon(F_0)$; solving for $F_+ = (1-\varepsilon)F_0 + \varepsilon H_+$ reveals that

$$H_+(x) = \frac{F_+(x) - (1-\varepsilon)F_0(x)}{\varepsilon} = \frac{1 - (1-\varepsilon)(F_0(x) + F_0(2m_* - x))}{\varepsilon} \mathbf{1}(x \geq m_*).$$

This is continuous with $H(-\infty) = 0$, $H(\infty) = 1$, and is nondecreasing since

$$F_0(x) + F_0(2m_* - x) = F_0(x) - F_0(x - 2m_*) + 1$$

is decreasing on $x \geq m_*$, which makes it a valid *cdf*. The argument for F_- , which is a reflection of F_+ about zero, is similar.

Now, if $b_* < m_*$, then m_* must be strictly greater than the right hand side of $(\star)$, hence there exists a sequence $\{T_n\} \subset \mathcal{T}$ such that

$$\max\left(\limsup_{n \rightarrow \infty} |\mathbb{E}_{F_+}[T_n]|, \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_-}[T_n]|\right) < m_*.$$

However, translation invariance of T_n forces $\mathbb{E}_{F_+}[T_n] = \mathbb{E}_{F_-}[T_n] + 2m_*$, hence

$$2m_* = \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_+}[T_n] - \mathbb{E}_{F_-}[T_n]| \leq \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_+}[T_n]| + \limsup_{n \rightarrow \infty} |\mathbb{E}_{F_-}[T_n]| < 2m_*,$$

a contradiction! □

Remark. Some further technical conditions are required to ensure that $\mathbb{E}_F[M_n] \rightarrow F^-(\frac{1}{2})$, which we omit here.

1.2. Minimax Asymptotic Variance

Instead of relying on asymptotic bias, we will now examine the asymptotic variance of a special class of estimators.

Definition 1.7 (M-estimator of Location). We say that T_n is an M-estimator of location if

$$T_n \in \arg \min_t \sum_{i=1}^n \rho(X_i - t)$$

for some function ρ . When ρ is differentiable, we denote $\psi := \rho'$, and T_n equivalently satisfies

$$\sum_{i=1}^n \psi(X_i - T_n) = 0.$$

Note that M-estimators are automatically translation invariant. We will generally assume that the loss function ρ is symmetric and convex from now on.

Lemma 1.8 (Asymptotic Normality of M-estimators). *Let $\{T_n\}$ be the sequence of M estimators associated with ψ . Define the maps $\lambda(t) := \mathbb{E}_F[\psi(X - t)]$, $\sigma^2(t) := \text{var}_F[\psi(X - t)]$, and suppose that there exists $t_0 \in \mathbb{R}$ such that the following are satisfied.*

1. λ is continuous in a neighborhood of t_0 , with $\lambda(t_0) = 0$.
2. λ is differentiable at t_0 , with $\lambda'(t_0) < 0$.
3. σ^2 is continuous, finite, non-zero in a neighborhood of t_0 .

Then $\sqrt{n}(T_n - t_0) \xrightarrow{d} \mathcal{N}(0, V(\psi, F))$, where

$$V(\psi, F) := \frac{\sigma^2(t_0)}{(\lambda'(t_0))^2}$$

is the asymptotic variance of $\{T_n\}$.

Example 1.9. The M-estimator associated with $\rho(x) = x^2$ is the sample mean, with asymptotic variance $\text{var}_F(X)$.

Example 1.10. The M-estimator associated with $\rho(x) = |x|$ is the sample median, with asymptotic variance $1/4(f(F^{-1}(\frac{1}{2})))^2$.

Example 1.11. The M-estimator associated with $\rho(x) = -\log f(x)$ is the MLE of ξ in the location family $F(\cdot - \xi)$.

From now on, we will restrict ourselves to the contamination neighborhood

$$\mathcal{P}_\varepsilon^{\text{sym}}(F_0) := \{(1 - \varepsilon)F_0 + \varepsilon H : \text{symmetric } H\}$$

to ensure that $F \in \mathcal{P}_\varepsilon^{\text{sym}}(F_0)$ is symmetric (about zero), and further select $F_0 = \Phi$.

Consider the problem

$$V_* := \min_{\psi} \max_{F \in \mathcal{P}_\varepsilon^{\text{sym}}(\Phi)} V(\psi, F),$$

where ψ ranges over functions satisfying the requirements of [Lemma 1.8](#), along with symmetry and convexity so that $t_0 = 0$.

Theorem 1.12. Define the Huber loss

$$\rho_c(x) = \begin{cases} \frac{1}{2}x^2 & \text{if } |x| < c, \\ c|x| - \frac{1}{2}c^2 & \text{if } |x| \geq c. \end{cases}$$

The sequence of M-estimators associated with ρ_c solve the minimax asymptotic variance problem when c satisfies

$$2\left(\frac{\phi(c)}{c} - \Phi(c)\right) = \frac{\varepsilon}{1 - \varepsilon}.$$

Remark. This corresponds to

$$\psi_c(x) := \rho'_c(x) = \begin{cases} x & \text{if } |x| < c, \\ c \text{sign}(x) & \text{if } |x| \geq c. \end{cases}$$

Lemma 1.13. Suppose that $F_* \in \mathcal{P}_\varepsilon^{\text{sym}}(\Phi)$ has density f_* such that

$$F_* \in \arg \max_{F \in \mathcal{P}_\varepsilon^{\text{sym}}(\Phi)} V(\psi_*, F)$$

where $\psi_* = -f'_*/f_*$. Then ψ_* solves the minimax asymptotic variance solution, and

$$\min_{\psi} \max_{F \in \mathcal{P}_\varepsilon^{\text{sym}}(\Phi)} V(\psi, F) = V(\psi_*, F_*).$$

Proof. We already have

$$\min_{\psi} \max_{F \in \mathcal{P}_{\varepsilon}^{\text{sym}}(\Phi)} V(\psi, F) \leq \max_{F \in \mathcal{P}_{\varepsilon}^{\text{sym}}(\Phi)} V(\psi_*, F) = V(\psi_*, F_*).$$

To establish the other direction, we will show that $\psi_* \in \arg \min_{\psi} V(\psi, F_*)$, whence

$$\min_{\psi} \max_{F \in \mathcal{P}_{\varepsilon}^{\text{sym}}(\Phi)} V(\psi, F) \geq \min_{\psi} V(\psi, F_*) = V(\psi_*, F_*).$$

To this end, compute

$$\mathbb{E}_{F_*}[\psi'] = \int \psi' f_* = - \int \psi f_*' = \int \psi \psi_* f_* = \mathbb{E}_{F_*}[\psi \psi_*] \leq \sqrt{\mathbb{E}_{F_*}[\psi^2] \mathbb{E}_{F_*}[\psi_*^2]}.$$

In particular, $\mathbb{E}_{F_*}[\psi'] = \mathbb{E}_{F_*}[\psi_*^2]$. Thus,

$$V(\psi, F_*) = \frac{\mathbb{E}_{F_*}[\psi^2]}{(\mathbb{E}_{F_*}[\psi'])^2} \geq \frac{\mathbb{E}_{F_*}[\psi^2]}{\mathbb{E}_{F_*}[\psi^2] \mathbb{E}_{F_*}[\psi_*^2]} = \frac{1}{\mathbb{E}_{F_*}[\psi_*^2]} = V(\psi_*, F_*). \quad \square$$

Proof of Theorem 1.12. Set

$$f_c(x) := \frac{1-\varepsilon}{\sqrt{2\pi}} \exp(-\rho_c(x)) = \begin{cases} (1-\varepsilon)\phi(x) & \text{if } |x| < c, \\ \frac{1-\varepsilon}{\sqrt{2\pi}} \exp(-c|x| + \frac{1}{2}c^2) & \text{if } |x| \geq c, \end{cases}$$

and check that our choice of c ensures that f_c is a probability density function. The corresponding distribution F_c can be shown to belong to $\mathcal{P}_{\varepsilon}^{\text{sym}}(\Phi)$, by isolating

$$h_c(x) := \frac{f_c(x) - (1-\varepsilon)\phi}{\varepsilon} \geq 0$$

and checking that this is also a probability density function. Now, for any $F := (1-\varepsilon)\Phi + \varepsilon H \in \mathcal{P}_{\varepsilon}^{\text{sym}}(\Phi)$, we have

$$V(\psi_c, F) = \frac{\mathbb{E}_F[\psi_c^2]}{(\mathbb{E}_F[\psi_c'])^2} = \frac{(1-\varepsilon)\mathbb{E}_{\Phi}[\psi_c^2] + \varepsilon\mathbb{E}_H[\psi_c^2]}{((1-\varepsilon)\mathbb{E}_{\Phi}[\psi_c'] + \varepsilon\mathbb{E}_H[\psi_c'])^2}.$$

which attains its maximum precisely when $\mathbb{E}_H[\psi_c^2] = c^2$ and $\mathbb{E}_H[\psi_c'] = 0$. However, these are satisfied by H_c (supported on $\{|x| \geq c\}$, where $\psi_c(x) = c \text{sign}(x)$), whence $V(\psi_c, F) \leq V(\psi_c, F_c)$. We have shown that F_c satisfies the conditions of Lemma 1.13, which directly gives the result. $\square$

2. Quantification of Robustness

2.1. Influence Functions

Definition 2.1 (Influence Function). The influence function of a functional T at x with respect to a distribution F is defined by

$$\text{IF}(x; T, F) = \lim_{\varepsilon \rightarrow 0^+} \frac{T((1 - \varepsilon)F + \varepsilon\delta_x) - T(F)}{\varepsilon} = \left. \frac{d}{d\varepsilon} T((1 - \varepsilon)F + \varepsilon\delta_x) \right|_{\varepsilon=0}.$$

Remark. The influence function $\text{IF}(x; T, F)$ is the Gateaux derivative of T at F in the direction $\delta_x - F$.

The influence function $\text{IF}(x; T, F)$ quantifies infinitesimal effect on $T(F)$ of adding a new point x to F . We will call functionals with bounded influence functions *infinitesimally robust*.

Example 2.2. Let T be a linear functional. Denoting $F_t := (1 - t)F + t\delta_x$, we have

$$T(F_t) = (1 - t)T(F) + tT(\delta_x) \implies \text{IF}(x; T, F) = T(\delta_x) - T(F).$$

For instance, the influence function of the mean functional $\mu(F) := \mathbb{E}_F[X]$ is described by

$$\text{IF}(\mu(F) + x; \mu, F) = x.$$

Note that this is unbounded!

Example 2.3. Let $m(G) := G^{-1}(\frac{1}{2})$ be the median functional. Denoting $F_t := (1 - t)F + t\delta_x$, note that we must have $F_t(m(F_t)) = \frac{1}{2}$, which expands to

$$(1 - t)F(m(F_t)) + t\mathbf{1}(m(F_t) \geq x).$$

Differentiating this implicit equation and setting $t = 0$ gives

$$-\frac{1}{2} + f(m(F)) \text{IF}(x; m, F) + \mathbf{1}(m(F) \geq x) = 0 \implies \text{IF}(x; m, F) = \frac{\text{sign}(x - m(F))}{2f(m(F))}.$$

Example 2.4. Let T be an M-estimator associated with ψ , obeying $\mathbb{E}_F[\psi(X, T(F))] = 0$. Plugging in $F_t := (1 - t)F + t\delta_x$ gives

$$(1 - t)\mathbb{E}_F[\psi(X, T(F_t))] + t\psi(x, T(F_t)) = 0.$$

Differentiating at $t = 0$ gives

$$-\cancel{\mathbb{E}_F[\psi(X, T(F))]} + (1 - t)\mathbb{E}_F[\psi'(X, T(F))] \text{IF}(x; T, F) + \psi(x, T(F)) + t\psi'(x, T(F)) \text{IF}(x; T, F) = 0,$$

hence

$$\text{IF}(x; T, F) = M^{-1}\mathbb{E}_F[\psi(x, T(F))], \quad M := -\mathbb{E}_F[\psi'(x, T(F))].$$

The influence function can be interpreted as a certain limit of the *sensitivity curve*. For $X_i \stackrel{\text{iid}}{\sim} F$, with estimators $T_n \xrightarrow{p} T(F)$, and under suitable regularity conditions, we have

$$\text{SC}(x, T) := n(T_{n+1}(X_1, \dots, X_n, x) - T_n(X_1, \dots, X_n)) \rightarrow \text{IF}(x; T, F).$$

2.2. Breakdown Points

Definition 2.5 (Breakdown Point, Asymptotic). The asymptotic breakdown point of T is

$$\varepsilon^*(T, F) := \sup \left\{ \varepsilon : \sup_H \|T((1 - \varepsilon)F + \varepsilon H) - T(F)\| < \infty \right\}.$$

Example 2.6. Consider the mean functional $\mu(F) := \mathbb{E}_F[X]$. Then,

$$\mu((1 - \varepsilon)F + \varepsilon H) - \mu(F) = \varepsilon(\mu(H) - \mu(F))$$

is unbounded over all distributions H whenever $\varepsilon > 0$. This means that $\varepsilon^*(\mu, F) = 0$.

Theorem 2.7. Let T be an M -estimator of location associated with ψ , with ψ nondecreasing, $k_1 := -\psi(-\infty)$, $k_2 := \psi(\infty)$ both finite. Then,

$$\varepsilon^*(T, F) = \frac{\min(k_1, k_2)}{k_1 + k_2}.$$

In particular, $\varepsilon^*(T, F) = \frac{1}{2}$ when $\psi(-\infty) + \psi(\infty) = 0$.

Example 2.8. For the median functional $m(F) := F^{-1}(\frac{1}{2})$, $\varepsilon^*(m, F) = \frac{1}{2}$.

Definition 2.9 (Breakdown Point, Finite Sample). The finite sample breakdown point of a statistic T_n at $X := (X_1, \dots, X_n)$ is

$$\hat{\varepsilon}(T_n, X) := \sup \left\{ \frac{m}{n} : \sup_{d_H(X, X') \leq m} \|T_n(X') - T_n(X)\| < \infty \right\}.$$

Remark. Here, d_H denotes the *Hamming distance*, with

$$d_H(X, X') := \sum_{i=1}^n \mathbf{1}(X_i \neq X'_i).$$

We may call X' an m -replacement of X when $d_H(X, X') = m$.

Remark. The sample median M_n achieves the largest possible finite sample breakdown point

$$\hat{\varepsilon}(M_n, X) = \frac{1}{n} \left\lfloor \frac{n}{2} - 1 \right\rfloor$$

among all translation invariant estimators.

3. Location and Scale Estimation

3.1. The Univariate Setting

Pure scale estimation involves considering iid $X_1, \dots, X_n$ from the the scale family

$$f(x; \sigma) = \frac{1}{\sigma} f\left(\frac{x}{\sigma}\right).$$

An MLE of σ would satisfy the score equation

$$\sum_{i=1}^n \frac{\partial}{\partial x} \log f(x; \sigma) = \sum_{i=1}^n -\frac{x}{\sigma^2} \frac{f'(x/\sigma)}{f(x/\sigma)} - \frac{1}{\sigma} = 0.$$

Definition 3.1 (M-estimator of Scale). We say that S_n is an M-estimator of scale if it satisfies

$$\sum_{i=1}^n \chi\left(\frac{X_i}{S_n}\right) = 0$$

for some function χ .

This corresponds to the population functional S satisfying

$$\mathbb{E}_F\left[\chi\left(\frac{X}{S(F)}\right)\right] = 0.$$

Remark. We can compute

$$\text{IF}(x; S, F) = \frac{\chi(x/S(F))}{\mathbb{E}_F[\chi'(X/S(F))(X/S(F))]}.$$

An infinitesimally robust M-estimator of scale requires bounded χ .

Example 3.2. Let $X_i \stackrel{\text{iid}}{\sim} N(0, \sigma^2)$, and consider the M-estimator described by

$$\chi_c(x) := \begin{cases} x^2 + \kappa_c & \text{if } |x| \leq c, \\ c^2 + \kappa_c & \text{if } |x| > c, \end{cases}$$

where κ_c is a *Fisher consistency constant* chosen such that $\mathbb{E}_{N(0,1)}[\chi_c(Z)] = 0$. Note that the symmetry of the location problem saved us from this complication earlier!

More generally, for iid $X_1, \dots, X_n$ from a location-scale family of the form

$$f(x; \sigma) = \frac{1}{\sigma} f\left(\frac{x - \mu}{\sigma}\right),$$

we may consider estimators (T_n, S_n) which satisfy equations

$$\sum_{i=1}^n \psi\left(\frac{X_i - T_n}{S_n}\right) = 0, \quad \sum_{i=1}^n \chi\left(\frac{X_i - T_n}{S_n}\right) = 0$$

as joint M-estimators of location and scale.

Example 3.3. Picking $\rho = \rho_c$, $\chi = \chi_c$ amounts to solving

$$(T_n, S_n) \in \arg \min_{\mu, \sigma} \left\{ \sum_{i=1}^n \sigma \rho_c \left(\frac{X_i - \mu}{\sigma} \right) + \sigma \beta n \right\}$$

with an appropriate Fisher consistency constant β .

Example 3.4. It is also possible to perform estimation in two stages, by first estimating the scale, then plugging that into a location estimation problem. The *Mean Absolute Deviation (MAD)* estimator is a common choice for this initial scale estimate.

3.2. The Multivariate Setting

Here, we will consider multivariate distributions on $\mathbb{R}^p$.

Definition 3.5. An elliptical family of distributions has densities of the form

$$f(x; \mu, \Sigma) = \frac{1}{|\Sigma|^{\frac{1}{2}}} f((x - \mu)^\top \Sigma^{-1} (x - \mu)).$$

The quantity $d(\cdot; \mu, \Sigma) := \sqrt{\|x - \mu\|_{\Sigma^{-1}}^2}$ is the *Mahalanobis distance*.

We may consider estimators (T_n, S_n) satisfying equations of the form

$$\begin{aligned} \sum_{i=1}^n w_1(d_i^2)(X_i - T_n) &= 0, \\ \frac{1}{n} \sum_{i=1}^n w_2(d_i^2)(X_i - T_n)(X_i - T_n)^\top &= S_n, \end{aligned}$$

where $d_i^2 := d^2(X_i; T_n, S_n)$. Note that these produce *affine invariant estimators*. As usual, we may construct the corresponding population-level functionals

$$\begin{aligned} \mathbb{E}_F[w_1(d^2(X))(X - T(F))] &= 0, \\ \mathbb{E}_F[w_2(d^2(X))(X - T(F))(X - T(F))^\top] &= S(F). \end{aligned}$$

Theorem 3.6 (Maronna (1976)). Suppose that $w_2(d^2)d^2$ is nondecreasing with a finite supremum k , and $w_1(d^2)d < \infty$. Then, the corresponding location-scale functionals (T, S) have breakdown point

$$\varepsilon^*((T, S); F) \leq \min \left\{ \frac{1}{k}, 1 - \frac{p}{k} \right\} \leq \frac{1}{p+1}.$$

In this setting, the right notion of breakdown involves T escaping to ∞ as well as the minimum/maximum eigenvalues of S escaping to $0/\infty$:

$$\varepsilon^*(T, S; F) := \sup \left\{ \varepsilon : \sup_{G \in \mathcal{P}_\varepsilon(F)} \|T(G) - T(F)\| < \infty, \right. \\ \left. \sup_{G \in \mathcal{P}_\varepsilon(F)} \lambda_{\max}(S(G)) < \infty, \sup_{G \in \mathcal{P}_\varepsilon(F)} \lambda_{\min}(S(G)) > 0 \right\}.$$

The Mahalanobis distance in the previous construction can be swapped out for other notions of outlyingness. The Stahel-Donoho estimator uses a projection-based outlyingness of the form

$$o(x; F) := \sup_{u \in S^{d-1}} \left| \frac{u^\top x - T_n^1(u^\top x; F_u)}{S_n^1(u^\top x; F_u)} \right|$$

where F_u is the distribution of $u^\top X$ with $X \sim F$, and T_n^1, S_n^1 are (robust) univariate location and scale estimators. This leads to weighted mean and covariance estimators

$$T_n(X) := \frac{\sum_{i=1}^n w_1(o(X_i)) X_i}{\sum_{i=1}^n w_1(o(X_i))}, \\ S_n(X) := \frac{\sum_{i=1}^n w_2(o(X_i)) (X_i - T_n(X))(X_i - T_n(X))^\top}{\sum_{i=1}^n w_2(o(X_i))}.$$

Under conditions analogous to [Theorem 3.6](#), these estimators can be shown to achieve a breakdown point of $\frac{1}{2}$, avoiding the dimensional penalty.

4. Robust Regression

Consider

$$y_i = x_i^\top \beta + \varepsilon_i$$

where $\{\varepsilon_i\}$ are iid random variables such that $\varepsilon_i \perp x_i$, $\mathbb{E}[\varepsilon_i] = 0$, and $\text{var}[\varepsilon_i] = \sigma^2$. Recall that ordinary least-squares linear regression solves

$$\hat{\beta}_{\text{OLS}} \in \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^n \|y_i - x_i^\top \beta\|_2^2.$$

One obvious extension is to instead optimize

$$\hat{\beta} \in \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^n \rho(y_i - x_i^\top \beta). \iff \sum_{i=1}^n \psi(x_i - x_i^\top \hat{\beta}) x_i = 0.$$

Under suitable conditions on the loss ρ and the noise ε , we recover asymptotic normality

$$\sqrt{n}(\hat{\beta} - \beta) \xrightarrow{d} \mathcal{N}\left(0, \frac{\mathbb{E}_F[\psi^2(\varepsilon_1)]}{(\mathbb{E}_F[\psi'(\varepsilon_1)])^2} \mathbb{E}_F[x_1 x_1^\top]^{-1}\right).$$

The general theory of M-estimators shows that using the Huber loss gives the minimax variance optimal estimator for symmetric contaminated noise ε_i . The influence function for (the population version of) these estimators take the form

$$\text{IF}((x, y); T, F) = -(\mathbb{E}_F[\psi'(\varepsilon_1)] \mathbb{E}_F[x_1 x_1^\top])^{-1} \psi(y - x^\top T(F)) x.$$

This shows that this setup is infinitesimally robust in the space of residuals, but *not* in the space of covariates x !

Lemma 4.1. *Let ψ be nondecreasing. The breakdown point of the regression estimator solving $\sum_{i=1}^n \psi(y_i - x_i^\top T_n) x_i = 0$ is zero.*

4.1. Mean-Shift Model

Consider an extension of the previous model of the form

$$y_i = x_i^\top \beta + \gamma_i + \varepsilon_i,$$

introducing a sparse parameter γ ; we intend for $\gamma_i \neq 0$ when the linear model does not hold (for instance, because of the presence of outliers). One choice is a LASSO-type estimator

$$(\hat{\beta}_\lambda, \hat{\gamma}_\lambda) \in \arg \min_{\beta, \gamma} \left\{ \frac{1}{2} \|y - X\beta - \gamma\|_2^2 + \lambda \|\gamma\|_1 \right\}$$

for some $\lambda \geq 0$.

Theorem 4.2. *The solution $\hat{\beta}_\lambda$ coincides with the Huber estimator obtained from*

$$\hat{\beta} \in \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^n \rho_\lambda(y_i - x_i^\top \beta).$$

Bibliography

- Hampel, F. R. (1974). The Influence Curve and Its Role in Robust Estimation. *Journal of the American Statistical Association*, 69(346), 383–393. <https://doi.org/10.2307/2285666>
- Huber, P. J. (1964). Robust Estimation of a Location Parameter. *The Annals of Mathematical Statistics*, 35(1), 73–101. <https://doi.org/10.1214/aoms/1177703732>
- Maronna, R. A. (1976). Robust M-Estimators of Multivariate Location and Scatter. *The Annals of Statistics*, 4(1), 51–67. <https://www.jstor.org/stable/2957994>