

# Olympiad Inequalities

Satvik Saha

## Theorems

**Theorem 1 (Trivial Inequality)** For all  $x \in \mathbb{R}$ :

$$x^2 \geq 0$$

**Theorem 2 (QM-AM-GM-HM)** For non-negative real numbers  $a_1, a_2, \dots, a_n$ , the following inequality holds:

$$\sqrt{\frac{a_1^2 + a_2^2 + \dots + a_n^2}{n}} \geq \frac{a_1 + a_2 + \dots + a_n}{n} \geq \sqrt[n]{a_1 a_2 \dots a_n} \geq \frac{n}{\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n}}$$

Equality holds if and only if  $a_1 = a_2 = \dots = a_n$ .

**Theorem 3 (Bernoulli)** For all  $r \geq 1$  and  $x \geq -1$ :

$$(1+x)^r \geq 1+rx$$

**Theorem 4 (Schur)** Let  $a, b, c$  be positive real numbers and  $r > 0$ . The following inequality holds:

$$a^r(a-b)(a-c) + b^r(b-c)(b-a) + c^r(c-a)(c-b) \geq 0$$

**Theorem 5 (Nesbitt)** Let  $a, b, c$  be positive real numbers. The following inequality holds:

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2}$$

Equality holds if  $a = b = c$ .

**Theorem 6 (Titu's Lemma)** Let  $(a_1, a_2, \dots, a_n)$  and  $(b_1, b_2, \dots, b_n)$  be sequences of non-negative real numbers. The following inequality holds:

$$\frac{a_1^2}{b_1} + \frac{a_2^2}{b_2} + \dots + \frac{a_n^2}{b_n} \geq \frac{(a_1 + a_2 + \dots + a_n)^2}{b_1 + b_2 + \dots + b_n}$$

Equality holds if  $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \dots = \frac{a_n}{b_n}$ .

**Theorem 7 (Cauchy-Schwarz)** Let  $(a_1, a_2, \dots, a_n)$  and  $(b_1, b_2, \dots, b_n)$  be sequences of non-negative real numbers. The following inequality holds:

$$(a_1^2 + a_2^2 + \dots + a_n^2)(b_1^2 + b_2^2 + \dots + b_n^2) \geq (a_1b_1 + a_2b_2 + \dots + a_nb_n)^2$$

Equality holds if and only if  $\frac{a_1}{b_1} = \frac{a_2}{b_2} = \dots = \frac{a_n}{b_n}$ .

**Theorem 8 (Hölder)** Let  $(a_1, a_2, \dots, a_n), (b_1, b_2, \dots, b_n), \dots, (z_1, z_2, \dots, z_n)$  be sequences of non-negative real numbers, and  $\lambda_a, \lambda_b, \dots, \lambda_z$  be positive reals which sum to 1. The following inequality holds:

$$(a_1 + a_2 + \dots + a_n)^{\lambda_a} (b_1 + b_2 + \dots + b_n)^{\lambda_b} \dots (z_1 + z_2 + \dots + z_n)^{\lambda_z} \geq (a_1^{\lambda_a} b_1^{\lambda_b} \dots z_1^{\lambda_z} + \dots + a_n^{\lambda_a} b_n^{\lambda_b} \dots z_n^{\lambda_z})$$

**Theorem 9 (Rearrangement)** Let  $(a_1 \leq a_2 \leq \dots \leq a_n)$  and  $(b_1 \leq b_2 \leq \dots \leq b_n)$  be sequences of non-decreasing real numbers. For any permutation  $\pi$  of  $\{1, 2, \dots, n\}$ , the following inequality holds:

$$a_1b_1 + a_2b_2 + \dots + a_nb_n \geq a_1b_{\pi(1)} + a_2b_{\pi(2)} + \dots + a_nb_{\pi(n)} \geq a_1b_n + a_2b_{n-1} + \dots + a_nb_1$$

**Theorem 10 (Chebyshev)** Let  $(a_1 \leq a_2 \leq \dots \leq a_n)$  and  $(b_1 \leq b_2 \leq \dots \leq b_n)$  be sequences of non-decreasing real numbers. The following inequality holds:

$$\frac{a_1b_1 + a_2b_2 + \dots + a_nb_n}{n} \geq \frac{a_1 + a_2 + \dots + a_n}{n} \cdot \frac{b_1 + b_2 + \dots + b_n}{n} \geq \frac{a_1b_n + a_2b_{n-1} + \dots + a_nb_1}{n}$$

**Theorem 11 (Muirhead)** Let the sequence  $(a_1, a_2, \dots, a_n)$  majorize<sup>1</sup>  $(b_1, b_2, \dots, b_n)$ . Then, for a sequence of positive reals  $(x_1, x_2, \dots, x_n)$ , the following inequality holds:

$$\sum_{sym} x_1^{a_1} x_2^{a_2} \dots x_n^{a_n} \geq \sum_{sym} x_1^{b_1} x_2^{b_2} \dots x_n^{b_n}$$

Here, the sums are taken over all permutations of the  $n$  variables.

---

<sup>1</sup>A sequence  $(x_1 \geq x_2 \geq \dots \geq x_n)$  majorizes  $(y_1 \geq y_2 \geq \dots \geq y_n)$  if  $\sum x_i = \sum y_i$  and for all  $k = 1, 2, \dots, n-1$

$$(x_1 + x_2 + \dots + x_k) \geq (y_1 + y_2 + \dots + y_k)$$