Problem L

Let XI be a Time Series such that

Xt = a. Xty +Et, Et NON(0,1), lale 1.

For MLE we need the L1:

 $L(x_{1/x_{2/2}}, x_{1/2}, x_$ 

 $f_{X_{i}}(x_{i}) = \int_{x_{i}=2}^{+} f(x_{i}|x_{i-1},...,x_{i}) \cdot f_{X_{i}}(x_{i})$ 

 $X_{14} \mid X_{16}, \dots X_{l} \sim N(aX_{16}, 1)$ 

The real of

 $X_{+}(X_{+-1}, -- X_{1}) = (X_{+-1}, -- X_{1})$ 

Mear of

 $X_{+}$  is E[X+]= Fa. X++1+ 0

= aX+-1+E+

 $\begin{cases}
X_{17} \mid X_{16}, \dots, X_{1} \\
X_{17} \mid X_{16}, X_{15}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{1}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$   $\begin{cases}
Colon, 2, 2 \\
X_{17} \mid X_{16}, \dots, X_{16}
\end{cases}$ 

( Polizatin little X of R.Y X17 X16, ... X1  $AP(2): X_{t} = \Phi_{1} X_{t-1} + \Phi_{2} X_{t-2} + \varepsilon_{1} \qquad \varepsilon_{t} N(o_{1})$   $X_{t} | X_{t-1} X_{t-2} \sim N(\Phi_{1} X_{t-1} + \Phi_{2} X_{2-2}, 1)$   $f_{X_{t} | X_{t-1} \sim X_{t}} = f_{N(o_{1})} (X_{t-1} - \Phi_{1} X_{t-1} - \Phi_{2} X_{2-1})$  detsify

Box - Cox Cor be Implemented

## Problem 3

AIC = Akoike Information Criterion
-2 lop LL +2k
10# of paramaters
Likelyhood

Fit for differ Models and find lowest AIC

There Are three weys to test best models

- 1-) Info Crievia
- 2-) Pestodoprom and ACF PACF & lostering
- 3-) Filting different models and doing statistical fests to parameters if they are significantly different than "0"

7 incressing

coefficient

course overfitting

this formula

gives the

best coefficient

to add by

perelizing puting

More porouses

and Minputering

Log Richthood

Difference by Sarina and undelling seasonably with during variousless

Sorina undels

Seasonal port os

stochastre varioble.