



1. Heat flows from system A to system B.

- 3 (a) ____ During the heat flow, the entropy of A
A) increases **B)** stays the same **C)** decreases




- 3 (b) If U_A and U_B are the energies of the two systems, what is $\frac{dU_A}{dU_B}$?


- 3 2. ____ The chemical potential of sodium inside a cell is $\mu_{\text{Na}} = -30 \text{ eV}$; outside the cell, $\mu_{\text{Na}} = -28 \text{ eV}$. For the two sides to reach equilibrium, sodium would have to flow
A) into the cell **B)** out of the cell



- 3 3. ____ System A has a temperature of $T = -300 \text{ K}$, and system B has a temperature of $T = 300 \text{ K}$. If the systems are placed in contact,
A) heat will flow from A to B
B) heat will flow from B to A
C) they will undergo matter-antimatter annihilation
D) none of the above



- 2 4. ____ A paramagnet in a magnetic field has an entropy $S(U)$ as shown. It has negative temperature when its energy is
A) low **B)** high **C)** never



$S(U)$ for a paramagnet


- 3 5. ____ A miserly system at $T = 300\text{ K}$ is placed in contact with a normal system at $T = 280\text{ K}$. What happens initially?
A) Both systems get hotter **B)** Both systems get colder
C) M gets colder, N gets hotter **D)** None of these.

M
300K

N
280K

6. The figure shows a process of an ideal gas with $f = 3$, on a PV diagram. The internal energy decreases by -120 kJ during the process.


- 3 (a) ____ How much work is done during this process?
 (Positive means work flows into the gas.)
A) $+10\text{ kJ}$ **B)** $+20\text{ kJ}$ **C)** $+80\text{ kJ}$
D) -10 kJ **E)** -20 kJ **F)** -80 kJ



- 3 (b) ____ *How much* heat Q flows into or out of the gas?
A) 0 kJ
B) -80 kJ **C)** -120 kJ **D)** -200 kJ
E) $+80\text{ kJ}$ **F)** $+120\text{ kJ}$ **G)** $+200\text{ kJ}$


- 3 7. _____ Which is faster?
A) an adiabatic process B) an isothermal process

- 3 8. _____ Suppose gas is in a container sealed by an ideal piston that is free to move. The gas is heated, and the gas expands. This is an ... expansion.
A) adiabatic B) isobaric C) isothermal



9. This figure shows a heat engine operating between 200 K and 800 K.

- 3 (a) _____ What is this engine's efficiency?
A) 25% B) 33% C) 67% D) 75%




- 3 (b) How much work would this engine produce for the same amount of heat $Q_{in} = 45 \text{ J}$, if it were a Carnot engine?

- 3 10. ____ Nitrogen gas ($f = 5$) at standard pressure ($P = 1 \text{ atm}$) expands adiabatically to twice its volume. What is the pressure of the gas after the expansion?
A) 0.38 atm B) 0.5 atm C) 1 atm D) 2 atm E) 2.6 atm

11. This figure shows a refrigerator.

- 3 (a) ____ How much heat Q_{out} flows into the hot reservoir?
A) 5 J B) 15 J
C) 20 J D) 35 J



- 3 (b) ____ What is its Coefficient of Performance?
A) 0.33 B) 0.43 C) 0.75
D) 1.33 E) 2 F) 2.33

- 3 12. ____ Which of these is an intensive quantity?
A) internal energy B) pressure C) volume

3 13. ____ A car has two identical hubcaps: one is shiny and new, the other is rusty. Which has the larger Gibbs free energy?

- A) the shiny one B) the rusty one


3 14. ____ How does the van der Waals model differ from the ideal gas model?

- A) it models a phase transition
B) it takes into account the minimum volume of atoms
C) it explains the critical point
D) all of the above

15. During a chemical process at constant temperature and pressure, 30 J of heat flows into the system, the internal energy increases by $\Delta U = +10$ J, and 20 J of compression work flows outward to move the air out of the way.

2 (a) ____ What is the change ΔH in the enthalpy?

- A) -20 J B) -10 J C) 0 J
D) $+10$ J E) $+20$ J F) $+30$ J



2 (b) ____ What is the change ΔG in the Gibbs free energy?

- A) -20 J B) -10 J C) 0 J
D) $+10$ J E) $+20$ J F) $+30$ J

16. We can write the thermodynamic identity of a rubber band as

$$dU = T dS + F dL$$

where L is the length of the rubber band, and F is the tension in the rubber band.

3 (a) What are the natural variables of U *in this case*?

3 (b) What is $\left(\frac{\partial S}{\partial L}\right)_U$?

3 (c) _____ Which of these is a pair of conjugate variables?
A) T & U **B)** S & L **C)** F & L **D)** U & S

3 17. What is $\left(\frac{\partial H}{\partial S}\right)_{P,N}$?

18. In the following equation, fill in the blanks.

$$\left(\frac{\partial \boxed{\text{a}}}{\partial V}\right)_{T,N} = \boxed{\text{c}} \left(\frac{\partial P}{\partial N}\right)_{V,\boxed{\text{b}}}$$

$$\boxed{2} \quad \text{(a)} \quad \begin{array}{cccccc} \text{F)} & F & \mu) & \mu & \text{N)} & N & \text{P)} & P & \text{S)} & S & \text{T)} & T & \text{V)} & V \end{array}$$


2 (b) _____

F) F μ) μ N) N P) P S) S T) T V) V

2 (c) _____

+) + _____ -) - _____

5] 19. This figure is the phase diagram of water. Label the following regions and points on the diagram. (You can use the letter or the word as you like.)
S) solid **L)** liquid **G)** gas **T)** triple point **C)** critical point



20. Consider a new material called *bowlingrene*. At standard atmospheric pressure, the Gibbs free energy (in joules per mole) of the solid and liquid phases of bowlingrene, as functions of temperature, are

$$\text{solid: } G_s(T) = -8000 - 30T \quad \text{liquid: } G_l(T) = +8000 - 80T$$

The entropy of each phase is independent of temperature, and the pressure is held constant.

- 3 (a) Find the entropy (per mole) of solid bowlingrene.

- 3 (b) _____ At 300 K, bowlingrene is
 A) solid **B)** liquid

- 3 (c) Find the melting/freezing temperature of bowlingrene. (*Note: your answer shouldn't contradict your answer to part (b).*)

- 3 XC 21. **Extra Credit:** The figure shows the boundary between the solid and liquid phases of a material. At $P = 1 \text{ atm}$, the solid phase has a density of $\rho = 900 \text{ kg/m}^3$, the liquid phase has a density of $\rho = 800 \text{ kg/m}^3$, and the latent heat of fusion is $L = 4000 \text{ J}$. Find the slope of the boundary (with the correct units).

