STATISTICAL ANALYSIS PLAN

A Phase 1 Open-label, Dose-escalation Study of Kelvaparib in Adults with Advanced Solid Tumours

Study KLV-7702

Indication: Advanced solid tumours with a homologous recombination repair defect

Sponsor	Marnholt Oncology GmbH
Study number	KLV-7702
Phase	Phase 1
Plan version	Version 2.1
Plan date	1 April 2019
Author	Biostatistics, Marnholt
Status	Final, approved before database lock

REVISION HISTORY

This plan has been reissued. Each version supersedes the one before it in full; the analyses described here are those in force at database lock.

Version	Date	Principal change
1.0	31 January 2019	Original approved version.
2.1	1 April 2019	Censoring conventions and the sensitivity analyses aligned with regulatory guidance.

CHANGES IN VERSION 2.1

Version 2.1 revises version 1.0. It restates the censoring conventions and the sensitivity analyses, the principal change recorded in the revision history above; no endpoint definition or analysis set is changed by it. The version in force at database lock is the one recorded on the cover of this document.

1. INTRODUCTION

This plan describes the analyses to be performed for study KLV-7702. It expands the statistical section of the protocol and takes precedence where the two differ. It was finalised and approved before the database was locked on 1 May 2019 and before any unblinded analysis.

2. OBJECTIVES AND ENDPOINTS

The primary objective is maximum tolerated dose and recommended phase 2 dose. The primary endpoint is incidence of dose-limiting toxicity during the first 21 days at the assigned cohort dose, and the maximum tolerated dose derived from it.

Secondary endpoints are duration of response, progression-free survival, overall survival and the incidence and severity of treatment-emergent adverse events. Progression-free survival is defined as the time from the first dose to the first documented disease progression or death from any cause, whichever occurs first. Overall survival is defined as the time from the first dose to death from any cause. Duration of response is defined, for a subject with a confirmed complete or partial response, as the time from the first dose of study treatment to the end of response recorded for the subject in the analysis dataset.

3. STUDY DESIGN

An open-label, multicentre, non-randomised, sequential dose-escalation study using a 3+3 design, followed by expansion at the dose selected as the recommended phase 2 dose. Approximately 58 subjects across 4 dose cohorts.

4. ANALYSIS POPULATIONS

Population	Definition	Used for
Safety analysis set	All subjects who received at least one dose of study treatment.	All safety analyses and exposure.
Full analysis set	All subjects in the safety analysis set with at least one post-baseline disease assessment.	All efficacy analyses.
Per-protocol set	Full analysis set subjects without a protocol deviation leading to discontinuation of study treatment.	Sensitivity analysis of the primary endpoint.
Pharmacokinetic analysis set	Subjects with at least one evaluable post-dose concentration.	Pharmacokinetic summaries.
DLT-evaluable set	Subjects who received at least 21 days of study treatment or had a dose-limiting toxicity.	Determination of the maximum tolerated dose.

5. GENERAL CONSIDERATIONS

Continuous variables are summarised by n, mean, standard deviation, median, first and third quartiles, minimum and maximum. Categorical variables by frequency and percentage with the denominator stated. Percentages are rounded to one decimal place and suppressed where the denominator is fewer than five. Analyses are performed using SAS version 9.4 or later.

Baseline is the last non-missing assessment on or before the first dose. A treatment-emergent adverse event is any event with onset on or after the first dose, or a pre-existing event that worsens after the first dose.

6. DISPOSITION AND BASELINE CHARACTERISTICS

Disposition is summarised by dose cohort, giving the numbers screened, treated, ongoing and discontinued with reasons. Demographic and baseline disease characteristics are summarised descriptively; no formal comparison is performed.

7. EFFICACY ANALYSES

7.1 Primary endpoint

The number and proportion experiencing a dose-limiting toxicity is summarised by dose cohort in the DLT-evaluable set. The maximum tolerated dose is the highest dose level at which fewer than 33 per cent of DLT-evaluable subjects had a dose-limiting toxicity; the expansion subjects treated at a dose level are included. No hypothesis is tested.

7.2 Secondary endpoints

Duration of response, Progression-free survival, Overall survival are estimated by the Kaplan-Meier method with medians and 95 per cent confidence limits by the Brookmeyer-Crowley method. Binomial proportions are summarised with exact 95 per cent confidence intervals.

7.3 Multiplicity

No multiplicity adjustment is required because only one primary comparison is performed and all secondary analyses are descriptive. Nominal p-values, where presented, are not adjusted and are not used to claim significance.

7.4 Subgroup analyses

The primary endpoint is summarised within subgroups defined by BRCA mutation status, homologous recombination status, prior lines of therapy, age category, sex, ECOG performance status. Subgroup analyses are exploratory, are not adjusted for multiplicity, and no significance is claimed from them.

7.5 Sensitivity analyses

The primary analysis is repeated in the per-protocol set.

8. SAFETY ANALYSES

Treatment-emergent adverse events are summarised by system organ class and preferred term using MedDRA version 22.0, with subjects counted once per term at their highest grade. Separate summaries cover events of grade 3 or higher, serious events, related events, events leading to dose modification or discontinuation, and fatal events. Adverse events of special interest (myelodysplastic syndrome or acute myeloid leukaemia, pneumonitis, severe anaemia) are summarised individually with time to onset and resolution.

Laboratory data are summarised by shift from baseline to worst post-baseline CTCAE grade. Vital signs are summarised by change from baseline.

9. PHARMACOKINETIC ANALYSES

Concentrations are summarised by nominal sampling time using n, arithmetic mean, standard deviation, geometric mean, geometric coefficient of variation, median, minimum and maximum. Derived parameters are summarised by dose. Concentrations below the limit of quantification are set to zero for descriptive summaries and excluded from geometric means.

10. INTERIM ANALYSES

No interim analysis is performed. Dose-escalation decisions are taken cohort by cohort under the rules of the protocol and are not interim analyses of the primary endpoint.

11. CHANGES FROM THE PROTOCOL-SPECIFIED ANALYSES

This plan clarified which categories of protocol deviation are considered important for the per-protocol set and extended the subgroup list. Neither change affects the primary analysis. No change was made after database lock.

12. TABLE, LISTING AND FIGURE SPECIFICATIONS

The index of outputs and their shells is provided in the appendix. Numbering follows ICH E3: Section 14.1 demographic and disposition, 14.2 efficacy (including the subgroup analyses of the primary endpoint), 14.3 safety, 14.4 pharmacokinetics.

13 STUDY OBJECTIVES AND ENDPOINTS

The objectives and endpoints analysed under this plan are those in the protocol. Where this plan gives a more precise definition of an endpoint than the protocol, this plan governs the analysis and the difference is recorded in Section 19.

14 GENERAL ANALYSIS CONVENTIONS

14.1 Definition of baseline

Baseline is the last non-missing assessment obtained on or before the date and time of the first dose of study treatment. Where an assessment on the day of first dose has no time recorded it is taken as pre-dose unless another record shows otherwise.

14.2 Definition of study day

Study day is calculated relative to the date of first dose, which is day 1. There is no day 0; the day before first dose is day -1. This convention is applied in every dataset and every output.

14.3 Treatment-emergent period

An event is treatment-emergent if its onset is on or after the first dose, or if it was present before the first dose and worsened after it. Adverse events are collected to the end of study participation (protocol Section 7.3).

14.4 Handling of partial and missing dates

A partial date is completed conservatively: a missing day is set to the first of the month for a start date and the last of the month for an end date; a missing month is set to January for a start date and December for an end date. Where completion would place a start date before the first dose, the event is treated as treatment-emergent unless other information shows otherwise.

14.5 Presentation conventions

Continuous variables are summarised by the number of observations, mean, standard deviation, median, minimum and maximum. Percentages are presented to one decimal place and calculated on the number of subjects in the analysis set unless a different denominator is stated in the table. Confidence intervals are two-sided at the 95 per cent level unless stated otherwise.

15 ANALYSIS SETS

Table 1 Analysis sets

Set	Definition	Used for
Enrolled	All subjects assigned to treatment.	Disposition and deviations.
Safety	All subjects who received at least one dose, analysed as treated.	All safety analyses.
DLT evaluable	Subjects who received at least 21 days of study treatment or had a dose-limiting toxicity.	Primary endpoint.
Full analysis	All subjects in the safety analysis set with at least one post-baseline disease assessment, analysed as assigned.	Primary and secondary efficacy.
Per protocol	Full analysis set subjects without a protocol deviation leading to discontinuation of study treatment.	Sensitivity analysis of the primary endpoint.
Efficacy evaluable	The same subjects as the full analysis set.	Response analyses.
Pharmacokinetic	Subjects with at least one measurable post-dose concentration and no deviation affecting it.	Pharmacokinetic summaries.

16 ESTIMANDS

The estimand for the primary endpoint is defined by its population, variable, the handling of intercurrent events and the population-level summary. The population is the full analysis set. The variable is the primary endpoint as defined in Section 13. Intercurrent events are handled as follows: discontinuation of study treatment for any reason is handled by the treatment policy strategy, so data after discontinuation continue to contribute; no data on subsequent anti-disease therapy were collected, so its start is not used to censor; death without the event of interest is handled as described in Section 17.2. The summary measure is the proportion of subjects with the event, with its exact confidence interval.

17 STATISTICAL METHODS

17.1 Primary analysis

Incidence of dose-limiting toxicity during the first 21 days at the assigned cohort dose, and the maximum tolerated dose derived from it.

No formal hypothesis testing is planned. Sample size follows the operating characteristics of the 3+3 design and the number of subjects required to characterise safety at the selected dose. Proportions are presented with exact Clopper-Pearson 95 per cent confidence intervals and time-to-event endpoints estimated by the Kaplan-Meier method.

No data on subsequent anti-disease therapy were collected. A subject who starts a subsequent therapy is therefore not censored at its start: follow-up continues regardless of it (the treatment policy strategy for that intercurrent event).

17.2 Censoring rules

Progression-free survival: a subject with neither documented progression nor death is censored at the date of the last assessment at which absence of progression was documented or, if still in follow-up at the data cut-off, at the data cut-off date; a subject with no post-baseline assessment is censored at day 1; death without documented progression is an event at the date of death. Overall survival: a subject not known to have died is censored at the date last known alive or, if still in follow-up at the data cut-off, at the data cut-off date. Duration of response: measured from the first dose of study treatment to the recorded end of response; the response and progression dates are as held in the analysis datasets, which hold no assessment-level tumour data from which they could be re-derived; a responder whose response has not ended is censored at the date of the last assessment or, if still in follow-up at the data cut-off, at the data cut-off date. No subject is censored at the start of a subsequent anti-disease therapy.

17.3 Sensitivity and supplementary analyses

- Repeat of the primary analysis on the per-protocol set.
- Repeat of the primary analysis treating subjects with no post-baseline assessment as non-responders.

17.4 Subgroup analyses

The primary endpoint is examined in the prespecified subgroups listed in the table shells. These analyses are descriptive, are not adjusted for multiplicity, and are not used to claim an effect in any subgroup.

17.5 Multiplicity

No multiplicity adjustment is required because only one primary comparison is performed and all secondary analyses are descriptive. Nominal p-values, where presented, are not adjusted and are not used to claim significance.

17.6 Interim analyses

Any interim analysis is described here before it is performed, including its timing, the quantities to be examined, the alpha spent if any, and the actions available to the sponsor. No interim analysis is performed (Section 10).

17.7 Safety analyses

Safety analyses are descriptive and are performed on the safety set. Adverse events are coded with the Medical Dictionary for Regulatory Activities and graded with the Common Terminology Criteria for Adverse Events. A subject with more than one occurrence of the same preferred term is counted once at the highest grade. Laboratory results are graded and presented as shifts from baseline to the worst post-baseline grade.

17.8 Pharmacokinetic analyses

Concentrations below the lower limit of quantification are set to zero before the first measurable concentration and treated as missing thereafter. Parameters are summarised by geometric mean and geometric coefficient of variation. No formal comparison between doses is performed unless stated in the protocol.

18 TABLE, LISTING AND FIGURE INVENTORY

The outputs below are produced under this plan. Each is specified by its population, its content and its sort order. Shells are held in the output specification document.

Table 2 Planned tables and figures

Number	Title	Population	Content
14.1.1	Demographic and Baseline Characteristics	Safety	Descriptive statistics by treatment group and overall.
14.1.2	Subject Disposition	Safety	Number and percentage by disposition category and treatment group.
14.1.3	Extent of Exposure	Safety	Duration of treatment, cycles started, relative dose intensity, dose reductions and interruptions.
14.2.1	Best Overall Response by RECIST 1.1	Full analysis	Best overall response by RECIST 1.1 and objective response rate with its exact 95 per cent confidence interval.
14.2.2	Kaplan-Meier Estimates of Time-to-event Endpoints	Full analysis	Kaplan-Meier median and quartiles of duration of response, progression-free survival and overall survival.
14.2.3	Overall Response Rate and Progression-free Survival by BRCA Mutation Status	Full analysis	Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.
14.2.4	Overall Response Rate and Progression-free Survival by Homologous Recombination Status	Full analysis	Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.
14.2.5	Overall Response Rate and Progression-free Survival by Prior Lines of Therapy	Full analysis	Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Number	Title	Population	Content
14.2.6	Overall Response Rate and Progression-free Survival by Age Category	Full analysis	Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.
14.2.7	Overall Response Rate and Progression-free Survival by Sex	Full analysis	Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.
14.2.8	Overall Response Rate and Progression-free Survival by ECOG Performance Status	Full analysis	Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.
14.3.1	Overview of Treatment-emergent Adverse Events	Safety	Number and percentage with any event, grade 3 or higher, serious, fatal, leading to dose modification or discontinuation.
14.3.2	Treatment-emergent Adverse Events Occurring in at Least 2% of Subjects by System Organ Class and Preferred Term	Safety	Number and percentage by system organ class and preferred term, sorted by descending frequency.
14.3.3	Treatment-emergent Adverse Events of Grade 3 or Higher by System Organ Class and Preferred Term	Safety	As the table by system organ class and preferred term, restricted to grade 3 or higher.
14.3.4	Treatment-related Treatment-emergent Adverse Events Occurring in at Least 2% of Subjects	Safety	Number and percentage of events assessed as related, by preferred term.
14.3.5	Serious Treatment-emergent Adverse Events by Preferred Term	Safety	Number and percentage by preferred term.

Number	Title	Population	Content
14.3.6	Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment	Safety	Number and percentage by preferred term.
14.3.7	Deaths	Safety	Number and percentage by primary cause.
14.3.8	Shift from Baseline to Worst Post-baseline CTCAE Grade for Selected Laboratory Parameters	Safety	Worst post-baseline CTCAE grade and shifts of two or more grades from baseline, by parameter.
14.3.9	Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events	Safety	Median time to onset and to resolution of the most frequent events.
14.3.10	Overview of adverse events by BRCA mutation status	Safety	Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.
14.3.11	Overview of adverse events by homologous recombination status	Safety	Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.
14.3.12	Overview of adverse events by prior lines of therapy	Safety	Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.
14.3.13	Overview of adverse events by age category	Safety	Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.
14.3.14	Overview of adverse events by sex	Safety	Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Number	Title	Population	Content
14.3.15	Overview of adverse events by ECOG performance status	Safety	Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.
14.3.16	Exposure-Adjusted Adverse Events	Safety	Exposure in patient-years and exposure-adjusted event rates, overall and by subgroup.
14.4.1	Summary of Steady-state Pharmacokinetic Parameters by Dose Cohort	Pharmacokinetic	Summary statistics of steady-state pharmacokinetic parameters by dose.
14.4.2	Accumulation Ratio and Trough Concentration by Dose Cohort	Pharmacokinetic	Accumulation ratio and trough concentration by dose.

18.1 Shells

A shell is given for every output. The shell fixes the title, the column structure, the denominators, the sort order and the footnotes, so that the output can be programmed and reviewed before any data are seen.

Table 14.1.1 Demographic and Baseline Characteristics

Population: Safety analysis set

Content: Descriptive statistics by treatment group and overall.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.1.2 Subject Disposition

Population: Safety analysis set

Content: Number and percentage by disposition category and treatment group.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.1.3 Extent of Exposure

Population: Safety analysis set

Content: Duration of treatment, cycles started, relative dose intensity, dose reductions and interruptions.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.1 Best Overall Response by RECIST 1.1

Population: Full analysis analysis set

Content: Best overall response by RECIST 1.1 and objective response rate with its exact 95 per cent confidence interval.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.2 Kaplan-Meier Estimates of Time-to-event Endpoints

Population: Full analysis analysis set

Content: Kaplan-Meier median and quartiles of duration of response, progression-free survival and overall survival.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.3 Overall Response Rate and Progression-free Survival by BRCA Mutation Status

Population: Full analysis analysis set

Content: Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Subgroup	n	Responders, n (%)	95% CI	Median PFS (months)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.4 Overall Response Rate and Progression-free Survival by Homologous Recombination Status

Population: Full analysis analysis set

Content: Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Subgroup	n	Responders, n (%)	95% CI	Median PFS (months)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.5 Overall Response Rate and Progression-free Survival by Prior Lines of Therapy

Population: Full analysis analysis set

Content: Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Subgroup	n	Responders, n (%)	95% CI	Median PFS (months)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.6 Overall Response Rate and Progression-free Survival by Age Category

Population: Full analysis analysis set

Content: Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Subgroup	n	Responders, n (%)	95% CI	Median PFS (months)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.7 Overall Response Rate and Progression-free Survival by Sex

Population: Full analysis analysis set

Content: Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Subgroup	n	Responders, n (%)	95% CI	Median PFS (months)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.2.8 Overall Response Rate and Progression-free Survival by ECOG Performance Status

Population: Full analysis analysis set

Content: Objective response rate with its exact 95 per cent confidence interval and median progression-free survival in each subgroup level.

Subgroup	n	Responders, n (%)	95% CI	Median PFS (months)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.1 Overview of Treatment-emergent Adverse Events

Population: Safety analysis set

Content: Number and percentage with any event, grade 3 or higher, serious, fatal, leading to dose modification or discontinuation.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.2 Treatment-emergent Adverse Events Occurring in at Least 2% of Subjects by System Organ Class and Preferred Term

Population: Safety analysis set

Content: Number and percentage by system organ class and preferred term, sorted by descending frequency.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.3 Treatment-emergent Adverse Events of Grade 3 or Higher by System Organ Class and Preferred Term

Population: Safety analysis set

Content: As the table by system organ class and preferred term, restricted to grade 3 or higher.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.4 Treatment-related Treatment-emergent Adverse Events Occurring in at Least 2% of Subjects

Population: Safety analysis set

Content: Number and percentage of events assessed as related, by preferred term.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.5 Serious Treatment-emergent Adverse Events by Preferred Term

Population: Safety analysis set

Content: Number and percentage by preferred term.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.6 Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment

Population: Safety analysis set

Content: Number and percentage by preferred term.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.7 Deaths

Population: Safety analysis set

Content: Number and percentage by primary cause.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.8 Shift from Baseline to Worst Post-baseline CTCAE Grade for Selected Laboratory Parameters

Population: Safety analysis set

Content: Worst post-baseline CTCAE grade and shifts of two or more grades from baseline, by parameter.

Baseline grade	Post-baseline grade 0 to 1	Grade 2	Grade 3 or higher
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.9 Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events

Population: Safety analysis set

Content: Median time to onset and to resolution of the most frequent events.

Preferred term	Events, n	Median time to onset (days)	Resolved, n (%)	Median time to resolution (days)
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.10 Overview of adverse events by BRCA mutation status

Population: Safety analysis set

Content: Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Category	Subgroup level 1	Subgroup level 2	Subgroup level 3
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.11 Overview of adverse events by homologous recombination status

Population: Safety analysis set

Content: Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Category	Subgroup level 1	Subgroup level 2	Subgroup level 3
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.12 Overview of adverse events by prior lines of therapy

Population: Safety analysis set

Content: Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Category	Subgroup level 1	Subgroup level 2	Subgroup level 3
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.13 Overview of adverse events by age category

Population: Safety analysis set

Content: Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Category	Subgroup level 1	Subgroup level 2	Subgroup level 3
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.14 Overview of adverse events by sex

Population: Safety analysis set

Content: Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Category	Subgroup level 1	Subgroup level 2	Subgroup level 3
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.15 Overview of adverse events by ECOG performance status

Population: Safety analysis set

Content: Subjects with any event, grade 3 or higher, serious, leading to discontinuation and fatal, in each subgroup level.

Category	Subgroup level 1	Subgroup level 2	Subgroup level 3
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.3.16 Exposure-Adjusted Adverse Events

Population: Safety analysis set

Content: Exposure in patient-years and exposure-adjusted event rates, overall and by subgroup.

Category	100 mg	200 mg	300 mg	400 mg	Total
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.4.1 Summary of Steady-state Pharmacokinetic Parameters by Dose Cohort

Population: Pharmacokinetic analysis set

Content: Summary statistics of steady-state pharmacokinetic parameters by dose.

Parameter	n	Geometric mean (CV%)	Range
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

Table 14.4.2 Accumulation Ratio and Trough Concentration by Dose Cohort

Population: Pharmacokinetic analysis set

Content: Accumulation ratio and trough concentration by dose.

Parameter	n	Geometric mean (CV%)	Range
Row label 1	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 2	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 3	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 4	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 5	xx (xx.x)	xx (xx.x)	xx (xx.x)
Row label 6	xx (xx.x)	xx (xx.x)	xx (xx.x)

xx denotes a count and xx.x a percentage, to be replaced at production. Percentages are calculated on the number of subjects in the analysis set unless stated otherwise. Sort order: descending frequency in the total column, then alphabetically.

19 CHANGES FROM THE PROTOCOL

Where this plan differs from the analysis described in the protocol, the difference and the reason for it are recorded here. Any change made after database lock is reported in the clinical study report and identified as post hoc.

20 PROGRAMMING CONSIDERATIONS

Analyses are produced from the analysis datasets described in the dataset specification, which follow the Analysis Data Model. Outputs are produced by validated programs, and every output is independently reproduced from the same analysis datasets by a second programmer before it is released.