Study KLV-7701 — Tables, Listings and Figures

Kelvaparib (KLV-3308) — Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

Table 14.3.9 Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events – Safety Analysis Set

Preferred term	Events, n	Median time to onset (days)	Resolved, n (%)	Median time to resolution (days)
Nausea	393	32.0	330 (84.0)	11.0
Anaemia	317	29.0	265 (83.6)	10.0
Fatigue	299	32.0	239 (79.9)	10.0
Vomiting	163	34.0	130 (79.8)	11.0
Constipation	163	29.0	131 (80.4)	9.0
Decreased appetite	152	38.0	127 (83.6)	10.0
Thrombocytopenia	152	41.0	129 (84.9)	12.0
Neutropenia	127	36.0	98 (77.2)	9.5
Headache	127	25.0	105 (82.7)	11.0
Diarrhoea	119	37.0	97 (81.5)	11.0
Abdominal pain	118	25.5	102 (86.4)	11.5
Dysgeusia	115	24.0	92 (80.0)	10.0
Dizziness	98	23.5	76 (77.6)	12.0
Asthenia	87	33.0	74 (85.1)	10.0
Cough	81	33.0	65 (80.2)	10.0
Back pain	80	28.0	67 (83.8)	10.0
Nasopharyngitis	77	29.0	61 (79.2)	11.0
Insomnia	76	38.0	65 (85.5)	9.0
Arthralgia	74	36.0	65 (87.8)	8.0
Leukopenia	73	30.0	57 (78.1)	11.0

Events are counted at the event level rather than the subject level. NE = not estimable.