Study KLV-7701 — Tables, Listings and Figures

Kelvaparib (KLV-3308) — Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

Table 14.3.6 Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Anaemia	11 (2.6)	8 (3.8)	19 (3.0)
Malignant neoplasm progression	6 (1.4)	7 (3.3)	13 (2.1)
Sepsis	8 (1.9)	3 (1.4)	11 (1.8)
Neutropenia	5 (1.2)	3 (1.4)	8 (1.3)
Pneumonia	7 (1.7)	1 (0.5)	8 (1.3)
Thrombocytopenia	7 (1.7)	0 (0.0)	7 (1.1)
Intestinal obstruction	2 (0.5)	2 (1.0)	4 (0.6)
Ascites	2 (0.5)	1 (0.5)	3 (0.5)
Dyspnoea	3 (0.7)	0 (0.0)	3 (0.5)
Leukopenia	1 (0.2)	2 (1.0)	3 (0.5)
Myelodysplastic syndrome	1 (0.2)	2 (1.0)	3 (0.5)
Pneumonitis	1 (0.2)	2 (1.0)	3 (0.5)
Fatigue	2 (0.5)	0 (0.0)	2 (0.3)
Lymphopenia	2 (0.5)	0 (0.0)	2 (0.3)
Pulmonary embolism	1 (0.2)	1 (0.5)	2 (0.3)
Abdominal pain	1 (0.2)	0 (0.0)	1 (0.2)
Asthenia	1 (0.2)	0 (0.0)	1 (0.2)
Constipation	0 (0.0)	1 (0.5)	1 (0.2)
Decreased appetite	1 (0.2)	0 (0.0)	1 (0.2)
Dizziness	0 (0.0)	1 (0.5)	1 (0.2)
Nausea	1 (0.2)	0 (0.0)	1 (0.2)
Pleural effusion	0 (0.0)	1 (0.5)	1 (0.2)
Pyrexia	1 (0.2)	0 (0.0)	1 (0.2)

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)

Percentages are based on N = 627.