Study KLV-7701 — Tables, Listings and Figures

Kelvaparib (KLV-3308) — Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

Table 14.1.1 Demographic and Baseline Characteristics – Safety Analysis Set

Characteristic	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Age (years)			
Mean (SD)	65.3 (9.00)	64.7 (9.21)	65.1 (9.07)
Median	65.0	65.0	65.0
Min, max	38, 88	42, 88	38, 88
>= 65 years, n (%)	225 (53.8)	108 (51.7)	333 (53.1)
Sex, n (%)			
Male	0 (0.0)	0 (0.0)	0 (0.0)
Female	418 (100.0)	209 (100.0)	627 (100.0)
Race, n (%)			
White	342 (81.8)	161 (77.0)	503 (80.2)
Asian	38 (9.1)	25 (12.0)	63 (10.0)
Black or African American	20 (4.8)	11 (5.3)	31 (4.9)
Other	18 (4.3)	12 (5.7)	30 (4.8)
ECOG performance status, n (%)			
0	193 (46.2)	103 (49.3)	296 (47.2)
1	184 (44.0)	89 (42.6)	273 (43.5)
2	41 (9.8)	17 (8.1)	58 (9.3)
BRCA mutation status, n (%)			
BRCA1 mutated	101 (24.2)	49 (23.4)	150 (23.9)
BRCA2 mutated	62 (14.8)	33 (15.8)	95 (15.2)
BRCA wild-type	255 (61.0)	127 (60.8)	382 (60.9)
Homologous recombination deficiency, n (%)	222 (53.1)	104 (49.8)	326 (52.0)
Platinum-sensitive disease, n (%)	265 (63.4)	135 (64.6)	400 (63.8)
Residual disease after cytoreductive surgery, n (%)	178 (42.6)	88 (42.1)	266 (42.4)
Number of prior lines of platinum-based therapy			
Median	2.0	2.0	2.0
>= 3, n (%)	122 (29.2)	61 (29.2)	183 (29.2)

SD = standard deviation; ECOG = Eastern Cooperative Oncology Group. Percentages are based on the number of subjects in the safety analysis set within each group.