TABLES, LISTINGS AND FIGURES

Study KLV-7703 — A Phase 3 Randomised, Placebo-controlled Study of Kelvaparib in Recurrent High-grade Serous Ovarian Carcinoma

Kelvaparib (KLV-3308)

Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

This pack contains the post-text tables, listings and figures supporting the clinical study report for study KLV-7703. All displays derive from the analysis datasets locked on 7 February 2023.

Contents

- Table 14.1.1 Demographic and Baseline Characteristics
- Table 14.1.2 Subject Disposition
- Table 14.1.3 Extent of Exposure
- Table 14.2.1 Best Overall Response
- Table 14.2.2 Time-to-Event Analyses
- Table 14.3.1 Overview of Adverse Events
- Table 14.3.2a Adverse Events by System Organ Class and Preferred Term
- Table 14.3.5 Serious Adverse Events
- Table 14.3.4a Treatment-Related Adverse Events
- Table 14.3.6 Adverse Events Leading to Discontinuation
- Table 14.3.7 Deaths
- Table 14.3.8 Laboratory Shift from Baseline to Worst Grade
- Table 14.3.9 Adverse Event Onset Timing

Table 14.1.1 Demographic and Baseline Characteristics

Table 14.1.1 Demographic and Baseline Characteristics – Safety Analysis Set

Characteristic	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Age (years)			
Mean (SD)	65.0 (8.44)	66.4 (8.93)	65.5 (8.62)
Median	65.0	66.5	65.0
Min, max	44, 88	45, 88	44, 88
>= 65 years, n (%)	125 (53.0)	69 (58.5)	194 (54.8)
Sex, n (%)			
Male	0 (0.0)	0 (0.0)	0 (0.0)
Female	236 (100.0)	118 (100.0)	354 (100.0)
Race, n (%)			
White	192 (81.4)	98 (83.1)	290 (81.9)
Asian	19 (8.1)	6 (5.1)	25 (7.1)
Black or African American	16 (6.8)	6 (5.1)	22 (6.2)
Other	9 (3.8)	8 (6.8)	17 (4.8)
ECOG performance status, n (%)			
0	98 (41.5)	50 (42.4)	148 (41.8)
1	118 (50.0)	53 (44.9)	171 (48.3)
2	20 (8.5)	15 (12.7)	35 (9.9)
BRCA mutation status, n (%)			
BRCA1 mutated	55 (23.3)	29 (24.6)	84 (23.7)
BRCA2 mutated	44 (18.6)	14 (11.9)	58 (16.4)
BRCA wild-type	137 (58.1)	75 (63.6)	212 (59.9)
Homologous recombination deficiency, n (%)	122 (51.7)	63 (53.4)	185 (52.3)
Platinum-sensitive disease, n (%)	155 (65.7)	82 (69.5)	237 (66.9)
Residual disease after cytoreductive surgery, n (%)	101 (42.8)	46 (39.0)	147 (41.5)
Number of prior lines of platinum-based therapy			
Median	2.0	2.0	2.0

Characteristic	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
>= 3, n (%)	64 (27.1)	33 (28.0)	97 (27.4)

SD = standard deviation; ECOG = Eastern Cooperative Oncology Group. Percentages are based on the number of subjects in the safety analysis set within each group.

Table 14.1.2 Subject Disposition

Table 14.1.2 Subject Disposition – Safety Analysis Set

Disposition	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Subjects treated, n	236	118	354
Treatment ongoing at data cut-off, n (%)	61 (25.8)	21 (17.8)	82 (23.2)
Discontinued study treatment, n (%)	175 (74.2)	97 (82.2)	272 (76.8)
Disease progression	40 (16.9)	26 (22.0)	66 (18.6)
Adverse event	11 (4.7)	6 (5.1)	17 (4.8)
Death	20 (8.5)	8 (6.8)	28 (7.9)
Withdrawal by subject	37 (15.7)	24 (20.3)	61 (17.2)
Physician decision	35 (14.8)	19 (16.1)	54 (15.3)
Lost to follow-up	17 (7.2)	6 (5.1)	23 (6.5)
Protocol deviation	15 (6.4)	8 (6.8)	23 (6.5)

Percentages are based on the number of subjects treated within each group.

Table 14.1.3 Extent of Exposure

Table 14.1.3 Extent of Exposure – Safety Analysis Set

Exposure parameter	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Duration of treatment (days)			
Median	134.0	127.5	131.5
Q1, Q3	82.0, 181.8	73.8, 165.5	80.0, 179.0
Min, max	8, 307	3, 280	3, 307
Cycles started, median	5.0	4.5	5.0
Relative dose intensity (%), median	85.7	85.9	85.8
Any dose reduction, n (%)	42 (17.8)	23 (19.5)	65 (18.4)
Any dose interruption, n (%)	51 (21.6)	32 (27.1)	83 (23.4)

Q1 = first quartile; Q3 = third quartile. Relative dose intensity is the total dose received divided by the total dose planned over the treatment period.

Table 14.2.1 Best Overall Response

Table 14.2.1 Best Overall Response by RECIST 1.1 – Full Analysis Set

Best overall response	Kelvaparib 300 mg (N=227)	Placebo (N=109)	Total (N=336)
Complete response (CR)	28 (12.3)	2 (1.8)	30 (8.9)
Partial response (PR)	101 (44.5)	26 (23.9)	127 (37.8)
Stable disease	57 (25.1)	52 (47.7)	109 (32.4)
Progressive disease	41 (18.1)	29 (26.6)	70 (20.8)
Not evaluable	0 (0.0)	0 (0.0)	0 (0.0)
Objective response rate, n (%)	129 (56.8)	28 (25.7)	157 (46.7)
95% CI	(50.1, 63.4)	(17.8, 34.9)	(41.3, 52.2)

CI = confidence interval. Assessed according to RECIST version 1.1. Confidence intervals are exact (Clopper-Pearson).

Table 14.2.2 Time-to-Event Analyses

Table 14.2.2 Kaplan-Meier Estimates of Time-to-event Endpoints – Full Analysis Set

Endpoint	Kelvaparib 300 mg (N=227)	Placebo (N=109)	Total (N=336)
Duration of response, median (months)	11.4	9.3	11.4
25th, 75th percentile	(6.1, 15.5)	(4.6, 19.6)	(5.4, 15.5)
Progression-free survival, median (months)	8.6	11.1	8.8
25th, 75th percentile	(2.8, 12.5)	(5.9, 21.2)	(3.7, 14.2)
Overall survival, median (months)	26.0	NE	28.7
25th, 75th percentile	(20.2, NE)	(21.9, NE)	(21.3, NE)

NE = not estimable. Duration of response is analysed among responding subjects only.

Table 14.3.1 Overview of Adverse Events

Table 14.3.1 Overview of Treatment-emergent Adverse Events – Safety Analysis Set

Category, n (%)	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Any TEAE	236 (100.0)	118 (100.0)	354 (100.0)
Any treatment-related TEAE	234 (99.2)	118 (100.0)	352 (99.4)
Any grade >= 3 TEAE	122 (51.7)	57 (48.3)	179 (50.6)
Any serious TEAE	57 (24.2)	26 (22.0)	83 (23.4)
Any TEAE leading to dose reduction	42 (17.8)	22 (18.6)	64 (18.1)
Any TEAE leading to discontinuation	30 (12.7)	13 (11.0)	43 (12.1)
Any fatal TEAE	22 (9.3)	6 (5.1)	28 (7.9)

TEAE = treatment-emergent adverse event. A subject is counted once within each category.

Table 14.3.2a Adverse Events by System Organ Class and Preferred Term

Table 14.3.2a Treatment-emergent Adverse Events Occurring in at Least 10% of Subjects by System Organ Class and Preferred Term – Safety Analysis Set

System organ class / Preferred term	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Gastrointestinal disorders	201 (85.2)	107 (90.7)	308 (87.0)
Nausea	147 (62.3)	76 (64.4)	223 (63.0)
Vomiting	72 (30.5)	42 (35.6)	114 (32.2)
Constipation	66 (28.0)	46 (39.0)	112 (31.6)
Diarrhoea	49 (20.8)	20 (16.9)	69 (19.5)
Abdominal pain	29 (12.3)	22 (18.6)	51 (14.4)
Blood and lymphatic system disorders	183 (77.5)	95 (80.5)	278 (78.5)
Anaemia	126 (53.4)	65 (55.1)	191 (54.0)
Thrombocytopenia	61 (25.8)	34 (28.8)	95 (26.8)
Neutropenia	55 (23.3)	31 (26.3)	86 (24.3)
General disorders and administration site conditions	144 (61.0)	88 (74.6)	232 (65.5)
Fatigue	94 (39.8)	62 (52.5)	156 (44.1)
Asthenia	38 (16.1)	27 (22.9)	65 (18.4)
Oedema peripheral	24 (10.2)	22 (18.6)	46 (13.0)
Nervous system disorders	111 (47.0)	59 (50.0)	170 (48.0)
Headache	54 (22.9)	22 (18.6)	76 (21.5)
Dysgeusia	41 (17.4)	29 (24.6)	70 (19.8)
Dizziness	42 (17.8)	21 (17.8)	63 (17.8)
Musculoskeletal and connective tissue disorders	67 (28.4)	39 (33.1)	106 (29.9)
Arthralgia	29 (12.3)	16 (13.6)	45 (12.7)
Back pain	25 (10.6)	17 (14.4)	42 (11.9)
Investigations	67 (28.4)	37 (31.4)	104 (29.4)
Aspartate aminotransferase increased	24 (10.2)	16 (13.6)	40 (11.3)
Infections and infestations	63 (26.7)	33 (28.0)	96 (27.1)
Nasopharyngitis	28 (11.9)	15 (12.7)	43 (12.1)

System organ class / Preferred term	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Metabolism and nutrition disorders	58 (24.6)	33 (28.0)	91 (25.7)
Decreased appetite	58 (24.6)	33 (28.0)	91 (25.7)
Respiratory, thoracic and mediastinal disorders	58 (24.6)	22 (18.6)	80 (22.6)
Cough	28 (11.9)	12 (10.2)	40 (11.3)
Psychiatric disorders	24 (10.2)	15 (12.7)	39 (11.0)
Insomnia	24 (10.2)	15 (12.7)	39 (11.0)

Subjects are counted once per preferred term and once per system organ class at their highest reported grade. MedDRA version 25.0.

Table 14.3.5 Serious Adverse Events

Table 14.3.5 Serious Treatment-emergent Adverse Events by Preferred Term – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Anaemia	27 (11.4)	9 (7.6)	36 (10.2)
Malignant neoplasm progression	3 (1.3)	3 (2.5)	6 (1.7)
Myelodysplastic syndrome	2 (0.8)	4 (3.4)	6 (1.7)
Pneumonia	2 (0.8)	4 (3.4)	6 (1.7)
Neutropenia	3 (1.3)	2 (1.7)	5 (1.4)
Pneumonitis	5 (2.1)	0 (0.0)	5 (1.4)
Sepsis	4 (1.7)	0 (0.0)	4 (1.1)
Asthenia	1 (0.4)	1 (0.8)	2 (0.6)
Intestinal obstruction	2 (0.8)	0 (0.0)	2 (0.6)
Thrombocytopenia	2 (0.8)	0 (0.0)	2 (0.6)
Vomiting	2 (0.8)	0 (0.0)	2 (0.6)
Abdominal pain	0 (0.0)	1 (0.8)	1 (0.3)
Alanine aminotransferase increased	1 (0.4)	0 (0.0)	1 (0.3)
Arthralgia	1 (0.4)	0 (0.0)	1 (0.3)
Ascites	1 (0.4)	0 (0.0)	1 (0.3)
Aspartate aminotransferase increased	0 (0.0)	1 (0.8)	1 (0.3)
Blood creatinine increased	0 (0.0)	1 (0.8)	1 (0.3)
Dyspnoea	1 (0.4)	0 (0.0)	1 (0.3)
Headache	1 (0.4)	0 (0.0)	1 (0.3)
Pleural effusion	1 (0.4)	0 (0.0)	1 (0.3)
Pulmonary embolism	1 (0.4)	0 (0.0)	1 (0.3)

Subjects are counted once per preferred term. Percentages are based on N = 354.

Table 14.3.4a Treatment-Related Adverse Events

Table 14.3.4a Treatment-related Treatment-emergent Adverse Events Occurring in at Least 5% of Subjects – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Nausea	85 (36.0)	47 (39.8)	132 (37.3)
Anaemia	79 (33.5)	37 (31.4)	116 (32.8)
Fatigue	64 (27.1)	32 (27.1)	96 (27.1)
Constipation	41 (17.4)	30 (25.4)	71 (20.1)
Vomiting	40 (16.9)	27 (22.9)	67 (18.9)
Thrombocytopenia	39 (16.5)	22 (18.6)	61 (17.2)
Neutropenia	35 (14.8)	23 (19.5)	58 (16.4)
Decreased appetite	32 (13.6)	19 (16.1)	51 (14.4)
Headache	32 (13.6)	14 (11.9)	46 (13.0)
Dysgeusia	23 (9.7)	18 (15.3)	41 (11.6)
Dizziness	22 (9.3)	15 (12.7)	37 (10.5)
Asthenia	19 (8.1)	16 (13.6)	35 (9.9)
Diarrhoea	20 (8.5)	13 (11.0)	33 (9.3)
Back pain	19 (8.1)	13 (11.0)	32 (9.0)
Arthralgia	18 (7.6)	13 (11.0)	31 (8.8)
Oedema peripheral	15 (6.4)	15 (12.7)	30 (8.5)
Abdominal pain	15 (6.4)	15 (12.7)	30 (8.5)
Cough	18 (7.6)	8 (6.8)	26 (7.3)
Nasopharyngitis	18 (7.6)	7 (5.9)	25 (7.1)
Aspartate aminotransferase increased	13 (5.5)	12 (10.2)	25 (7.1)
Insomnia	13 (5.5)	11 (9.3)	24 (6.8)
Pyrexia	16 (6.8)	6 (5.1)	22 (6.2)
Alanine aminotransferase increased	16 (6.8)	6 (5.1)	22 (6.2)
Dyspepsia	11 (4.7)	8 (6.8)	19 (5.4)
Leukopenia	10 (4.2)	8 (6.8)	18 (5.1)

Relationship assessed by the investigator. Percentages are based on the number of subjects treated in each group. Overall N = 354.

Table 14.3.6 Adverse Events Leading to Discontinuation

Table 14.3.6 Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Anaemia	5 (2.1)	2 (1.7)	7 (2.0)
Malignant neoplasm progression	3 (1.3)	3 (2.5)	6 (1.7)
Myelodysplastic syndrome	1 (0.4)	3 (2.5)	4 (1.1)
Pneumonia	2 (0.8)	2 (1.7)	4 (1.1)
Sepsis	4 (1.7)	0 (0.0)	4 (1.1)
Intestinal obstruction	2 (0.8)	0 (0.0)	2 (0.6)
Neutropenia	2 (0.8)	0 (0.0)	2 (0.6)
Thrombocytopenia	2 (0.8)	0 (0.0)	2 (0.6)
Vomiting	1 (0.4)	1 (0.8)	2 (0.6)
Abdominal pain	1 (0.4)	0 (0.0)	1 (0.3)
Ascites	1 (0.4)	0 (0.0)	1 (0.3)
Aspartate aminotransferase increased	1 (0.4)	0 (0.0)	1 (0.3)
Asthenia	0 (0.0)	1 (0.8)	1 (0.3)
Back pain	0 (0.0)	1 (0.8)	1 (0.3)
Dyspnoea	1 (0.4)	0 (0.0)	1 (0.3)
Pleural effusion	1 (0.4)	0 (0.0)	1 (0.3)
Pneumonitis	1 (0.4)	0 (0.0)	1 (0.3)
Pulmonary embolism	1 (0.4)	0 (0.0)	1 (0.3)
Urinary tract infection	1 (0.4)	0 (0.0)	1 (0.3)

Percentages are based on N = 354.

Table 14.3.7 Deaths

Table 14.3.7 Deaths – Safety Analysis Set

Category, n (%)	Kelvaparib 300 mg (N=236)	Placebo (N=118)	Total (N=354)
Any death	51 (21.6)	21 (17.8)	72 (20.3)
Death within 30 days of last dose	20 (8.5)	8 (6.8)	28 (7.9)
Primary cause of death			
Disease progression	24 (10.2)	15 (12.7)	39 (11.0)
Adverse event	22 (9.3)	6 (5.1)	28 (7.9)
Other	5 (2.1)	0 (0.0)	5 (1.4)

Percentages are based on the number of subjects treated within each group.

Table 14.3.8 Laboratory Shift from Baseline to Worst Grade

Table 14.3.8 Shift from Baseline to Worst Post-baseline CTCAE Grade for Selected Laboratory Parameters – Safety Analysis Set

Laboratory parameter	Kelvaparib 300 mg: worst grade 3 n (%)	Kelvaparib 300 mg: worst grade 4 n (%)	Kelvaparib 300 mg: shift >= 2 grades n (%)	Placebo: worst grade 3 n (%)	Placebo: worst grade 4 n (%)	Placebo: shift >= 2 grades n (%)	Total: worst grade 3 n (%)	Total: worst grade 4 n (%)	Total: shift >= 2 grades n (%)
Haemoglobin	46 (19.5)	5 (2.1)	89 (37.7)	13 (11.0)	5 (4.2)	33 (28.0)	59 (16.7)	10 (2.8)	122 (34.5)
Platelets	26 (11.0)	7 (3.0)	51 (21.6)	10 (8.5)	1 (0.8)	17 (14.4)	36 (10.2)	8 (2.3)	68 (19.2)
Neutrophils	24 (10.2)	6 (2.5)	54 (22.9)	13 (11.0)	1 (0.8)	25 (21.2)	37 (10.5)	7 (2.0)	79 (22.3)
Lymphocytes	19 (8.1)	2 (0.8)	40 (16.9)	7 (5.9)	1 (0.8)	22 (18.6)	26 (7.3)	3 (0.8)	62 (17.5)
CA-125	34 (14.4)	15 (6.4)	74 (31.4)	23 (19.5)	1 (0.8)	45 (38.1)	57 (16.1)	16 (4.5)	119 (33.6)
ALT	11 (4.7)	5 (2.1)	26 (11.0)	6 (5.1)	1 (0.8)	16 (13.6)	17 (4.8)	6 (1.7)	42 (11.9)
AST	11 (4.7)	3 (1.3)	18 (7.6)	3 (2.5)	1 (0.8)	6 (5.1)	14 (4.0)	4 (1.1)	24 (6.8)
Creatinine	10 (4.2)	3 (1.3)	31 (13.1)	4 (3.4)	3 (2.5)	14 (11.9)	14 (4.0)	6 (1.7)	45 (12.7)
Lactate dehydrogenase	8 (3.4)	5 (2.1)	30 (12.7)	3 (2.5)	3 (2.5)	15 (12.7)	11 (3.1)	8 (2.3)	45 (12.7)
Albumin	6 (2.5)	5 (2.1)	23 (9.7)	4 (3.4)	1 (0.8)	13 (11.0)	10 (2.8)	6 (1.7)	36 (10.2)

CTCAE = Common Terminology Criteria for Adverse Events, version 5.0. Percentages are based on the number of subjects treated in each group. Overall N = 354.

Table 14.3.9 Adverse Event Onset Timing

Table 14.3.9 Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events – Safety Analysis Set

Preferred term	Events, n	Median time to onset (days)	Resolved, n (%)	Median time to resolution (days)
Nausea	223	36.0	197 (88.3)	9.0
Anaemia	191	32.0	151 (79.1)	12.0
Fatigue	156	32.0	133 (85.3)	12.0
Vomiting	114	24.5	100 (87.7)	10.0
Constipation	112	32.5	97 (86.6)	9.0
Thrombocytopenia	95	31.0	81 (85.3)	10.0
Decreased appetite	91	31.0	71 (78.0)	9.0
Neutropenia	88	39.5	70 (79.5)	10.0
Headache	76	37.5	66 (86.8)	8.5
Dysgeusia	70	35.5	62 (88.6)	12.0
Diarrhoea	69	25.0	57 (82.6)	9.0
Asthenia	65	18.0	55 (84.6)	9.0
Dizziness	63	29.0	54 (85.7)	10.5
Abdominal pain	51	39.0	42 (82.4)	9.0
Oedema peripheral	46	35.5	41 (89.1)	10.0
Arthralgia	45	27.0	37 (82.2)	12.0
Nasopharyngitis	43	35.0	35 (81.4)	12.0
Back pain	42	18.0	34 (81.0)	11.5
Cough	40	42.5	36 (90.0)	12.0
Aspartate aminotransferase increased	40	41.5	32 (80.0)	12.5

Events are counted at the event level rather than the subject level. NE = not estimable.