TABLES, LISTINGS AND FIGURES

Study KLV-7702 — A Phase 1 Open-label, Dose-escalation Study of Kelvaparib in Adults with Advanced Solid Tumours

Kelvaparib (KLV-3308)

Marnholt Oncology GmbH

Indication: Advanced solid tumours with a homologous recombination repair defect

This pack contains the post-text tables, listings and figures supporting the clinical study report for study KLV-7702. All displays derive from the analysis datasets locked on 1 May 2019.

Contents

- Table 14.1.1 Demographic and Baseline Characteristics
- Table 14.1.2 Subject Disposition
- Table 14.1.3 Extent of Exposure
- Table 14.2.1 Best Overall Response
- Table 14.2.2 Time-to-Event Analyses
- Table 14.3.1 Overview of Adverse Events
- Table 14.3.2a Adverse Events by System Organ Class and Preferred Term
- Table 14.3.5 Serious Adverse Events
- Table 14.3.4a Treatment-Related Adverse Events
- Table 14.3.6 Adverse Events Leading to Discontinuation
- Table 14.3.7 Deaths
- Table 14.3.8 Laboratory Shift from Baseline to Worst Grade
- Table 14.3.9 Adverse Event Onset Timing

Table 14.1.1 Demographic and Baseline Characteristics

Table 14.1.1 Demographic and Baseline Characteristics – Safety Analysis Set

Characteristic	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Age (years)					
Mean (SD)	65.8 (2.04)	66.2 (10.96)	66.4 (8.86)	66.6 (9.79)	66.3 (8.78)
Median	66.0	65.0	68.0	67.5	67.0
Min, max	63, 68	50, 86	50, 84	47, 80	47, 86
>= 65 years, n (%)	4 (66.7)	5 (55.6)	20 (64.5)	8 (66.7)	37 (63.8)
Sex, n (%)					
Male	3 (50.0)	3 (33.3)	21 (67.7)	10 (83.3)	37 (63.8)
Female	3 (50.0)	6 (66.7)	10 (32.3)	2 (16.7)	21 (36.2)
Race, n (%)					
White	6 (100.0)	5 (55.6)	26 (83.9)	7 (58.3)	44 (75.9)
Asian	0 (0.0)	1 (11.1)	1 (3.2)	3 (25.0)	5 (8.6)
Black or African American	0 (0.0)	2 (22.2)	3 (9.7)	1 (8.3)	6 (10.3)
Other	0 (0.0)	1 (11.1)	1 (3.2)	1 (8.3)	3 (5.2)
ECOG performance status, n (%)					
0	3 (50.0)	4 (44.4)	15 (48.4)	6 (50.0)	28 (48.3)
1	2 (33.3)	4 (44.4)	12 (38.7)	5 (41.7)	23 (39.7)
2	1 (16.7)	1 (11.1)	4 (12.9)	1 (8.3)	7 (12.1)
BRCA mutation status, n (%)					
BRCA1 mutated	0 (0.0)	2 (22.2)	10 (32.3)	3 (25.0)	15 (25.9)
BRCA2 mutated	3 (50.0)	0 (0.0)	7 (22.6)	2 (16.7)	12 (20.7)
BRCA wild-type	3 (50.0)	7 (77.8)	14 (45.2)	7 (58.3)	31 (53.4)
Homologous recombination deficiency, n (%)	3 (50.0)	4 (44.4)	17 (54.8)	8 (66.7)	32 (55.2)
Number of prior lines of therapy					
Median	1.5	1.0	2.0	1.0	1.0
>= 3, n (%)	2 (33.3)	0 (0.0)	6 (19.4)	1 (8.3)	9 (15.5)

SD = standard deviation; ECOG = Eastern Cooperative Oncology Group. Percentages are based on the number of subjects in the safety analysis set within each group.

Table 14.1.2 Subject Disposition

Table 14.1.2 Subject Disposition – Safety Analysis Set

Disposition	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Subjects treated, n	6	9	31	12	58
Treatment ongoing at data cut-off, n (%)	3 (50.0)	2 (22.2)	8 (25.8)	4 (33.3)	17 (29.3)
Discontinued study treatment, n (%)	3 (50.0)	7 (77.8)	23 (74.2)	8 (66.7)	41 (70.7)
Disease progression	1 (16.7)	3 (33.3)	7 (22.6)	3 (25.0)	14 (24.1)
Adverse event	0 (0.0)	2 (22.2)	1 (3.2)	1 (8.3)	4 (6.9)
Death	1 (16.7)	0 (0.0)	4 (12.9)	1 (8.3)	6 (10.3)
Withdrawal by subject	1 (16.7)	1 (11.1)	5 (16.1)	1 (8.3)	8 (13.8)
Physician decision	0 (0.0)	1 (11.1)	2 (6.5)	2 (16.7)	5 (8.6)
Lost to follow-up	0 (0.0)	0 (0.0)	3 (9.7)	0 (0.0)	3 (5.2)
Protocol deviation	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)

Percentages are based on the number of subjects treated within each group.

Table 14.1.3 Extent of Exposure

Table 14.1.3 Extent of Exposure – Safety Analysis Set

Exposure parameter	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Duration of treatment (days)					
Median	171.0	117.0	118.0	147.0	121.5
Q1, Q3	107.8, 198.2	74.0, 129.0	83.5, 156.0	110.5, 226.2	86.0, 164.8
Min, max	65, 236	8, 308	8, 238	8, 448	8, 448
Cycles started, median	6.0	4.0	4.0	5.5	4.0
Relative dose intensity (%), median	83.8	83.9	84.7	82.9	84.2
Any dose reduction, n (%)	0 (0.0)	2 (22.2)	6 (19.4)	4 (33.3)	12 (20.7)
Any dose interruption, n (%)	1 (16.7)	3 (33.3)	5 (16.1)	1 (8.3)	10 (17.2)

Q1 = first quartile; Q3 = third quartile. Relative dose intensity is the total dose received divided by the total dose planned over the treatment period.

Table 14.2.1 Best Overall Response

Table 14.2.1 Best Overall Response by RECIST 1.1 – Full Analysis Set

Best overall response	100 mg (N=6)	200 mg (N=8)	300 mg (N=30)	400 mg (N=11)	Total (N=55)
Complete response (CR)	0 (0.0)	1 (12.5)	2 (6.7)	1 (9.1)	4 (7.3)
Partial response (PR)	0 (0.0)	2 (25.0)	6 (20.0)	4 (36.4)	12 (21.8)
Stable disease	4 (66.7)	1 (12.5)	13 (43.3)	3 (27.3)	21 (38.2)
Progressive disease	2 (33.3)	4 (50.0)	9 (30.0)	3 (27.3)	18 (32.7)
Not evaluable	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Objective response rate, n (%)	0 (0.0)	3 (37.5)	8 (26.7)	5 (45.5)	16 (29.1)
95% CI	(0.0, 45.9)	(8.5, 75.5)	(12.3, 45.9)	(16.7, 76.6)	(17.6, 42.9)

CI = confidence interval. Assessed according to RECIST version 1.1. Confidence intervals are exact (Clopper-Pearson).

Table 14.2.2 Time-to-Event Analyses

Table 14.2.2 Kaplan-Meier Estimates of Time-to-event Endpoints – Full Analysis Set

Endpoint	100 mg (N=6)	200 mg (N=8)	300 mg (N=30)	400 mg (N=11)	Total (N=55)
Duration of response, median (months)	NE	NE	9.8	NE	NE
25th, 75th percentile	NE	NE	(9.8, NE)	(7.1, NE)	(9.8, NE)
Progression-free survival, median (months)	3.1	12.4	7.1	8.7	8.5
25th, 75th percentile	(2.9, 3.3)	(9.7, 12.7)	(3.0, 15.9)	(2.5, NE)	(3.0, 12.7)
Overall survival, median (months)	NE	NE	NE	NE	NE
25th, 75th percentile	NE	(12.7, NE)	NE	NE	NE

NE = not estimable. Duration of response is analysed among responding subjects only.

Table 14.3.1 Overview of Adverse Events

Table 14.3.1 Overview of Treatment-emergent Adverse Events – Safety Analysis Set

Category, n (%)	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Any TEAE	6 (100.0)	9 (100.0)	31 (100.0)	12 (100.0)	58 (100.0)
Any treatment-related TEAE	5 (83.3)	9 (100.0)	31 (100.0)	11 (91.7)	56 (96.6)
Any grade >= 3 TEAE	3 (50.0)	7 (77.8)	21 (67.7)	10 (83.3)	41 (70.7)
Any serious TEAE	1 (16.7)	2 (22.2)	12 (38.7)	7 (58.3)	22 (37.9)
Any TEAE leading to dose reduction	0 (0.0)	2 (22.2)	6 (19.4)	4 (33.3)	12 (20.7)
Any TEAE leading to discontinuation	1 (16.7)	2 (22.2)	5 (16.1)	2 (16.7)	10 (17.2)
Any fatal TEAE	1 (16.7)	0 (0.0)	4 (12.9)	1 (8.3)	6 (10.3)

TEAE = treatment-emergent adverse event. A subject is counted once within each category.

Table 14.3.2a Adverse Events by System Organ Class and Preferred Term

Table 14.3.2a Treatment-emergent Adverse Events Occurring in at Least 10% of Subjects by System Organ Class and Preferred Term – Safety Analysis Set

System organ class / Preferred term	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Gastrointestinal disorders	5 (83.3)	9 (100.0)	30 (96.8)	10 (83.3)	54 (93.1)
Nausea	2 (33.3)	5 (55.6)	21 (67.7)	9 (75.0)	37 (63.8)
Constipation	1 (16.7)	2 (22.2)	16 (51.6)	3 (25.0)	22 (37.9)
Vomiting	1 (16.7)	3 (33.3)	12 (38.7)	2 (16.7)	18 (31.0)
Abdominal pain	3 (50.0)	4 (44.4)	7 (22.6)	3 (25.0)	17 (29.3)
Diarrhoea	3 (50.0)	2 (22.2)	7 (22.6)	1 (8.3)	13 (22.4)
Dyspepsia	1 (16.7)	2 (22.2)	6 (19.4)	3 (25.0)	12 (20.7)
Blood and lymphatic system disorders	4 (66.7)	8 (88.9)	23 (74.2)	10 (83.3)	45 (77.6)
Anaemia	2 (33.3)	4 (44.4)	19 (61.3)	10 (83.3)	35 (60.3)
Neutropenia	2 (33.3)	4 (44.4)	8 (25.8)	3 (25.0)	17 (29.3)
Thrombocytopenia	1 (16.7)	2 (22.2)	10 (32.3)	3 (25.0)	16 (27.6)
Lymphopenia	1 (16.7)	2 (22.2)	4 (12.9)	3 (25.0)	10 (17.2)
Leukopenia	1 (16.7)	1 (11.1)	2 (6.5)	2 (16.7)	6 (10.3)
General disorders and administration site conditions	4 (66.7)	6 (66.7)	25 (80.6)	8 (66.7)	43 (74.1)
Fatigue	4 (66.7)	4 (44.4)	18 (58.1)	7 (58.3)	33 (56.9)
Asthenia	0 (0.0)	2 (22.2)	6 (19.4)	3 (25.0)	11 (19.0)
Pyrexia	1 (16.7)	0 (0.0)	7 (22.6)	1 (8.3)	9 (15.5)
Nervous system disorders	2 (33.3)	3 (33.3)	16 (51.6)	5 (41.7)	26 (44.8)
Headache	0 (0.0)	1 (11.1)	10 (32.3)	4 (33.3)	15 (25.9)
Dizziness	2 (33.3)	1 (11.1)	6 (19.4)	0 (0.0)	9 (15.5)
Dysgeusia	1 (16.7)	1 (11.1)	4 (12.9)	1 (8.3)	7 (12.1)
Infections and infestations	2 (33.3)	4 (44.4)	14 (45.2)	5 (41.7)	25 (43.1)
Urinary tract infection	1 (16.7)	2 (22.2)	4 (12.9)	4 (33.3)	11 (19.0)
Nasopharyngitis	1 (16.7)	2 (22.2)	5 (16.1)	1 (8.3)	9 (15.5)
Metabolism and nutrition disorders	2 (33.3)	5 (55.6)	12 (38.7)	5 (41.7)	24 (41.4)

System organ class / Preferred term	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Decreased appetite	2 (33.3)	5 (55.6)	12 (38.7)	5 (41.7)	24 (41.4)
Investigations	1 (16.7)	4 (44.4)	13 (41.9)	6 (50.0)	24 (41.4)
Alanine aminotransferase increased	0 (0.0)	1 (11.1)	6 (19.4)	2 (16.7)	9 (15.5)
Aspartate aminotransferase increased	1 (16.7)	0 (0.0)	4 (12.9)	3 (25.0)	8 (13.8)
Weight decreased	0 (0.0)	2 (22.2)	3 (9.7)	2 (16.7)	7 (12.1)
Musculoskeletal and connective tissue disorders	0 (0.0)	4 (44.4)	13 (41.9)	3 (25.0)	20 (34.5)
Back pain	0 (0.0)	2 (22.2)	7 (22.6)	0 (0.0)	9 (15.5)
Arthralgia	0 (0.0)	1 (11.1)	5 (16.1)	2 (16.7)	8 (13.8)
Respiratory, thoracic and mediastinal disorders	0 (0.0)	1 (11.1)	8 (25.8)	3 (25.0)	12 (20.7)
Dyspnoea	0 (0.0)	1 (11.1)	3 (9.7)	2 (16.7)	6 (10.3)
Cough	0 (0.0)	0 (0.0)	5 (16.1)	1 (8.3)	6 (10.3)
Psychiatric disorders	2 (33.3)	1 (11.1)	7 (22.6)	1 (8.3)	11 (19.0)
Insomnia	2 (33.3)	1 (11.1)	7 (22.6)	1 (8.3)	11 (19.0)
Vascular disorders	0 (0.0)	1 (11.1)	7 (22.6)	0 (0.0)	8 (13.8)
Hypertension	0 (0.0)	1 (11.1)	7 (22.6)	0 (0.0)	8 (13.8)

Subjects are counted once per preferred term and once per system organ class at their highest reported grade. MedDRA version 22.0.

Table 14.3.5 Serious Adverse Events

Table 14.3.5 Serious Treatment-emergent Adverse Events by Preferred Term – Safety Analysis Set

Preferred term	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Anaemia	0 (0.0)	1 (11.1)	4 (12.9)	4 (33.3)	9 (15.5)
Blood creatinine increased	0 (0.0)	2 (22.2)	0 (0.0)	0 (0.0)	2 (3.4)
Malignant neoplasm progression	0 (0.0)	0 (0.0)	1 (3.2)	1 (8.3)	2 (3.4)
Pneumonia	0 (0.0)	0 (0.0)	2 (6.5)	0 (0.0)	2 (3.4)
Ascites	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.7)
Aspartate aminotransferase increased	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Cough	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Intestinal obstruction	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Lymphopenia	0 (0.0)	0 (0.0)	0 (0.0)	1 (8.3)	1 (1.7)
Myelodysplastic syndrome	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Nausea	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Neutropenia	0 (0.0)	0 (0.0)	0 (0.0)	1 (8.3)	1 (1.7)
Sepsis	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Thrombocytopenia	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)

Subjects are counted once per preferred term. Percentages are based on N = 58.

Table 14.3.4a Treatment-Related Adverse Events

Table 14.3.4a Treatment-related Treatment-emergent Adverse Events Occurring in at Least 5% of Subjects – Safety Analysis Set

Preferred term	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Fatigue	3 (50.0)	2 (22.2)	12 (38.7)	4 (33.3)	21 (36.2)
Anaemia	0 (0.0)	2 (22.2)	13 (41.9)	6 (50.0)	21 (36.2)
Decreased appetite	1 (16.7)	5 (55.6)	8 (25.8)	2 (16.7)	16 (27.6)
Nausea	1 (16.7)	1 (11.1)	9 (29.0)	5 (41.7)	16 (27.6)
Constipation	0 (0.0)	1 (11.1)	10 (32.3)	2 (16.7)	13 (22.4)
Abdominal pain	3 (50.0)	3 (33.3)	5 (16.1)	1 (8.3)	12 (20.7)
Vomiting	0 (0.0)	3 (33.3)	7 (22.6)	1 (8.3)	11 (19.0)
Insomnia	2 (33.3)	1 (11.1)	6 (19.4)	1 (8.3)	10 (17.2)
Urinary tract infection	1 (16.7)	2 (22.2)	4 (12.9)	3 (25.0)	10 (17.2)
Neutropenia	0 (0.0)	1 (11.1)	5 (16.1)	3 (25.0)	9 (15.5)
Headache	0 (0.0)	1 (11.1)	6 (19.4)	1 (8.3)	8 (13.8)
Thrombocytopenia	0 (0.0)	2 (22.2)	4 (12.9)	2 (16.7)	8 (13.8)
Lymphopenia	1 (16.7)	1 (11.1)	3 (9.7)	1 (8.3)	6 (10.3)
Diarrhoea	1 (16.7)	1 (11.1)	3 (9.7)	1 (8.3)	6 (10.3)
Asthenia	0 (0.0)	1 (11.1)	3 (9.7)	2 (16.7)	6 (10.3)
Weight decreased	0 (0.0)	2 (22.2)	2 (6.5)	2 (16.7)	6 (10.3)
Dyspepsia	1 (16.7)	1 (11.1)	3 (9.7)	0 (0.0)	5 (8.6)
Leukopenia	1 (16.7)	1 (11.1)	1 (3.2)	2 (16.7)	5 (8.6)
Nasopharyngitis	0 (0.0)	2 (22.2)	2 (6.5)	1 (8.3)	5 (8.6)
Alanine aminotransferase increased	0 (0.0)	1 (11.1)	2 (6.5)	2 (16.7)	5 (8.6)
Arthralgia	0 (0.0)	1 (11.1)	2 (6.5)	2 (16.7)	5 (8.6)
Pyrexia	0 (0.0)	0 (0.0)	4 (12.9)	1 (8.3)	5 (8.6)
Dysgeusia	1 (16.7)	1 (11.1)	2 (6.5)	0 (0.0)	4 (6.9)
Dizziness	1 (16.7)	0 (0.0)	3 (9.7)	0 (0.0)	4 (6.9)
Back pain	0 (0.0)	1 (11.1)	3 (9.7)	0 (0.0)	4 (6.9)
Hypertension	0 (0.0)	1 (11.1)	3 (9.7)	0 (0.0)	4 (6.9)

Preferred term	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Rash	1 (16.7)	1 (11.1)	1 (3.2)	0 (0.0)	3 (5.2)
Myalgia	0 (0.0)	1 (11.1)	1 (3.2)	1 (8.3)	3 (5.2)
Dyspnoea	0 (0.0)	1 (11.1)	1 (3.2)	1 (8.3)	3 (5.2)

Relationship assessed by the investigator. Percentages are based on the number of subjects treated in each group. Overall N = 58.

Table 14.3.6 Adverse Events Leading to Discontinuation

Table 14.3.6 Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment – Safety Analysis Set

Preferred term	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Malignant neoplasm progression	0 (0.0)	0 (0.0)	1 (3.2)	1 (8.3)	2 (3.4)
Sepsis	0 (0.0)	0 (0.0)	2 (6.5)	0 (0.0)	2 (3.4)
Ascites	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.7)
Intestinal obstruction	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Lymphopenia	0 (0.0)	0 (0.0)	0 (0.0)	1 (8.3)	1 (1.7)
Myelodysplastic syndrome	0 (0.0)	1 (11.1)	0 (0.0)	0 (0.0)	1 (1.7)
Pneumonia	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)
Thrombocytopenia	0 (0.0)	1 (11.1)	0 (0.0)	0 (0.0)	1 (1.7)

Percentages are based on N = 58.

Table 14.3.7 Deaths

Table 14.3.7 Deaths – Safety Analysis Set

Category, n (%)	100 mg (N=6)	200 mg (N=9)	300 mg (N=31)	400 mg (N=12)	Total (N=58)
Any death	1 (16.7)	1 (11.1)	5 (16.1)	1 (8.3)	8 (13.8)
Death within 30 days of last dose	1 (16.7)	0 (0.0)	4 (12.9)	1 (8.3)	6 (10.3)
Primary cause of death					
Disease progression	0 (0.0)	1 (11.1)	0 (0.0)	0 (0.0)	1 (1.7)
Adverse event	1 (16.7)	0 (0.0)	4 (12.9)	1 (8.3)	6 (10.3)
Other	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	1 (1.7)

Percentages are based on the number of subjects treated within each group.

Table 14.3.8 Laboratory Shift from Baseline to Worst Grade

Table 14.3.8 Shift from Baseline to Worst Post-baseline CTCAE Grade for Selected Laboratory Parameters – Safety Analysis Set

Laboratory parameter	100 mg: worst grade 3 n (%)	100 mg: worst grade 4 n (%)	100 mg: shift >= 2 g rades n (%)	200 mg: worst grade 3 n (%)	200 mg: worst grade 4 n (%)	200 mg: shift >= 2 g rades n (%)	300 mg: worst grade 3 n (%)	300 mg: worst grade 4 n (%)	300 mg: shift >= 2 g rades n (%)	400 mg: worst grade 3 n (%)	400 mg: worst grade 4 n (%)	400 mg: shift >= 2 g rades n (%)	Total: worst grade 3 n (%)	Total: worst grade 4 n (%)	Total: shift >= 2 g rades n (%)
Haemoglobin	0 (0.0)	0 (0.0)	2 (33.3)	1 (11.1)	2 (22.2)	4 (44.4)	10 (32.3)	4 (12.9)	17 (54.8)	3 (25.0)	0 (0.0)	5 (41.7)	14 (24.1)	6 (10.3)	28 (48.3)
Platelets	2 (33.3)	0 (0.0)	2 (33.3)	1 (11.1)	0 (0.0)	4 (44.4)	2 (6.5)	1 (3.2)	6 (19.4)	0 (0.0)	0 (0.0)	2 (16.7)	5 (8.6)	1 (1.7)	14 (24.1)
Neutrophils	0 (0.0)	0 (0.0)	2 (33.3)	1 (11.1)	1 (11.1)	3 (33.3)	4 (12.9)	0 (0.0)	4 (12.9)	1 (8.3)	0 (0.0)	1 (8.3)	6 (10.3)	1 (1.7)	10 (17.2)
Lymphocytes	0 (0.0)	0 (0.0)	0 (0.0)	1 (11.1)	0 (0.0)	0 (0.0)	4 (12.9)	1 (3.2)	6 (19.4)	2 (16.7)	0 (0.0)	2 (16.7)	7 (12.1)	1 (1.7)	8 (13.8)
CA-125	2 (33.3)	0 (0.0)	1 (16.7)	0 (0.0)	0 (0.0)	2 (22.2)	4 (12.9)	2 (6.5)	10 (32.3)	2 (16.7)	0 (0.0)	4 (33.3)	8 (13.8)	2 (3.4)	17 (29.3)
ALT	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	5 (16.1)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.7)	0 (0.0)	5 (8.6)
AST	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (11.1)	1 (11.1)	2 (6.5)	0 (0.0)	3 (9.7)	2 (16.7)	1 (8.3)	3 (25.0)	4 (6.9)	2 (3.4)	7 (12.1)
Creatinine	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	3 (9.7)	0 (0.0)	3 (9.7)	1 (8.3)	0 (0.0)	2 (16.7)	4 (6.9)	0 (0.0)	5 (8.6)
Lactate dehydrogenase	1 (16.7)	0 (0.0)	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.2)	6 (19.4)	1 (8.3)	1 (8.3)	2 (16.7)	2 (3.4)	2 (3.4)	9 (15.5)
Albumin	0 (0.0)	0 (0.0)	1 (16.7)	0 (0.0)	0 (0.0)	0 (0.0)	1 (3.2)	0 (0.0)	2 (6.5)	1 (8.3)	1 (8.3)	2 (16.7)	2 (3.4)	1 (1.7)	5 (8.6)

CTCAE = Common Terminology Criteria for Adverse Events, version 5.0. Percentages are based on the number of subjects treated in each group. Overall N = 58.

Table 14.3.9 Adverse Event Onset Timing

Table 14.3.9 Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events – Safety Analysis Set

Preferred term	Events, n	Median time to onset (days)	Resolved, n (%)	Median time to resolution (days)
Nausea	37	24.0	29 (78.4)	9.0
Anaemia	35	40.0	28 (80.0)	14.0
Fatigue	33	23.0	31 (93.9)	9.0
Decreased appetite	24	44.5	20 (83.3)	12.0
Constipation	22	39.0	19 (86.4)	8.0
Vomiting	18	23.0	14 (77.8)	5.5
Abdominal pain	17	46.0	15 (88.2)	10.0
Neutropenia	17	17.0	12 (70.6)	9.0
Thrombocytopenia	16	19.0	15 (93.8)	7.0
Headache	15	44.0	12 (80.0)	16.5
Diarrhoea	13	44.0	12 (92.3)	8.0
Dyspepsia	12	22.5	10 (83.3)	10.5
Insomnia	11	1.0	10 (90.9)	9.5
Urinary tract infection	11	51.0	8 (72.7)	15.5
Asthenia	11	30.0	8 (72.7)	8.5
Lymphopenia	10	38.0	10 (100.0)	17.0
Nasopharyngitis	9	8.0	6 (66.7)	12.0
Dizziness	9	30.0	6 (66.7)	9.5
Pyrexia	9	22.0	7 (77.8)	13.0
Alanine aminotransferase increased	9	43.0	8 (88.9)	6.5

Events are counted at the event level rather than the subject level. NE = not estimable.