TABLES, LISTINGS AND FIGURES

Study KLV-7704 — A Phase 2 Study of Kelvaparib in Recurrent High-grade Serous Ovarian Carcinoma

Kelvaparib (KLV-3308)

Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

This pack contains the post-text tables, listings and figures supporting the clinical study report for study KLV-7704. All displays derive from the analysis datasets locked on 28 May 2020.

Contents

- Table 14.1.1 Demographic and Baseline Characteristics
- Table 14.1.2 Subject Disposition
- Table 14.1.3 Extent of Exposure
- Table 14.2.1 Best Overall Response
- Table 14.2.2 Time-to-Event Analyses
- Table 14.3.1 Overview of Adverse Events
- Table 14.3.2a Adverse Events by System Organ Class and Preferred Term
- Table 14.3.5 Serious Adverse Events
- Table 14.3.4a Treatment-Related Adverse Events
- Table 14.3.6 Adverse Events Leading to Discontinuation
- Table 14.3.7 Deaths
- Table 14.3.8 Laboratory Shift from Baseline to Worst Grade
- Table 14.3.9 Adverse Event Onset Timing

Table 14.1.1 Demographic and Baseline Characteristics

Table 14.1.1 Demographic and Baseline Characteristics – Safety Analysis Set

Characteristic	Kelvaparib 300 mg (N=81)	Total (N=81)
Age (years)		
Mean (SD)	64.9 (8.83)	64.9 (8.83)
Median	65.0	65.0
Min, max	38, 88	38, 88
>= 65 years, n (%)	41 (50.6)	41 (50.6)
Sex, n (%)		
Male	0 (0.0)	0 (0.0)
Female	81 (100.0)	81 (100.0)
Race, n (%)		
White	70 (86.4)	70 (86.4)
Asian	6 (7.4)	6 (7.4)
Black or African American	3 (3.7)	3 (3.7)
Other	2 (2.5)	2 (2.5)
ECOG performance status, n (%)		
0	26 (32.1)	26 (32.1)
1	47 (58.0)	47 (58.0)
2	8 (9.9)	8 (9.9)
BRCA mutation status, n (%)		
BRCA1 mutated	19 (23.5)	19 (23.5)
BRCA2 mutated	10 (12.3)	10 (12.3)
BRCA wild-type	52 (64.2)	52 (64.2)
Homologous recombination deficiency, n (%)	45 (55.6)	45 (55.6)
Platinum-sensitive disease, n (%)	56 (69.1)	56 (69.1)
Residual disease after cytoreductive surgery, n (%)	33 (40.7)	33 (40.7)
Number of prior lines of platinum-based therapy		
Median	2.0	2.0

Characteristic	Kelvaparib 300 mg (N=81)	Total (N=81)
>= 3, n (%)	25 (30.9)	25 (30.9)

SD = standard deviation; ECOG = Eastern Cooperative Oncology Group. Percentages are based on the number of subjects in the safety analysis set within each group.

Table 14.1.2 Subject Disposition

Table 14.1.2 Subject Disposition – Safety Analysis Set

Disposition	Kelvaparib 300 mg (N=81)	Total (N=81)
Subjects treated, n	81	81
Treatment ongoing at data cut-off, n (%)	20 (24.7)	20 (24.7)
Discontinued study treatment, n (%)	61 (75.3)	61 (75.3)
Disease progression	15 (18.5)	15 (18.5)
Adverse event	4 (4.9)	4 (4.9)
Death	5 (6.2)	5 (6.2)
Withdrawal by subject	11 (13.6)	11 (13.6)
Physician decision	17 (21.0)	17 (21.0)
Lost to follow-up	7 (8.6)	7 (8.6)
Protocol deviation	2 (2.5)	2 (2.5)

Percentages are based on the number of subjects treated within each group.

Table 14.1.3 Extent of Exposure

Table 14.1.3 Extent of Exposure – Full Analysis Set

Exposure parameter	Kelvaparib 300 mg (N=81)	Total (N=81)
Duration of treatment (weeks)		
Median	105.0	105.0
Q1, Q3	72.0, 152.0	72.0, 152.0
Min, max	8, 321	8, 321
Cycles started, median	4.0	4.0
Relative dose intensity (%), median	86.3	86.3
Any dose reduction, n (%)	8 (9.9)	8 (9.9)
Any dose interruption, n (%)	25 (30.9)	25 (30.9)

Q1 = first quartile; Q3 = third quartile. Relative dose intensity is the total dose received divided by the total dose planned over the treatment period.
Exposure is summarised to the end of Cycle 6 (day 168).

Table 14.2.1 Best Overall Response

Table 14.2.1 Best Overall Response by RECIST 1.1 – Full Analysis Set

Best overall response	Kelvaparib 300 mg (N=80)	Total (N=80)
Complete response (CR)	9 (11.2)	9 (11.2)
Partial response (PR)	24 (30.0)	24 (30.0)
Stable disease	29 (36.2)	29 (36.2)
Progressive disease	18 (22.5)	18 (22.5)
Not evaluable	0 (0.0)	0 (0.0)
Objective response rate, n (%)	33 (41.2)	33 (41.2)
95% CI	(30.4, 52.8)	(30.4, 52.8)

CI = confidence interval. Assessed according to RECIST version 1.1. Confidence intervals are exact (Clopper-Pearson).

Table 14.2.2 Time-to-Event Analyses

Table 14.2.2 Kaplan-Meier Estimates of Time-to-event Endpoints – Full Analysis Set

Endpoint	Kelvaparib 300 mg (N=80)	Total (N=80)
Duration of response, median (months)	9.0	9.0
25th, 75th percentile	(8.2, NE)	(8.2, NE)
Progression-free survival, median (months)	6.7	6.7
25th, 75th percentile	(3.1, 15.0)	(3.1, 15.0)
Overall survival, median (months)	21.7	21.7
25th, 75th percentile	(21.7, 21.7)	(21.7, 21.7)

NE = not estimable. Duration of response is analysed among responding subjects only.

Table 14.3.1 Overview of Adverse Events

Table 14.3.1 Overview of Treatment-emergent Adverse Events – Safety Analysis Set

Category, n (%)	Kelvaparib 300 mg (N=81)	Total (N=81)
Any TEAE	81 (100.0)	81 (100.0)
Any treatment-related TEAE	79 (97.5)	79 (97.5)
Any grade >= 3 TEAE	30 (37.0)	30 (37.0)
Any serious TEAE	17 (21.0)	17 (21.0)
Any TEAE leading to dose reduction	8 (9.9)	8 (9.9)
Any TEAE leading to discontinuation	8 (9.9)	8 (9.9)
Any fatal TEAE	6 (7.4)	6 (7.4)

TEAE = treatment-emergent adverse event. A subject is counted once within each category.

Table 14.3.2a Adverse Events by System Organ Class and Preferred Term

Table 14.3.2a Treatment-emergent Adverse Events Occurring in at Least 10% of Subjects by System Organ Class and Preferred Term – Safety Analysis Set

System organ class / Preferred term	Kelvaparib 300 mg (N=81)	Total (N=81)
Gastrointestinal disorders	70 (86.4)	70 (86.4)
Nausea	55 (67.9)	55 (67.9)
Vomiting	23 (28.4)	23 (28.4)
Constipation	21 (25.9)	21 (25.9)
Diarrhoea	14 (17.3)	14 (17.3)
Abdominal pain	10 (12.3)	10 (12.3)
Blood and lymphatic system disorders	54 (66.7)	54 (66.7)
Anaemia	34 (42.0)	34 (42.0)
Thrombocytopenia	17 (21.0)	17 (21.0)
Neutropenia	14 (17.3)	14 (17.3)
General disorders and administration site conditions	48 (59.3)	48 (59.3)
Fatigue	37 (45.7)	37 (45.7)
Asthenia	13 (16.0)	13 (16.0)
Oedema peripheral	10 (12.3)	10 (12.3)
Nervous system disorders	39 (48.1)	39 (48.1)
Headache	20 (24.7)	20 (24.7)
Dysgeusia	18 (22.2)	18 (22.2)
Dizziness	9 (11.1)	9 (11.1)
Respiratory, thoracic and mediastinal disorders	23 (28.4)	23 (28.4)
Cough	17 (21.0)	17 (21.0)
Musculoskeletal and connective tissue disorders	23 (28.4)	23 (28.4)
Arthralgia	14 (17.3)	14 (17.3)
Infections and infestations	22 (27.2)	22 (27.2)
Urinary tract infection	10 (12.3)	10 (12.3)
Nasopharyngitis	10 (12.3)	10 (12.3)

System organ class / Preferred term	Kelvaparib 300 mg (N=81)	Total (N=81)
Investigations	21 (25.9)	21 (25.9)
Metabolism and nutrition disorders	19 (23.5)	19 (23.5)
Decreased appetite	19 (23.5)	19 (23.5)
Psychiatric disorders	10 (12.3)	10 (12.3)
Insomnia	10 (12.3)	10 (12.3)

Subjects are counted once per preferred term and once per system organ class at their highest reported grade. MedDRA version 23.1.

Table 14.3.5 Serious Adverse Events

Table 14.3.5 Serious Treatment-emergent Adverse Events by Preferred Term – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=81)	Total (N=81)
Anaemia	5 (6.2)	5 (6.2)
Malignant neoplasm progression	2 (2.5)	2 (2.5)
Ascites	1 (1.2)	1 (1.2)
Aspartate aminotransferase increased	1 (1.2)	1 (1.2)
Diarrhoea	1 (1.2)	1 (1.2)
Intestinal obstruction	1 (1.2)	1 (1.2)
Lymphopenia	1 (1.2)	1 (1.2)
Myelodysplastic syndrome	1 (1.2)	1 (1.2)
Pneumonia	1 (1.2)	1 (1.2)
Pneumonitis	1 (1.2)	1 (1.2)
Sepsis	1 (1.2)	1 (1.2)
Vomiting	1 (1.2)	1 (1.2)

Subjects are counted once per preferred term. Percentages are based on N = 81.

Table 14.3.4a Treatment-Related Adverse Events

Table 14.3.4a Treatment-related Treatment-emergent Adverse Events Occurring in at Least 5% of Subjects – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=81)	Total (N=81)
Nausea	37 (45.7)	37 (45.7)
Fatigue	22 (27.2)	22 (27.2)
Anaemia	20 (24.7)	20 (24.7)
Vomiting	18 (22.2)	18 (22.2)
Thrombocytopenia	15 (18.5)	15 (18.5)
Decreased appetite	14 (17.3)	14 (17.3)
Headache	13 (16.0)	13 (16.0)
Dysgeusia	12 (14.8)	12 (14.8)
Cough	11 (13.6)	11 (13.6)
Asthenia	10 (12.3)	10 (12.3)
Constipation	10 (12.3)	10 (12.3)
Arthralgia	9 (11.1)	9 (11.1)
Diarrhoea	8 (9.9)	8 (9.9)
Insomnia	8 (9.9)	8 (9.9)
Oedema peripheral	8 (9.9)	8 (9.9)
Urinary tract infection	7 (8.6)	7 (8.6)
Abdominal pain	7 (8.6)	7 (8.6)
Lymphopenia	6 (7.4)	6 (7.4)
Dizziness	6 (7.4)	6 (7.4)
Neutropenia	6 (7.4)	6 (7.4)
Nasopharyngitis	5 (6.2)	5 (6.2)
Aspartate aminotransferase increased	5 (6.2)	5 (6.2)
Back pain	5 (6.2)	5 (6.2)

Relationship assessed by the investigator. Percentages are based on the number of subjects treated in each group. Overall N = 81.

Table 14.3.6 Adverse Events Leading to Discontinuation

Table 14.3.6 Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=81)	Total (N=81)
Malignant neoplasm progression	2 (2.5)	2 (2.5)
Ascites	1 (1.2)	1 (1.2)
Aspartate aminotransferase increased	1 (1.2)	1 (1.2)
Intestinal obstruction	1 (1.2)	1 (1.2)
Nausea	1 (1.2)	1 (1.2)
Pneumonia	1 (1.2)	1 (1.2)
Sepsis	1 (1.2)	1 (1.2)

Percentages are based on N = 81.

Table 14.3.7 Deaths

Table 14.3.7 Deaths – Safety Analysis Set

Category, n (%)	Kelvaparib 300 mg (N=81)	Total (N=81)
Any death	12 (14.8)	12 (14.8)
Death within 30 days of last dose	5 (6.2)	5 (6.2)
Primary cause of death		
Disease progression	6 (7.4)	6 (7.4)
Adverse event	6 (7.4)	6 (7.4)
Other	2 (2.5)	2 (2.5)

Percentages are based on the number of subjects treated within each group.

Table 14.3.8 Laboratory Shift from Baseline to Worst Grade

Table 14.3.8 Shift from Baseline to Worst Post-baseline CTCAE Grade for Selected Laboratory Parameters – Safety Analysis Set

Laboratory parameter	Kelvaparib 300 mg: worst grade 3 n (%)	Kelvaparib 300 mg: worst grade 4 n (%)	Kelvaparib 300 mg: shift >= 2 grades n (%)	Total: worst grade 3 n (%)	Total: worst grade 4 n (%)	Total: shift >= 2 grades n (%)
Haemoglobin	12 (14.8)	3 (3.7)	27 (33.3)	12 (14.8)	3 (3.7)	27 (33.3)
Platelets	9 (11.1)	1 (1.2)	18 (22.2)	9 (11.1)	1 (1.2)	18 (22.2)
Neutrophils	4 (4.9)	0 (0.0)	14 (17.3)	4 (4.9)	0 (0.0)	14 (17.3)
Lymphocytes	6 (7.4)	1 (1.2)	17 (21.0)	6 (7.4)	1 (1.2)	17 (21.0)
CA-125	11 (13.6)	2 (2.5)	26 (32.1)	11 (13.6)	2 (2.5)	26 (32.1)
ALT	1 (1.2)	1 (1.2)	8 (9.9)	1 (1.2)	1 (1.2)	8 (9.9)
AST	0 (0.0)	3 (3.7)	10 (12.3)	0 (0.0)	3 (3.7)	10 (12.3)
Creatinine	4 (4.9)	0 (0.0)	9 (11.1)	4 (4.9)	0 (0.0)	9 (11.1)
Lactate dehydrogenase	9 (11.1)	0 (0.0)	13 (16.0)	9 (11.1)	0 (0.0)	13 (16.0)
Albumin	1 (1.2)	1 (1.2)	4 (4.9)	1 (1.2)	1 (1.2)	4 (4.9)

CTCAE = Common Terminology Criteria for Adverse Events, version 5.0. Percentages are based on the number of subjects treated in each group. Overall N = 81.

Table 14.3.9 Adverse Event Onset Timing

Table 14.3.9 Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events – Safety Analysis Set

Preferred term	Events, n	Median time to onset (days)	Resolved, n (%)	Median time to resolution (days)
Nausea	55	31.0	47 (85.5)	14.0
Fatigue	37	30.0	34 (91.9)	14.0
Anaemia	34	41.5	29 (85.3)	12.0
Vomiting	23	25.0	20 (87.0)	11.0
Constipation	21	47.0	19 (90.5)	6.0
Headache	20	37.0	18 (90.0)	7.0
Decreased appetite	19	12.0	17 (89.5)	10.0
Dysgeusia	18	30.0	16 (88.9)	7.0
Cough	17	11.0	17 (100.0)	12.0
Thrombocytopenia	17	23.0	12 (70.6)	12.0
Diarrhoea	14	9.5	13 (92.9)	7.0
Arthralgia	14	44.0	14 (100.0)	13.0
Neutropenia	14	37.0	10 (71.4)	13.0
Asthenia	13	41.0	13 (100.0)	7.0
Urinary tract infection	10	20.0	9 (90.0)	11.0
Insomnia	10	10.0	9 (90.0)	16.0
Abdominal pain	10	5.5	7 (70.0)	5.0
Oedema peripheral	10	35.0	10 (100.0)	11.5
Nasopharyngitis	10	18.0	10 (100.0)	5.0
Dizziness	9	31.0	8 (88.9)	2.0

Events are counted at the event level rather than the subject level. NE = not estimable.