Study KLV-7701 — Tables, Listings and Figures

Kelvaparib (KLV-3308) — Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

Table 14.3.5 Serious Treatment-emergent Adverse Events by Preferred Term – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Anaemia	35 (8.4)	25 (12.0)	60 (9.6)
Pneumonia	11 (2.6)	4 (1.9)	15 (2.4)
Thrombocytopenia	13 (3.1)	1 (0.5)	14 (2.2)
Malignant neoplasm progression	6 (1.4)	7 (3.3)	13 (2.1)
Sepsis	7 (1.7)	4 (1.9)	11 (1.8)
Neutropenia	4 (1.0)	4 (1.9)	8 (1.3)
Pneumonitis	4 (1.0)	3 (1.4)	7 (1.1)
Myelodysplastic syndrome	3 (0.7)	3 (1.4)	6 (1.0)
Fatigue	3 (0.7)	1 (0.5)	4 (0.6)
Intestinal obstruction	2 (0.5)	2 (1.0)	4 (0.6)
Nausea	3 (0.7)	1 (0.5)	4 (0.6)
Ascites	2 (0.5)	1 (0.5)	3 (0.5)
Leukopenia	1 (0.2)	2 (1.0)	3 (0.5)
Alanine aminotransferase increased	1 (0.2)	1 (0.5)	2 (0.3)
Dyspnoea	2 (0.5)	0 (0.0)	2 (0.3)
Hypertension	2 (0.5)	0 (0.0)	2 (0.3)
Pulmonary embolism	1 (0.2)	1 (0.5)	2 (0.3)
Aspartate aminotransferase increased	1 (0.2)	0 (0.0)	1 (0.2)
Decreased appetite	0 (0.0)	1 (0.5)	1 (0.2)
Diarrhoea	1 (0.2)	0 (0.0)	1 (0.2)
Dizziness	0 (0.0)	1 (0.5)	1 (0.2)
Lymphopenia	1 (0.2)	0 (0.0)	1 (0.2)
Pleural effusion	0 (0.0)	1 (0.5)	1 (0.2)
Urinary tract infection	1 (0.2)	0 (0.0)	1 (0.2)
Vomiting	1 (0.2)	0 (0.0)	1 (0.2)

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)

Subjects are counted once per preferred term. Percentages are based on N = 627.