TABLES, LISTINGS AND FIGURES

Study KLV-7701 — A Phase 3 Randomised, Placebo-controlled Study of Kelvaparib as Maintenance Therapy in Recurrent High-grade Serous Ovarian Carcinoma

Kelvaparib (KLV-3308)

Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

This pack contains the post-text tables, listings and figures supporting the clinical study report for study KLV-7701. All displays derive from the analysis datasets locked on 13 June 2022.

Contents

- Table 14.1.1 Demographic and Baseline Characteristics
- Table 14.1.2 Subject Disposition
- Table 14.1.3 Extent of Exposure
- Table 14.2.1 Best Overall Response
- Table 14.2.2 Time-to-Event Analyses
- Table 14.3.1 Overview of Adverse Events
- Table 14.3.2a Adverse Events by System Organ Class and Preferred Term
- Table 14.3.5 Serious Adverse Events
- Table 14.3.4a Treatment-Related Adverse Events
- Table 14.3.6 Adverse Events Leading to Discontinuation
- Table 14.3.7 Deaths
- Table 14.3.8 Laboratory Shift from Baseline to Worst Grade
- Table 14.3.9 Adverse Event Onset Timing

Table 14.1.1 Demographic and Baseline Characteristics

Table 14.1.1 Demographic and Baseline Characteristics – Safety Analysis Set

Characteristic	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Age (years)			
Mean (SD)	65.3 (9.00)	64.7 (9.21)	65.1 (9.07)
Median	65.0	65.0	65.0
Min, max	38, 88	42, 88	38, 88
>= 65 years, n (%)	225 (53.8)	108 (51.7)	333 (53.1)
Sex, n (%)			
Male	0 (0.0)	0 (0.0)	0 (0.0)
Female	418 (100.0)	209 (100.0)	627 (100.0)
Race, n (%)			
White	342 (81.8)	161 (77.0)	503 (80.2)
Asian	38 (9.1)	25 (12.0)	63 (10.0)
Black or African American	20 (4.8)	11 (5.3)	31 (4.9)
Other	18 (4.3)	12 (5.7)	30 (4.8)
ECOG performance status, n (%)			
0	193 (46.2)	103 (49.3)	296 (47.2)
1	184 (44.0)	89 (42.6)	273 (43.5)
2	41 (9.8)	17 (8.1)	58 (9.3)
BRCA mutation status, n (%)			
BRCA1 mutated	101 (24.2)	49 (23.4)	150 (23.9)
BRCA2 mutated	62 (14.8)	33 (15.8)	95 (15.2)
BRCA wild-type	255 (61.0)	127 (60.8)	382 (60.9)
Homologous recombination deficiency, n (%)	222 (53.1)	104 (49.8)	326 (52.0)
Platinum-sensitive disease, n (%)	265 (63.4)	135 (64.6)	400 (63.8)
Residual disease after cytoreductive surgery, n (%)	178 (42.6)	88 (42.1)	266 (42.4)
Number of prior lines of platinum-based therapy			
Median	2.0	2.0	2.0

Characteristic	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
>= 3, n (%)	122 (29.2)	61 (29.2)	183 (29.2)

SD = standard deviation; ECOG = Eastern Cooperative Oncology Group. Percentages are based on the number of subjects in the safety analysis set within each group.

Table 14.1.2 Subject Disposition

Table 14.1.2 Subject Disposition – Safety Analysis Set

Disposition	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Subjects treated, n	418	209	627
Treatment ongoing at data cut-off, n (%)	101 (24.2)	43 (20.6)	144 (23.0)
Discontinued study treatment, n (%)	317 (75.8)	166 (79.4)	483 (77.0)
Disease progression	37 (8.9)	41 (19.6)	78 (12.4)
Adverse event	25 (6.0)	12 (5.7)	37 (5.9)
Death	38 (9.1)	23 (11.0)	61 (9.7)
Withdrawal by subject	82 (19.6)	37 (17.7)	119 (19.0)
Physician decision	67 (16.0)	27 (12.9)	94 (15.0)
Lost to follow-up	39 (9.3)	12 (5.7)	51 (8.1)
Protocol deviation	29 (6.9)	14 (6.7)	43 (6.9)

Percentages are based on the number of subjects treated within each group.

Table 14.1.3 Extent of Exposure

Table 14.1.3 Extent of Exposure – Safety Analysis Set

Exposure parameter	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Duration of treatment (days)			
Median	122.0	116.0	119.0
Q1, Q3	70.2, 166.5	62.0, 172.0	68.0, 169.0
Min, max	8, 340	8, 353	8, 353
Cycles started, median	4.0	4.0	4.0
Relative dose intensity (%), median	86.8	87.8	86.8
Any dose reduction, n (%)	73 (17.5)	40 (19.1)	113 (18.0)
Any dose interruption, n (%)	122 (29.2)	59 (28.2)	181 (28.9)

Q1 = first quartile; Q3 = third quartile. Relative dose intensity is the total dose received divided by the total dose planned over the treatment period.

Table 14.2.1 Best Overall Response

Table 14.2.1 Best Overall Response by RECIST 1.1 – Full Analysis Set

Best overall response	Kelvaparib 300 mg (N=405)	Placebo (N=199)	Total (N=604)
Complete response (CR)	49 (12.1)	15 (7.5)	64 (10.6)
Partial response (PR)	201 (49.6)	64 (32.2)	265 (43.9)
Stable disease	112 (27.7)	76 (38.2)	188 (31.1)
Progressive disease	43 (10.6)	44 (22.1)	87 (14.4)
Not evaluable	0 (0.0)	0 (0.0)	0 (0.0)
Objective response rate, n (%)	250 (61.7)	79 (39.7)	329 (54.5)
95% CI	(56.8, 66.5)	(32.8, 46.9)	(50.4, 58.5)

CI = confidence interval. Assessed according to RECIST version 1.1. Confidence intervals are exact (Clopper-Pearson).

Table 14.2.2 Time-to-Event Analyses

Table 14.2.2 Kaplan-Meier Estimates of Time-to-event Endpoints – Full Analysis Set

Endpoint	Kelvaparib 300 mg (N=405)	Placebo (N=199)	Total (N=604)
Duration of response, median (months)	12.0	9.9	11.7
25th, 75th percentile	(5.8, 16.2)	(3.3, 13.7)	(5.3, 15.9)
Progression-free survival, median (months)	9.2	6.7	8.2
25th, 75th percentile	(3.6, 14.8)	(2.6, 11.3)	(3.1, 13.2)
Overall survival, median (months)	28.5	30.4	30.4
25th, 75th percentile	(19.5, NE)	(24.5, NE)	(19.6, NE)

NE = not estimable. Duration of response is analysed among responding subjects only.

Table 14.3.1 Overview of Adverse Events

Table 14.3.1 Overview of Treatment-emergent Adverse Events – Safety Analysis Set

Category, n (%)	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Any TEAE	417 (99.8)	209 (100.0)	626 (99.8)
Any treatment-related TEAE	411 (98.3)	204 (97.6)	615 (98.1)
Any grade >= 3 TEAE	190 (45.5)	108 (51.7)	298 (47.5)
Any serious TEAE	98 (23.4)	61 (29.2)	159 (25.4)
Any TEAE leading to dose reduction	72 (17.2)	39 (18.7)	111 (17.7)
Any TEAE leading to discontinuation	61 (14.6)	35 (16.7)	96 (15.3)
Any fatal TEAE	43 (10.3)	25 (12.0)	68 (10.8)

TEAE = treatment-emergent adverse event. A subject is counted once within each category.

Table 14.3.2a Adverse Events by System Organ Class and Preferred Term

Table 14.3.2a Treatment-emergent Adverse Events Occurring in at Least 10% of Subjects by System Organ Class and Preferred Term – Safety Analysis Set

System organ class / Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Gastrointestinal disorders	364 (87.1)	180 (86.1)	544 (86.8)
Nausea	257 (61.5)	136 (65.1)	393 (62.7)
Vomiting	113 (27.0)	50 (23.9)	163 (26.0)
Constipation	105 (25.1)	58 (27.8)	163 (26.0)
Diarrhoea	81 (19.4)	38 (18.2)	119 (19.0)
Abdominal pain	84 (20.1)	34 (16.3)	118 (18.8)
Dyspepsia	44 (10.5)	23 (11.0)	67 (10.7)
Blood and lymphatic system disorders	310 (74.2)	162 (77.5)	472 (75.3)
Anaemia	202 (48.3)	112 (53.6)	314 (50.1)
Thrombocytopenia	101 (24.2)	50 (23.9)	151 (24.1)
Neutropenia	83 (19.9)	43 (20.6)	126 (20.1)
Leukopenia	44 (10.5)	29 (13.9)	73 (11.6)
General disorders and administration site conditions	253 (60.5)	129 (61.7)	382 (60.9)
Fatigue	198 (47.4)	101 (48.3)	299 (47.7)
Asthenia	54 (12.9)	33 (15.8)	87 (13.9)
Nervous system disorders	185 (44.3)	94 (45.0)	279 (44.5)
Headache	85 (20.3)	42 (20.1)	127 (20.3)
Dysgeusia	67 (16.0)	48 (23.0)	115 (18.3)
Dizziness	69 (16.5)	29 (13.9)	98 (15.6)
Musculoskeletal and connective tissue disorders	123 (29.4)	68 (32.5)	191 (30.5)
Back pain	50 (12.0)	30 (14.4)	80 (12.8)
Arthralgia	48 (11.5)	26 (12.4)	74 (11.8)
Infections and infestations	118 (28.2)	62 (29.7)	180 (28.7)
Nasopharyngitis	55 (13.2)	22 (10.5)	77 (12.3)
Urinary tract infection	39 (9.3)	25 (12.0)	64 (10.2)

System organ class / Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Investigations	106 (25.4)	57 (27.3)	163 (26.0)
Metabolism and nutrition disorders	95 (22.7)	57 (27.3)	152 (24.2)
Decreased appetite	95 (22.7)	57 (27.3)	152 (24.2)
Respiratory, thoracic and mediastinal disorders	96 (23.0)	47 (22.5)	143 (22.8)
Cough	55 (13.2)	26 (12.4)	81 (12.9)
Psychiatric disorders	46 (11.0)	30 (14.4)	76 (12.1)
Insomnia	46 (11.0)	30 (14.4)	76 (12.1)

Subjects are counted once per preferred term and once per system organ class at their highest reported grade. MedDRA version 25.0.

Table 14.3.5 Serious Adverse Events

Table 14.3.5 Serious Treatment-emergent Adverse Events by Preferred Term – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Anaemia	35 (8.4)	25 (12.0)	60 (9.6)
Pneumonia	11 (2.6)	4 (1.9)	15 (2.4)
Thrombocytopenia	13 (3.1)	1 (0.5)	14 (2.2)
Malignant neoplasm progression	6 (1.4)	7 (3.3)	13 (2.1)
Sepsis	7 (1.7)	4 (1.9)	11 (1.8)
Neutropenia	4 (1.0)	4 (1.9)	8 (1.3)
Pneumonitis	4 (1.0)	3 (1.4)	7 (1.1)
Myelodysplastic syndrome	3 (0.7)	3 (1.4)	6 (1.0)
Fatigue	3 (0.7)	1 (0.5)	4 (0.6)
Intestinal obstruction	2 (0.5)	2 (1.0)	4 (0.6)
Nausea	3 (0.7)	1 (0.5)	4 (0.6)
Ascites	2 (0.5)	1 (0.5)	3 (0.5)
Leukopenia	1 (0.2)	2 (1.0)	3 (0.5)
Alanine aminotransferase increased	1 (0.2)	1 (0.5)	2 (0.3)
Dyspnoea	2 (0.5)	0 (0.0)	2 (0.3)
Hypertension	2 (0.5)	0 (0.0)	2 (0.3)
Pulmonary embolism	1 (0.2)	1 (0.5)	2 (0.3)
Aspartate aminotransferase increased	1 (0.2)	0 (0.0)	1 (0.2)
Decreased appetite	0 (0.0)	1 (0.5)	1 (0.2)
Diarrhoea	1 (0.2)	0 (0.0)	1 (0.2)
Dizziness	0 (0.0)	1 (0.5)	1 (0.2)
Lymphopenia	1 (0.2)	0 (0.0)	1 (0.2)
Pleural effusion	0 (0.0)	1 (0.5)	1 (0.2)
Urinary tract infection	1 (0.2)	0 (0.0)	1 (0.2)
Vomiting	1 (0.2)	0 (0.0)	1 (0.2)

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)

Subjects are counted once per preferred term. Percentages are based on N = 627.

Table 14.3.4a Treatment-Related Adverse Events

Table 14.3.4a Treatment-related Treatment-emergent Adverse Events Occurring in at Least 5% of Subjects – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Nausea	179 (42.8)	84 (40.2)	263 (41.9)
Anaemia	117 (28.0)	75 (35.9)	192 (30.6)
Fatigue	106 (25.4)	50 (23.9)	156 (24.9)
Vomiting	73 (17.5)	34 (16.3)	107 (17.1)
Constipation	69 (16.5)	34 (16.3)	103 (16.4)
Decreased appetite	64 (15.3)	34 (16.3)	98 (15.6)
Thrombocytopenia	57 (13.6)	30 (14.4)	87 (13.9)
Headache	56 (13.4)	28 (13.4)	84 (13.4)
Neutropenia	50 (12.0)	30 (14.4)	80 (12.8)
Diarrhoea	51 (12.2)	29 (13.9)	80 (12.8)
Abdominal pain	48 (11.5)	25 (12.0)	73 (11.6)
Dysgeusia	43 (10.3)	27 (12.9)	70 (11.2)
Dizziness	40 (9.6)	21 (10.0)	61 (9.7)
Asthenia	37 (8.9)	19 (9.1)	56 (8.9)
Nasopharyngitis	37 (8.9)	15 (7.2)	52 (8.3)
Cough	35 (8.4)	17 (8.1)	52 (8.3)
Back pain	31 (7.4)	19 (9.1)	50 (8.0)
Insomnia	29 (6.9)	17 (8.1)	46 (7.3)
Arthralgia	24 (5.7)	20 (9.6)	44 (7.0)
Leukopenia	25 (6.0)	18 (8.6)	43 (6.9)
Urinary tract infection	28 (6.7)	14 (6.7)	42 (6.7)
Dyspepsia	26 (6.2)	15 (7.2)	41 (6.5)
Rash	18 (4.3)	21 (10.0)	39 (6.2)
Alanine aminotransferase increased	21 (5.0)	13 (6.2)	34 (5.4)

Relationship assessed by the investigator. Percentages are based on the number of subjects treated in each group. Overall N = 627.

Table 14.3.6 Adverse Events Leading to Discontinuation

Table 14.3.6 Treatment-emergent Adverse Events Leading to Discontinuation of Study Treatment – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Anaemia	11 (2.6)	8 (3.8)	19 (3.0)
Malignant neoplasm progression	6 (1.4)	7 (3.3)	13 (2.1)
Sepsis	8 (1.9)	3 (1.4)	11 (1.8)
Neutropenia	5 (1.2)	3 (1.4)	8 (1.3)
Pneumonia	7 (1.7)	1 (0.5)	8 (1.3)
Thrombocytopenia	7 (1.7)	0 (0.0)	7 (1.1)
Intestinal obstruction	2 (0.5)	2 (1.0)	4 (0.6)
Ascites	2 (0.5)	1 (0.5)	3 (0.5)
Dyspnoea	3 (0.7)	0 (0.0)	3 (0.5)
Leukopenia	1 (0.2)	2 (1.0)	3 (0.5)
Myelodysplastic syndrome	1 (0.2)	2 (1.0)	3 (0.5)
Pneumonitis	1 (0.2)	2 (1.0)	3 (0.5)
Fatigue	2 (0.5)	0 (0.0)	2 (0.3)
Lymphopenia	2 (0.5)	0 (0.0)	2 (0.3)
Pulmonary embolism	1 (0.2)	1 (0.5)	2 (0.3)
Abdominal pain	1 (0.2)	0 (0.0)	1 (0.2)
Asthenia	1 (0.2)	0 (0.0)	1 (0.2)
Constipation	0 (0.0)	1 (0.5)	1 (0.2)
Decreased appetite	1 (0.2)	0 (0.0)	1 (0.2)
Dizziness	0 (0.0)	1 (0.5)	1 (0.2)
Nausea	1 (0.2)	0 (0.0)	1 (0.2)
Pleural effusion	0 (0.0)	1 (0.5)	1 (0.2)
Pyrexia	1 (0.2)	0 (0.0)	1 (0.2)

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)

Percentages are based on N = 627.

Table 14.3.7 Deaths

Table 14.3.7 Deaths – Safety Analysis Set

Category, n (%)	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Any death	95 (22.7)	44 (21.1)	139 (22.2)
Death within 30 days of last dose	38 (9.1)	23 (11.0)	61 (9.7)
Primary cause of death			
Disease progression	47 (11.2)	15 (7.2)	62 (9.9)
Adverse event	43 (10.3)	25 (12.0)	68 (10.8)
Other	5 (1.2)	4 (1.9)	9 (1.4)

Percentages are based on the number of subjects treated within each group.

Table 14.3.8 Laboratory Shift from Baseline to Worst Grade

Table 14.3.8 Shift from Baseline to Worst Post-baseline CTCAE Grade for Selected Laboratory Parameters – Safety Analysis Set

Laboratory parameter	Kelvaparib 300 mg: worst grade 3 n (%)	Kelvaparib 300 mg: worst grade 4 n (%)	Kelvaparib 300 mg: shift >= 2 grades n (%)	Placebo: worst grade 3 n (%)	Placebo: worst grade 4 n (%)	Placebo: shift >= 2 grades n (%)	Total: worst grade 3 n (%)	Total: worst grade 4 n (%)	Total: shift >= 2 grades n (%)
Haemoglobin	60 (14.4)	15 (3.6)	137 (32.8)	36 (17.2)	12 (5.7)	78 (37.3)	96 (15.3)	27 (4.3)	215 (34.3)
Platelets	30 (7.2)	7 (1.7)	80 (19.1)	21 (10.0)	2 (1.0)	36 (17.2)	51 (8.1)	9 (1.4)	116 (18.5)
Neutrophils	35 (8.4)	16 (3.8)	84 (20.1)	20 (9.6)	11 (5.3)	47 (22.5)	55 (8.8)	27 (4.3)	131 (20.9)
Lymphocytes	34 (8.1)	12 (2.9)	74 (17.7)	12 (5.7)	4 (1.9)	36 (17.2)	46 (7.3)	16 (2.6)	110 (17.5)
CA-125	55 (13.2)	12 (2.9)	136 (32.5)	35 (16.7)	10 (4.8)	67 (32.1)	90 (14.4)	22 (3.5)	203 (32.4)
ALT	23 (5.5)	3 (0.7)	50 (12.0)	11 (5.3)	1 (0.5)	19 (9.1)	34 (5.4)	4 (0.6)	69 (11.0)
AST	14 (3.3)	1 (0.2)	33 (7.9)	9 (4.3)	3 (1.4)	19 (9.1)	23 (3.7)	4 (0.6)	52 (8.3)
Creatinine	22 (5.3)	8 (1.9)	49 (11.7)	10 (4.8)	1 (0.5)	19 (9.1)	32 (5.1)	9 (1.4)	68 (10.8)
Lactate dehydrogenase	26 (6.2)	5 (1.2)	58 (13.9)	14 (6.7)	4 (1.9)	24 (11.5)	40 (6.4)	9 (1.4)	82 (13.1)
Albumin	14 (3.3)	4 (1.0)	27 (6.5)	5 (2.4)	6 (2.9)	23 (11.0)	19 (3.0)	10 (1.6)	50 (8.0)

CTCAE = Common Terminology Criteria for Adverse Events, version 5.0. Percentages are based on the number of subjects treated in each group. Overall N = 627.

Table 14.3.9 Adverse Event Onset Timing

Table 14.3.9 Time to Onset and Resolution of the Most Frequent Treatment-emergent Adverse Events – Safety Analysis Set

Preferred term	Events, n	Median time to onset (days)	Resolved, n (%)	Median time to resolution (days)
Nausea	393	32.0	330 (84.0)	11.0
Anaemia	317	29.0	265 (83.6)	10.0
Fatigue	299	32.0	239 (79.9)	10.0
Vomiting	163	34.0	130 (79.8)	11.0
Constipation	163	29.0	131 (80.4)	9.0
Decreased appetite	152	38.0	127 (83.6)	10.0
Thrombocytopenia	152	41.0	129 (84.9)	12.0
Neutropenia	127	36.0	98 (77.2)	9.5
Headache	127	25.0	105 (82.7)	11.0
Diarrhoea	119	37.0	97 (81.5)	11.0
Abdominal pain	118	25.5	102 (86.4)	11.5
Dysgeusia	115	24.0	92 (80.0)	10.0
Dizziness	98	23.5	76 (77.6)	12.0
Asthenia	87	33.0	74 (85.1)	10.0
Cough	81	33.0	65 (80.2)	10.0
Back pain	80	28.0	67 (83.8)	10.0
Nasopharyngitis	77	29.0	61 (79.2)	11.0
Insomnia	76	38.0	65 (85.5)	9.0
Arthralgia	74	36.0	65 (87.8)	8.0
Leukopenia	73	30.0	57 (78.1)	11.0

Events are counted at the event level rather than the subject level. NE = not estimable.