Study KLV-7701 — Tables, Listings and Figures

Kelvaparib (KLV-3308) — Marnholt Oncology GmbH

Indication: High-grade serous ovarian carcinoma

Table 14.3.4a Treatment-related Treatment-emergent Adverse Events Occurring in at Least 5% of Subjects – Safety Analysis Set

Preferred term	Kelvaparib 300 mg (N=418)	Placebo (N=209)	Total (N=627)
Nausea	179 (42.8)	84 (40.2)	263 (41.9)
Anaemia	117 (28.0)	75 (35.9)	192 (30.6)
Fatigue	106 (25.4)	50 (23.9)	156 (24.9)
Vomiting	73 (17.5)	34 (16.3)	107 (17.1)
Constipation	69 (16.5)	34 (16.3)	103 (16.4)
Decreased appetite	64 (15.3)	34 (16.3)	98 (15.6)
Thrombocytopenia	57 (13.6)	30 (14.4)	87 (13.9)
Headache	56 (13.4)	28 (13.4)	84 (13.4)
Neutropenia	50 (12.0)	30 (14.4)	80 (12.8)
Diarrhoea	51 (12.2)	29 (13.9)	80 (12.8)
Abdominal pain	48 (11.5)	25 (12.0)	73 (11.6)
Dysgeusia	43 (10.3)	27 (12.9)	70 (11.2)
Dizziness	40 (9.6)	21 (10.0)	61 (9.7)
Asthenia	37 (8.9)	19 (9.1)	56 (8.9)
Nasopharyngitis	37 (8.9)	15 (7.2)	52 (8.3)
Cough	35 (8.4)	17 (8.1)	52 (8.3)
Back pain	31 (7.4)	19 (9.1)	50 (8.0)
Insomnia	29 (6.9)	17 (8.1)	46 (7.3)
Arthralgia	24 (5.7)	20 (9.6)	44 (7.0)
Leukopenia	25 (6.0)	18 (8.6)	43 (6.9)
Urinary tract infection	28 (6.7)	14 (6.7)	42 (6.7)
Dyspepsia	26 (6.2)	15 (7.2)	41 (6.5)
Rash	18 (4.3)	21 (10.0)	39 (6.2)
Alanine aminotransferase increased	21 (5.0)	13 (6.2)	34 (5.4)

Relationship assessed by the investigator. Percentages are based on the number of subjects treated in each group. Overall N = 627.