
Children under 5 who are stunted (height-for-age) †
Men age 15+ who use any kind of tobacco
Women whose Body Mass Index is below normal †
Children under 5 who are underweight (weight-for-age) †
Women age 20-24 married before age 18
Women age 15+ who use any kind of tobacco
Women age 15-19 already mothers or pregnant
Pregnant women age 15-49 who are anaemic
Children under 5 who are wasted (weight-for-height)
All women age 15-49 who are anaemic
Total unmet need for family planning (currently married wo...
Children 6-59 months who are anaemic
Women age 15+ who consume alcohol
Sex ratio at birth (females per 1,000 males)
Men age 15+ who consume alcohol
Households with a member covered by health insurance/finan...
Children 24-35 months with a second dose of measles-contai...
Population with an improved drinking-water source †
Children age 5 attending pre-primary school
Children under 5 who are overweight (weight-for-height)
Children 12-23 months fully vaccinated (card or mother's r...
Children 12-23 months with 3 doses of penta or DPT vaccine
Currently married women (15-49) using a modern family plan...
Men 15+ with elevated blood pressure or taking medicine
Women 15+ with elevated blood pressure or taking medicine
Children under 5 whose birth was registered
Population using an improved sanitation facility †
Mothers who had at least 4 antenatal care visits †
Deaths in the last 3 years registered
Births attended by skilled health personnel †
Population living in households with electricity †
Women (15-49) who are literate
Institutional births
Households using clean fuel for cooking †
−1.0
−0.5
0.0
0.5
1.0
Spearman's ρ (95% CI)