8/8/2018 Gradient Checking | Coursera

≮ Back to Week 5

X Lessons

This Course: Machine Learning

Next

Gradient Checking

Gradient checking will assure that our backpropagation works as intended. We can approximate the derivative of our cost function with:

$$rac{\partial}{\partial\Theta}J(\Theta)pproxrac{J(\Theta+\epsilon)-J(\Theta-\epsilon)}{2\epsilon}$$

With multiple theta matrices, we can approximate the derivative **with respect to** Θ_i as follows:

$$rac{\partial}{\partial \Theta_j} J(\Theta) pprox rac{J(\Theta_1, \ldots, \Theta_j + \epsilon, \ldots, \Theta_n) - J(\Theta_1, \ldots, \Theta_j - \epsilon, \ldots, \Theta_n)}{2\epsilon}$$

A small value for ϵ (epsilon) such as $\epsilon=10^{-4}$, guarantees that the math works out properly. If the value for ϵ is too small, we can end up with numerical problems.

Hence, we are only adding or subtracting epsilon to the Θ_j matrix. In octave we can do it as follows:

```
epsilon = 1e-4;
for i = 1:n,
   thetaPlus = theta;
 thetaFlus(i) += epsilon;
thetaMinus = theta;
thetaMinus(i) -= epsilon;
gradApprox(i) = (J(thetaPlus) - J(thetaMinus))/(2*epsilon)
```

We previously saw how to calculate the deltaVector. So once we compute our gradApprox vector, we can check that gradApprox ≈ deltaVector.

Once you have verified **once** that your backpropagation algorithm is correct, you don't need to compute gradApprox again. The code to compute gradApprox can be very slow.

✓ Complete







8/8/2018 Gradient Checking | Coursera