

002 - Motivating Examples

EPIB 607 - FALL 2020

Sahir Rai Bhatnagar

Department of Epidemiology, Biostatistics, and Occupational Health
McGill University

`sahir.bhatnagar@mcgill.ca`

slides compiled on September 1, 2020



Early phase COVID-19 vaccine trial¹

Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial



Pedro M Folegatti*, Katie J Ewer*, Parvinder K Aley, Brian Angus, Stephan Becker, Sandra Belij-Rammerstorfer, Duncan Bellamy, Sagida Bibi, Mustapha Bittaye, Elizabeth A Clutterbuck, Christina Dold, Saul N Faust, Adam Finn, Amy L Flaxman, Bassam Hallis, Paul Heath, Daniel Jenkin, Rajeka Lazarus, Rebecca Makinson, Angela M Minassian, Katrina M Pollock, Maheshi Ramasamy, Hannah Robinson, Matthew Snape, Richard Tarrant, Meryn Voysey, Catherine Green*, Alexander D Douglas*, Adrian V S Hill*, Teresa Lambe*, Sarah C Gilbert*, Andrew J Pollard*, on behalf of the Oxford COVID Vaccine Trial Group†

Summary

Background The pandemic of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) might be curtailed by vaccination. We assessed the safety, reactogenicity, and immunogenicity of a viral vectored coronavirus vaccine that expresses the spike protein of SARS-CoV-2.

Methods We did a phase 1/2, single-blind, randomised controlled trial in five trial sites in the UK of a chimpanzee adenovirus-vectored vaccine (ChAdOx1 nCoV-19) expressing the SARS-CoV-2 spike protein compared with a meningococcal conjugate vaccine (MenACWY) as control. Healthy adults aged 18–55 years with no history of laboratory confirmed SARS-CoV-2 infection or of COVID-19-like symptoms were randomly assigned (1:1) to receive ChAdOx1 nCoV-19 at a dose of 5×10^{10} viral particles or MenACWY as a single intramuscular injection. A protocol amendment in two of the five sites allowed prophylactic paracetamol to be administered before vaccination. Ten participants assigned to a non-randomised, unblinded ChAdOx1 nCoV-19 prime-boost group received a two-dose schedule, with the booster vaccine administered 28 days after the first dose. Humoral responses at baseline and following vaccination were assessed using a standardised total IgG ELISA against trimeric SARS-CoV-2 spike protein, a multiplexed immunoassay, three live SARS-CoV-2 neutralisation assays (a 50% plaque reduction neutralisation assay [PRNT₅₀]; a microneutralisation assay [MNA₅₀, MNA₉₀, and MNA₅₀]; and Marburg VN), and a pseudovirus neutralisation assay. Cellular responses were assessed using an ex-vivo interferon- γ enzyme-linked immunospot assay. The co-primary outcomes are to assess efficacy, as measured by cases of symptomatic virologically confirmed COVID-19, and safety, as measured by the occurrence of serious adverse events. Analyses were done by group allocation in participants who received the vaccine. Safety was assessed over 28 days after

Published Online

July 20, 2020
[https://doi.org/10.1016/S0140-6736\(20\)31604-4](https://doi.org/10.1016/S0140-6736(20)31604-4)

See Online/Comment
[https://doi.org/10.1016/S0140-6736\(20\)31611-1](https://doi.org/10.1016/S0140-6736(20)31611-1)

*Contributed equally
†Members are listed in the appendix

The Jenner Institute
(P M Folegatti MSc, K J Ewer PhD, S Belij-Rammerstorfer PhD, D Bellamy MSc, M Bittaye PhD, A L Flaxman DPhil, D Jenkin MRCP, R Makinson MSc, A M Minassian DPhil, A D Douglas DPhil, Prof A V S Hill FMedSci, T Lambe PhD,


¹[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)31604-4/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31604-4/fulltext)

Phase 1/2 trial

- The focus in phase 1/2 trials is looking at what the vaccine does to the body and what the body does with the vaccine in *healthy* individuals
- Healthy adults aged 18-55 years with no history of laboratory confirmed SARS-CoV-2 infection or of COVID-19-like symptoms were randomly assigned (1:1) to receive **ChAdOx1 nCoV-19** at a dose of 5×10^{10} viral particles or **MenACWY** as a single intramuscular injection
- Between April 23 and May 21, 2020, 1077 participants were enrolled and assigned to receive either ChAdOx1 nCoV-19 (n=543) or MenACWY (n=534)
- Convalescent plasma is collected from someone who has recovered from a virus. When a person is infected with a virus, their body starts making antibodies to fight it. It is believed these antibodies could be the key ingredient for a treatment to help others with the same virus.
- Safety was assessed over 28 days after vaccination


test

```
plot(rnorm(10))
```



```
include_graphics2("http://www.biostat.mcgill.ca/hanley/statbook/OntarioCremation")
```

Figure 2: Weekly mortality rate from January 1, 2020 to May 31, 2020. The time series model is represented by the blue solid line and 95% confidence intervals in the shaded blue band.



```
ds=read.table("http://www.biostat.mcgill.ca/hanley/statbook/immunogenicityChAdO
str(ds)
```

```
## 'data.frame': ~I307 obs. of 2 variables:
## $ RefIndexCategory : Factor w/ 2 levels "Convalescent",...: 1 1 1
## $ IgGResponse.log10.ElisaUnits: num 2.56 2.74 2.79 3.32 3.15 2.35 2.72 2.9
```

```
tail(ds)
```

```
##           RefIndexCategory IgGResponse.log10.ElisaUnits
## 302 Day28PostChAdOx1 nCoV-19                1.99
## 303 Day28PostChAdOx1 nCoV-19                1.99
## 304 Day28PostChAdOx1 nCoV-19                2.42
## 305 Day28PostChAdOx1 nCoV-19                2.46
## 306 Day28PostChAdOx1 nCoV-19                2.42
## 307 Day28PostChAdOx1 nCoV-19                1.17
```

Session Info

```
R version 3.6.2 (2019-12-12)
Platform: x86_64-pc-linux-gnu (64-bit)
Running under: Pop!_OS 19.10

Matrix products: default
BLAS: /usr/lib/x86_64-linux-gnu/openblas/libblas.so.3
LAPACK: /usr/lib/x86_64-linux-gnu/libopenblas-r0.3.7.so

attached base packages:
[1] tools      stats      graphics  grDevices  utils      datasets  methods
[8] base

other attached packages:
[1] forcats_0.5.0    stringr_1.4.0    dplyr_1.0.2      purrr_0.3.4
[5] readr_1.3.1      tidyr_1.1.2      tibble_3.0.3     ggplot2_3.3.2.9000
[9] tidyverse_1.3.0  knitr_1.29

loaded via a namespace (and not attached):
[1] Rcpp_1.0.4.6    highr_0.8        cellranger_1.1.0 pillar_1.4.6
[5] compiler_3.6.2 dbplyr_1.4.2     jsonlite_1.7.0  lubridate_1.7.4
[9] evaluate_0.14   lifecycle_0.2.0 gtable_0.3.0    pkgconfig_2.0.3
[13] rlang_0.4.7     reprex_0.3.0    cli_2.0.2       rstudioapi_0.11
[17] DBI_1.1.0       haven_2.3.1     xfun_0.16       withr_2.2.0
[21] xml2_1.3.0      httr_1.4.1      fs_1.3.2        generics_0.0.2
[25] vctrs_0.3.4     hms_0.5.3       grid_3.6.2      tidyrselect_1.1.0
[29] glue_1.4.2      R6_2.4.1        fansi_0.4.1     readxl_1.3.1
[33] modelr_0.1.5    magrittr_1.5     backports_1.1.9 scales_1.1.1
[37] ellipsis_0.3.1  rvest_0.3.5     assertthat_0.2.1 colorspace_1.4-1
[41] stringi_1.4.6   munsell_0.5.0   broom_0.7.0     crayon_1.3.4
```