Comprehensive Encyclopedia of Indian Aquatic Pathologies and Health States for Integrated Farm Management

Aquaculture in the Indian subcontinent is a sophisticated balance between traditional ecological knowledge and modern intensive production systems. The sector, dominated by Indian Major Carps (IMC) and high-value crustaceans like *Macrobrachium rosenbergii*, faces significant headwinds due to disease-related mortalities, which can account for a 10% to 15% reduction in total production value.[1] The following report provides an encyclopedic profile of the most critical health states and diseases encountered in Indian freshwater and brackish water environments. This document is structured to support diagnostic AI assistants by providing high-density clinical data, regional terminology, and culturally relevant remediation strategies.

Bacterial Red Disease

1. Pathogen & Regional Terminology

Causative Agent: The primary pathogen associated with Bacterial Red Disease in the Indian context is *Aeromonas hydrophila*, a Gram-negative, facultative anaerobic bacterium.[2] However, recent molecular investigations in Indian rohu farms have identified co-infections involving *Aeromonas veronii*, *Aeromonas dhakensis*, and *Pseudomonas fluorescens*, which contribute to the severity of the hemorrhagic septicemia.[5]

Common Global Names: Motile Aeromonas Septicemia (MAS), Bacterial Hemorrhagic Septicemia, Red Pest.

- Regional Indian Terms: In the Hindi-speaking belt and parts of Odisha and West Bengal, farmers frequently refer to this condition as **Lal Bimari** or **Lal Rog** (Red Disease).[1] In the Telugu heartlands of Andhra Pradesh, it is commonly identified as **Yerrati Tegulu** (Red Spots).[7] In Bengali culture, it is often grouped under **Pakhana Pacha** (Fin Rot) when lesions are localized to the appendages.[2]

2. Visual Symptoms & Clinical Signs

External Lesions & Marks: The definitive visual marker for this disease is the appearance of distinct hemorrhagic patches—bright red, bloody spots—across the body surface, particularly at the base of the pectoral and pelvic fins, the operculum, and around the vent.[2] As the

infection progresses, these patches evolve into deep, circular ulcers with necrotic centers that may expose the underlying muscle tissue.[3] Scale erosion is common, where the scales lose their luster and fall off easily, leaving the skin vulnerable.[3]

Gill Appearance: Gills in infected fish often show significant discoloration, ranging from pale pink to a muddy gray-white.[2] Excessive mucus secretion is a hallmark, often leading to a "sliminess" that traps suspended silt, further impairing the fish's ability to extract oxygen from the water.[2]

Behavioral Changes: Farmers observe abnormal surfacing, where fish crowd the top layer of the pond or the water inlet, gasping for air even when dissolved oxygen levels are technically sufficient—a sign of impaired branchial function.[1] Other signs include erratic swimming, "shimmying," and a total loss of appetite (anorexia), with the fish eventually becoming lethargic and non-responsive to external stimuli.[4]

Internal Symptoms: Upon dissection, a characteristic finding is the accumulation of a clear, straw-colored, or slightly bloody fluid (ascites) in the abdominal cavity.[3] The liver and kidney often appear swollen, congested, and may exhibit pinpoint hemorrhages known as petechiae.[3]

3. Susceptible Indian Species

Primary Targets: The Indian Major Carps—Rohu (*Labeo rohita*), Catla (*Catla catla*), and Mrigal (*Cirrhinus mrigala*)—are the primary targets in semi-intensive pond systems.[2] Additionally, air-breathing fishes like *Clarias magur* and high-density culture species like Tilapia and Pangasius are increasingly showing susceptibility under stressed conditions.[8]

Life Stage Vulnerability: While all life stages are at risk, the most devastating impacts are seen in fingerlings and yearlings during the transition from nursery to grow-out ponds, where mortality rates can reach 50% to 80% if untreated.[2]

4. Environmental Triggers in Indian Aquaculture

Water Quality Parameters: The primary driver for an *Aeromonas* outbreak is poor water quality, specifically high unionized ammonia ($> 0.05 \text{ mg/L}$) and low dissolved oxygen (DO) levels below 4.0 ppm .[1] Sudden drops in pH caused by heavy rain or excessive organic loading can also trigger the pathogen's transition from a commensal to a virulent state.[1]

Seasonal/Weather Factors: The Indian monsoon cycle plays a critical role; the pre-monsoon heat followed by a sudden influx of rainwater drastically changes the pond’s temperature and pH profile, stressing the fish.[2] Outbreaks are also highly prevalent in the post-monsoon period (September to November) and during the summer heat when biological oxygen demand (BOD)

peaks.[2]

Pond Management Failures: Overstocking beyond the carrying capacity of the pond (typically exceeding 5,000–8,000 fingerlings/ha in traditional ponds) and the accumulation of excess feed and raw cow dung lead to high microbial loads.[1]

5. Prevention & Prophylaxis

Traditional/Local Indian Methods: A standard preventive practice among rural farmers is the application of agricultural lime (**Chuna**) at a rate of 200–250 kg per hectare to stabilize the pH and act as a mild disinfectant.[1] Prophylactic use of turmeric powder (**Haldi**) mixed with salt during seed transport or stocking is common, leveraging curcumin's known antibacterial properties.[16]

Modern Aquaculture Practices: Regular application of **CIFAX**, a specialized formulation developed by ICAR-CIFA, at a dosage of 1 liter per hectare-meter of water depth, effectively suppresses *Aeromonas* loads.[18] The use of water-based probiotics containing *Bacillus subtilis* is also recommended to improve the environmental microbial balance.[9]

6. Treatment & Remediation Strategy

Immediate Action Plan: On Day 1 of spotting red patches, farmers must stop all fertilization and reduce feeding by 50-70%. Immediate aeration should be increased using paddlewheel aerators or diesel-powered pumps to splash water.[1]

Chemical/Medical Treatments: A whole-pond treatment with Potassium Permanganate ($KMnO_4$) at a dose of 2–4 mg/L (or 2–4 kg per hectare-meter) is the standard disinfectant response.[9] For therapeutic feed intervention, Oxytetracycline administered at 75–80 mg per kg of fish weight for a duration of 10–12 days is highly effective.[4]

Post-Treatment Care: After chemical application, the pond should be rested for 72 hours without heavy feeding. A follow-up application of lime is essential to neutralize the acidity of the chemicals and restore the buffering capacity of the water.[9]

7. Commonly Asked Questions

Q: Why are my Rohu fish gasping at the surface and showing red spots on their bellies? A: This is likely Bacterial Red Disease (Lal Bimari). It is caused by *Aeromonas* bacteria that flourish when the water has too much organic waste or too little oxygen.[4]

Q: How can I tell if my pond has too much waste causing this disease? A: If the water color is a very dark, thick green or has a foul smell, and you see bubbles rising from the bottom, the

organic waste is too high, creating a breeding ground for bacteria.[1]

Q: Can I use turmeric and salt to cure the red disease in a large pond? A: While turmeric and salt are good for prevention during stocking, they are usually not strong enough to stop a mass outbreak in a large pond. You should use $KMnO_4$ or CIFAX for better results.[9]

Q: Is this red disease contagious to other ponds on my farm? A: Yes. The bacteria can travel through shared nets, water pipes, or even on the feet of birds. Always disinfect your nets in a solution of salt or $KMnO_4$ before moving between ponds.[1]

Q: What should I do if I see dead fish floating in the morning? A: Immediately remove and bury the dead fish away from the pond. Stop all feeding for two days and maximize aeration. Contact a local fisheries officer to confirm if an antibiotic like Oxytetracycline is needed.[1]

Aeromoniasis

1. Pathogen & Regional Terminology

Causative Agent: Primarily *Aeromonas hydrophila*, a Gram-negative rod-shaped bacterium.[2] It is often part of a complex involving *Aeromonas sobria* and *Aeromonas liquefaciens*.[2]

Common Global Names: Dropsy, Infectious Dropsy of Carps, Motile Aeromonas Septicemia (MAS).

- Regional Indian Terms: In various parts of India, the most descriptive term used is **Jalodar** (meaning water in the belly).[2] Farmers in Odisha and West Bengal often call it **Pet Phula** (Swollen Belly Disease).[2] In Telugu-speaking regions, it is sometimes referred to as **Kunda Potta** (Pot-belly).[6]

2. Visual Symptoms & Clinical Signs

External Lesions & Marks: The most striking visual indicator is a severely distended or bloated abdomen.[3] This is frequently accompanied by "pine-cone" scales, where the scales stand out from the body at a sharp angle due to the internal pressure of accumulated fluid.[3] Exophthalmia, or "pop-eye," where the eyes bulge abnormally from the sockets, is another common clinical sign.[2]

Gill Appearance: Gills may appear pale or exhibit bloody streaks. In chronic cases, there is significant erosion of the gill lamellae, reducing the surface area for respiration.[2]

Behavioral Changes: Infected fish exhibit a marked loss of balance, often swimming in circles or hanging vertically with the head up or down. They become extremely lethargic and are often

found resting near the pond banks where water is shallower.[3]

Internal Symptoms: Upon necropsy, the abdominal cavity is found to be engorged with a straw-colored, yellowish fluid.[3] Internal organs such as the liver, kidney, and spleen are typically enlarged (organomegaly) and may show signs of softening or necrosis.[3]

3. Susceptible Indian Species

Primary Targets: Common Carp (*Cyprinus carpio*) and Indian Major Carps (especially Rohu and Catla) are the most susceptible species in the Indian freshwater sector.[9] It is also a significant concern in the intensive culture of Tilapia and Pangasius.[9]

Life Stage Vulnerability: Broodfish and large adults are particularly vulnerable during the spawning season when their immune systems are naturally suppressed by reproductive stress.[2]

4. Environmental Triggers in Indian Aquaculture

Water Quality Parameters: Excessively high biological oxygen demand (BOD) and an accumulation of nitrogenous wastes like nitrite ($> 0.1 \text{ mg/L}$) are major precursors.[11] Chronic low dissolved oxygen ($< 3 \text{ ppm}$) significantly increases the risk of a dropsy outbreak.[9]

Seasonal/Weather Factors: Outbreaks are most frequent during the post-monsoon cooling phase and through the winter months when the fish's metabolic rate drops, making them less able to fight off bacterial invasion.[2]

Pond Management Failures: The primary failure is the lack of "bottom cleaning." The buildup of thick black sludge at the bottom of the pond (metabolic waste and uneaten feed) creates an anaerobic environment that fosters *Aeromonas* growth.[1]

5. Prevention & Prophylaxis

Traditional/Local Indian Methods: Using bundles of neem leaves (**Azadirachta indica**) submerged in the pond is a time-tested prophylactic against many bacterial infections.[12] Regular application of small doses of salt (around 50-100 kg per hectare) helps maintain the osmotic balance of the fish, preventing fluid accumulation.[12]

Modern Aquaculture Practices: Application of soil and water probiotics containing *Nitrosomonas* and *Nitrobacter* to reduce ammonia and nitrite levels.[21] Use of **Immunoboost-C** (developed by ICAR-CIFA) to improve the overall health and resistance of broodfish.[21]

6. Treatment & Remediation Strategy

Immediate Action Plan: Isolate heavily bloated fish immediately as they are a significant source of bacterial shedding. Stop all organic fertilization (cow dung, poultry waste) and reduce feed intake.[1]

Chemical/Medical Treatments: Disinfect the pond with a 1 ppm solution of Potassium Permanganate ($KMnO_4$).[15] For individual valuable fish or broodstock, a short-term bath in 5 ppm $KMnO_4$ for 2 minutes can be used.[22] Medicated feed containing Sulfadiazine (100 mg/kg fish) or Terramycin (Oxytetracycline, 75–80 mg/kg fish) should be administered for 10–12 days.[22]

Post-Treatment Care: Gradually reintroduce feeding using high-quality floating pellets. Apply CIFAX at 1 liter per hectare-meter to "flash" the environment and ensure the removal of residual bacterial pathogens.[18]

7. Commonly Asked Questions

Q: Why does my fish look like a pine cone with its scales sticking out? A: This is a classic sign of Dropsy (Aeromoniasis). The fish's internal organs are failing, and fluid is filling its belly, pushing the scales outward.[3]

Q: Is there a natural way to treat Pet Phula (swollen belly)? A: Prevention is better; use neem extract and salt regularly. However, once the belly is severely swollen, you must use $KMnO_4$ and potentially antibiotics in the feed to save the fish.[9]

Q: My fish has "Pop-Eye" along with a swollen belly. Is it the same disease? A: Yes, exophthalmia (pop-eye) is a common symptom of systemic bacterial infections like Aeromoniasis that affect the internal fluid pressure of the fish.[2]

Q: Can this disease spread if I don't remove the sick fish? A: Yes. Sick fish shed millions of bacteria into the water, which can infect healthy fish, especially if they are stressed by poor water quality.[1]

Q: How can I prevent the "black soil" at the bottom from causing this? A: You should periodically use a heavy chain to "rake" the bottom (bottom raking) to allow trapped gases to escape, or use soil probiotics to help break down the organic waste.[1]

Bacterial Gill Disease

1. Pathogen & Regional Terminology

Causative Agent: Primarily species of the genus *Flavobacterium*, most notably *Flavobacterium branchiophilum*. Other filamentous bacteria such as *Cytophaga* and *Flexibacter* are often isolated from the gill tissue in Indian ponds.[2]

Common Global Names: BGD, Bacterial Gill Rot, Environmental Gill Disease.

- Regional Indian Terms: In West Bengal and Odisha, farmers use the term **Gills Pacha** or **Galphada Pacha** (Gill Rot).[2] In the Telugu aquaculture belt, it is often called **Shanku Puta**.[6] In North India, it is sometimes described as **Gills ki Bimari**.

2. Visual Symptoms & Clinical Signs

External Lesions & Marks: Unlike skin-based bacterial infections, BGD often shows few external lesions initially. However, the opercula (gill covers) may appear flared or slightly swollen.[3] In advanced cases, the head region may exhibit a grayish-white film or secondary fungal growth.[2]

Gill Appearance: This is the primary diagnostic site. Healthy bright red gills are replaced by pale, pinkish-gray, or brownish-white tissue.[2] The gills appear "clogged" with excessive mucus and trapped silt or debris.[2] Microscopic examination often reveals "clubbing" of the gill lamellae, where the delicate filaments fuse together, drastically reducing the respiratory surface.[1]

Behavioral Changes: The most obvious sign is "piping"—fish gasping for air at the water surface, particularly near the pond inlets or aerators.[1] Fish exhibit extreme lethargy, do not respond to feeding, and often congregate in areas of highest water movement.[2]

Internal Symptoms: Internal organs are usually normal in the early stages, but as hypoxia (lack of oxygen) sets in, the liver may become pale and the blood may appear thin and watery.[3]

3. Susceptible Indian Species

Primary Targets: All freshwater species are susceptible, but the disease is particularly devastating to Indian Major Carps (Rohu and Catla) in high-density nursery and rearing ponds.[2]

Life Stage Vulnerability: Fry and fingerlings are the most at risk due to their smaller gill surface area and higher metabolic oxygen requirements.[2] Broodfish can also be affected during the hot summer months.

4. Environmental Triggers in Indian Aquaculture

Water Quality Parameters: High turbidity (suspended silt or clay particles) is a major mechanical

irritant for the gills.[27] Elevated unionized ammonia ($> 0.05 \text{ ppm}$) causes chemical "burning" of the gill tissue, which then becomes a site for bacterial colonization.[1]

Seasonal/Weather Factors: Most prevalent during the winter months (October to January) when water exchange is minimal, and during the peak summer (April to June) when high temperatures reduce oxygen solubility.[2]

Pond Management Failures: Overcrowding in nursery hapas or tanks is the leading cause.[8] Failure to use properly filtered or settled water in hatcheries often introduces the fine silt that triggers the condition.[1]

5. Prevention & Prophylaxis

Traditional/Local Indian Methods: Using "Banana Stem Pith" or juice as a natural coagulant to help settle suspended silt in the water column.[28] Farmers also use agricultural lime (**Chuna**) at 100–150 kg per hectare to help clear the water and stabilize pH.[1]

Modern Aquaculture Practices: Maintaining a dissolved oxygen level above 5.0 ppm through continuous aeration during the night and early morning.[1] Use of **Benzalkonium Chloride (BKC)** as a prophylactic at 0.5 ppm in hatchery tanks to keep bacterial counts low.[9]

6. Treatment & Remediation Strategy

Immediate Action Plan: On spotting gasping fish, immediately conduct a 30–50% water exchange with clean, oxygenated water. Stop all fertilization and reduce feed significantly.[1]

Chemical/Medical Treatments: Salt (NaCl) baths are very effective; the fish are dipped in a 3-5% salt solution for 5-10 minutes before being returned to the pond.[15] For pond-wide treatment, apply **Potassium Permanganate** at 2-4 ppm or use specialized antibiotics like Tetracycline (administered through the water in intensive systems).[6]

Post-Treatment Care: Apply **CIFAX** at 1 L/ha-m to "flash" the gill tissues and remove dead bacteria and mucus buildup.[19] Supplement the diet with Vitamin C (500 mg/kg feed) to help repair damaged gill membranes.[33]

7. Commonly Asked Questions

Q: My fish are gasping at the surface even though my aerator is running. Why? A: This is likely Bacterial Gill Disease. Even if there is oxygen in the water, the fish's gills are so clogged with mucus and bacteria that they cannot "breathe" properly.[2]

Q: How does mud in the water cause a bacterial disease? A: Fine mud or silt irritates the gills, causing the fish to produce extra mucus. Bacteria like *Flavobacterium* stick to this mucus and grow rapidly, eventually destroying the gill tissue.[1]

Q: Can I use salt to treat my whole pond for gill rot? A: It is hard to put enough salt in a large pond to kill the bacteria. It is much better to catch the sick fish and give them a "salt bath" in a bucket for 5 minutes before putting them back.[15]

Q: What is the best way to prevent gill rot in my nursery? A: Do not put too many fry in one tank or hapa. Make sure your water is clear and not muddy. Use a small dose of lime or BKC every week to keep the water clean.[1]

Q: Is CIFAX useful for gill problems? A: Yes. CIFAX is designed to "clean" the gills and skin by killing the bacteria that live in the mucus. It helps the fish breathe better almost immediately.[19]

Saprolegniasis

1. Pathogen & Regional Terminology

Causative Agent: Various species of the oomycete genus *Saprolegnia*, most commonly *Saprolegnia parasitica*. These are often called "water molds" and are opportunistic pathogens that colonize injured or stressed aquatic organisms.[2]

Common Global Names: Cotton Wool Disease, Winter Fungus, Water Mold Infection.

- Regional Indian Terms: In Andhra Pradesh and Telangana, it is known as **Rui Tegulu** or **Kapahi Tegulu** (Cottony Disease).[7] In Bengal and Odisha, farmers call it **Toon** or **Tula Rog** (Cotton disease).[2] It is also simply called **Fungus ki Bimari** in many Hindi-speaking regions.[1]

2. Visual Symptoms & Clinical Signs

External Lesions & Marks: The hallmark of this infection is the appearance of white, grayish, or brownish cotton-wool-like tufts on the skin, fins, mouth, or eyes.[2] These tufts are actually fungal hyphae growing out from the fish tissue.[35] If left untreated, these mats can cover large areas of the body, and the underlying skin becomes necrotic and ulcerated.[2]

Gill Appearance: The fungus can also invade the gills, appearing as white fuzzy patches that trap silt. This leads to rapid respiratory failure as the gill filaments are physically smothered and eventually destroyed.[2]

Behavioral Changes: Infected fish become very lethargic and often separate themselves from the school. They may be seen rubbing their bodies against pond walls, bamboo poles, or the

bottom to relieve the irritation caused by the fungal growth.[3]

Internal Symptoms: While primarily a superficial infection, in chronic cases, the fungus can penetrate deep into the muscle tissue and eventually reach the internal organs, leading to systemic death.[2]

3. Susceptible Indian Species

Primary Targets: Mrigal (*Cirrhinus mrigala*) is famously the most susceptible among the Indian Major Carps, followed by Rohu and Catla.[2] It is also a devastating problem for fish eggs in all freshwater hatcheries.[8]

Life Stage Vulnerability: Fish eggs are the most at risk; a single dead egg can serve as a nursery for the fungus, which then spreads and kills the entire batch.[8] Broodfish and large adults are susceptible after handling or spawning.[2]

4. Environmental Triggers in Indian Aquaculture

Water Quality Parameters: Accumulation of uneaten feed and dead organic matter provides the perfect food source for *Saprolegnia*.[2] Low dissolved oxygen and high ammonia weaken the fish's protective mucus layer, allowing the fungal spores to attach.[10]

Seasonal/Weather Factors: This is primarily a winter disease in India. It is most severe when water temperatures drop below $15^\circ\mathrm{C}$ to $20^\circ\mathrm{C}$.[2] The transition period from late monsoon to winter is a high-risk time for outbreaks.[2]

Pond Management Failures: The biggest failure is rough handling during netting or transport. Any injury, such as a lost scale or a skin scrape, is an immediate entry point for fungal spores.[1] Overstocking in cold weather also significantly increases the risk.[10]

5. Prevention & Prophylaxis

Traditional/Local Indian Methods: Using a salt bath (**Namak ka pani**) at 3% concentration is the most effective traditional preventive after handling or before stocking.[12] Many farmers also add small amounts of Potassium Permanganate ($\mathrm{KMnO_4}$) to their transport water.[22]

Modern Aquaculture Practices: Treatment of eggs with **Methylene Blue** or **Formalin** in the hatchery to prevent fungal colonization.[35] Use of **CIFAX** (1 L/ha-m) to keep the pond environment clear of fungal spores.[19] Ensuring that fish are handled with soft, wet nets and kept in the water as much as possible.[1]

6. Treatment & Remediation Strategy

Immediate Action Plan: Identify and remove the source of stress (e.g., stop rough handling, improve water quality). Remove heavily infected fish or dead eggs as they act as a source of spores.[35]

Chemical/Medical Treatments: Salt (NaCl) baths (3-4% solution) for 5–10 minutes are very effective for early skin infections.[15] For pond-wide disinfection, use Potassium Permanganate at 2–5 ppm.[22] Formalin can be used at 25 ppm (25 ml in 1,000 liters) for 1 hour, or as an indefinite bath at 12.5–25 mg/L in tanks.[36]

Post-Treatment Care: Gradually increase the water temperature if possible (in hatcheries). Feed a high-protein diet enriched with Vitamin C to speed up the healing of skin lesions.[33]

7. Commonly Asked Questions

Q: Why does my fish have white cotton-like stuff on its skin in the winter? A: This is Saprolegniasis (Winter Fungus). It happens when the water is cold and the fish's immune system is weak, especially if they were hurt by a net or a parasite.[2]

Q: My fish eggs in the hatchery are turning into a white fuzzy ball. What can I do? A: The fungus is eating the dead eggs and spreading to the live ones. You must remove the dead eggs immediately and use Methylene Blue or a light Formalin bath to protect the rest.[8]

Q: Is salt enough to cure a fish covered in fungus? A: If the infection is small, a 3% salt bath will work. But if the fungus is covering a lot of the body, you will need $\mathrm{KMnO_4}$ or Formalin to save the fish.[15]

Q: Can I catch fungus from my fish? A: No, *Saprolegnia* does not infect humans. However, you should not eat heavily infected fish as the meat will be poor quality and may have other bacteria.[35]

Q: How can I prevent fungus after I harvest some fish for the market? A: The fish remaining in the pond often get hurt by the nets during harvest. To prevent fungus, apply a dose of salt or CIFAX to the pond immediately after you finish harvesting.[1]

Healthy Fish (Optimal Health Baseline)

1. Physical Indicators & Baseline Parameters

Ideal Visual State: A healthy fish is a reflection of its environment. For Indian Major Carps (IMC)

and Scampi, the body should be streamlined and the skin must have a natural, lustrous shine.[1] Scales must be intact, laying flat against the body with no missing patches or discolored spots.[1] Eyes must be clear, bright, and alert—never cloudy, sunken, or bulging.[10] The mucus layer should be thin, translucent, and not excessively thick or "slimy".[10]

Gill Appearance: Gills are the most sensitive indicator of health. They must be a bright, vibrant cherry-red color.[1] The filaments should be clearly defined and free of mud, excess mucus, or white spots.[9] Any sign of pale pink or gray gills suggests a health problem.[2]

Behavioral State: Healthy fish show a vigorous feeding response, rushing to the surface or feed area when food is provided.[1] They should display active schooling behavior and a strong "flee response" when approached.[1] Swimming should be rhythmic and horizontal; "piping" at the surface or "flashing" against the bottom are signs of distress.[1]

Internal Indicators: In a healthy fish, the abdominal cavity is free of excess fluid. The liver is reddish-brown and firm, and the gut is usually full of food, indicating active metabolism.[3] The hepatopancreas in shrimp should be a dark brown or tan color, never pale or white.[38]

2. Physiological & Environmental Baselines

The following table summarizes the biochemical and environmental ranges that define a "Healthy State" for Indian aquaculture systems.

Parameter	Ideal Range (IMCs)	Significance
Blood Glucose	$18.00 -$	High levels indicate acute stress or poor diet.[42]
Total Blood Protein	$2.90 -$	Low levels suggest chronic illness or starvation.[42]
Blood Cholesterol	$202 -$	Indicator of healthy lipid metabolism.[42]
Dissolved Oxygen (DO)	$>$	Critical for respiration and growth.[1]
Water pH	$7.5 -$	Slightly alkaline is best for fish immunity.[1]

Water Temperature	$25^\circ C -$	Optimal range for Indian tropical species.[1]
Ammonia (NH_3)	$<$	Levels above this cause tissue damage.[9]
Nitrite (NO_2)	$<$	Prevents "brown blood" disease.[11]

3. Susceptible Indian Species (Welfare Focus)

Primary Indicators for IMCs: Rohu, Catla, and Mrigal should achieve a weight of 800g to 1kg within 10–12 months under healthy conditions.[15] The modern G3 Rohu strain can even reach this size in 6–8 months.[44]

Primary Indicators for Shrimp: For *Penaeus vannamei* and *Macrobrachium rosenbergii*, the gut should be visibly full (a dark line along the back) and the shell should be hard and free of black spots.[38]

4. Environmental Triggers for Maintaining Health

Water Quality: Regular morning testing of DO and pH is essential. A sudden drop in pH below 7.0 is often an early warning that the pond is becoming "sour" from too much waste.[1]

Management: Maintaining proper stocking density—typically 5,000–8,000 fingerlings per hectare in carps—prevents the chronic stress that leads to disease outbreaks.[1]

Seasonal Awareness: During the Indian summer, increase aeration to counteract the low oxygen solubility of hot water. During the winter, reduce feeding as the fish's metabolic rate slows down.[2]

5. Prevention & Prophylaxis (Baseline Care)

Traditional Wisdom: Monthly application of agricultural lime (**Chuna**) at 50-100 kg per hectare and occasional salt applications help maintain a stable environment.[1]

Modern Science: Regular water exchange (10-20% every month) and the use of high-quality, floating pellets with at least 25% protein ensure consistent growth and high immunity.[9]

6. Treatment & Remediation Strategy (For Stress)

Immediate Action: If fish look "dull" or stop eating, check water quality immediately. Increase

aeration and perform a partial water exchange.

Chemical Aid: A prophylactic dose of CIFAX (1 L/ha-m) can be used to "reset" the pond environment if a minor stress event (like a sudden heavy rain) occurs.[19]

Post-Stress Care: Add Vitamin C and B-complex to the feed for 7 days to help the fish recover their strength.[33]

7. Commonly Asked Questions

Q: How can I tell if my fish are healthy without a laboratory? A: Healthy fish are "greedy." If they rush to the surface to eat when you throw feed, they are likely healthy. Also, their skin should be shiny and their gills bright red.[1]

Q: What is the best color for pond water? A: A light brownish-green color is ideal. This shows you have a good balance of "plankton" (natural food) and enough oxygen.[1]

Q: My fish look healthy but they are not growing. Why? A: You may have too many fish in the pond (overstocking) or the feed quality is poor. Check your stocking density and try a higher-protein feed.[1]

Q: Is it normal for fish to jump occasionally? A: Yes, but if they jump many times or rub themselves against the bottom, it usually means they have parasites or the oxygen is low.[10]

Q: How often should I check my pond's water? A: You should check the oxygen and pH every morning. If the water smells bad, check the ammonia immediately.[1]

Parasitic Disease (Argulosis & Lernaeasis)

1. Pathogen & Regional Terminology

Causative Agent: Primarily species of the crustacean genus *Argulus* (Fish Lice), such as *Argulus siamensis* and *Argulus bengalensis*, and *Lernaea cyprinacea* (Anchor Worm).[2]

Common Global Names: Argulosis, Lernaeasis, Fish Lice, Anchor Worm.

- Regional Indian Terms: In Andhra Pradesh and Telangana, *Argulus* is called **Shanku Jalaga**.[7] In Bengal and Odisha, it is simply **Macher Uun** or **Uun Rog**.[13] Anchor worms are often called **Chuchia** or **Kanta** (meaning thorn or needle) because of how they stick out from the fish.[15]

2. Visual Symptoms & Clinical Signs

External Lesions & Marks:

- **Argulus:** These appear as tiny, flat, disc-like parasites (about 0.5 to 1 cm wide) that move rapidly across the skin and fins.[9] The attachment sites often show red, irritated "bite" marks or small hemorrhages.[13]
- **Lernaea:** These look like white or grayish "threads" sticking out of the fish's body.[15] The point of entry is usually a red, inflamed ulcer or sore, often near the base of the fins or on the head.[2]

Gill Appearance: Parasites frequently attach inside the gill covers (opercula), causing excessive mucus production and a pale, irritated appearance of the gill filaments.[2]

Behavioral Changes: "Flashing" or "Scratching" is the most common sign, where fish rub their bodies against the pond bottom, bamboo poles, or the banks to dislodge the parasites.[1] Affected fish may jump repeatedly out of the water (especially Catla) and exhibit erratic swimming behavior.[12]

Internal Symptoms: Generally few internal signs, but chronic infestation leads to severe anemia (thin, watery blood) and secondary bacterial infections that can cause organ swelling.[2]

3. Susceptible Indian Species

Primary Targets: All Indian Major Carps are affected, but Rohu (*Labeo rohita*) is particularly susceptible to *Argulus*.[2] Catla and Mrigal are also common hosts. In the ornamental sector, Goldfish and Koi are highly vulnerable.[15]

Life Stage Vulnerability: Broodfish are most susceptible due to their slower movement and the physiological stress of the breeding season.[13] However, heavy infestations can kill small fingerlings through blood loss and stress.[9]

4. Environmental Triggers in Indian Aquaculture

Water Quality Parameters: High organic load and low water exchange facilitate the rapid breeding of these parasites.[13] Excessive nutrients lead to "thick" water where parasites can easily find hosts.

Seasonal/Weather Factors: Infestations are most severe during the rainy season (June to September) and the transition into winter.[2] June, July, and August are the peak months for *Argulus* in most of India.[13]

Pond Management Failures: The use of "bamboo poles" as support for feeding or markers is a

major risk, as *Argulus* lay their eggs on these surfaces.[1] Failure to disinfect nets and moving equipment from infested ponds are also common errors.[1]

5. Prevention & Prophylaxis

Traditional/Local Indian Methods: A common traditional practice is placing bamboo poles in the pond to serve as "egg traps." These poles are removed every 10–12 days and dried in the sun for 24 hours to kill the parasite eggs before being put back.[1] Using bundles of neem leaves submerged near the water inlet also helps repel parasites.[12]

Modern Aquaculture Practices: Application of **Cypermethrin** at very low doses (around 0.94–2.84 $\mu g/L$) as a prophylactic in broodfish ponds.[47] Regular sampling of fish (every 15 days during peak season) to check for early signs of lice.[1]

6. Treatment & Remediation Strategy

Immediate Action Plan: On spotting lice, remove any wooden poles or aquatic weeds from the pond as these are egg-laying sites. Increase water exchange if possible.[1]

Chemical/Medical Treatments:

- **For Argulus:** Apply **Malathion (50 EC)** at a rate of 1 liter per acre-meter (for carps) or use a combined treatment of Cypermethrin (2.84 $\mu g/L$ in water) and **Emamectin benzoate** (161 $\mu g/kg$ in feed) for 7 days for complete eradication.[15]
- **For Lernaea:** Salt baths (1-2%) are effective for killing the free-swimming stages. Potassium Permanganate ($KMnO_4$) at 2–5 ppm is also used to treat the secondary sores.[15]

Post-Treatment Care: Apply **CIFAX** at 1 L/ha-m to heal the tiny bite wounds and prevent secondary bacterial infections.[19]

7. Commonly Asked Questions

Q: Why are my fish jumping out of the water like they are being chased? A: This is usually a sign of Fish Lice (*Argulus*). The lice bite the fish, causing sharp pain and irritation, making them jump to try and shake the parasites off.[12]

Q: I see tiny green discs moving on my fish's skin. What are they? A: Those are *Argulus* (Fish Lice). They are blood-sucking parasites that make the fish weak and stop them from growing.[9]

Q: Can I use salt to kill anchor worms? A: Salt baths help kill the young worms, but the adult

"anchor" stage is very tough. You may need to manually remove them from valuable fish or use specialized chemicals for the whole pond.[15]

Q: How do I stop lice from coming back every year? A: Clean your pond bottom during the summer. Do not let wild fish enter. If you use bamboo poles, take them out and dry them in the sun every 10 days to kill the eggs.[1]

Q: Is Malathion safe for my fish? A: Malathion is used in India for lice, but you must be very careful with the dose (1 liter per acre). Using too much can kill the fish. Always ask an expert before using it.[15]

White Tail Disease

1. Pathogen & Regional Terminology

Causative Agent: A complex of two RNA viruses: *Macrobrachium rosenbergii* nodavirus (MrNV) and its associated Extra Small Virus (XSV). MrNV is the primary pathogen belonging to the Family Nodaviridae.[50]

Common Global Names: WTD, White Muscle Disease (WMD).

- Regional Indian Terms: In the scampi-farming hubs of Andhra Pradesh and Tamil Nadu, it is widely known as **Safed Ponch** (White Tail) or **Milky Tail Disease**.[54] Farmers also use the terms **Swollen Head Syndrome** or **Branchiostegite Blister Disease (BBD)** to describe the symptoms seen in adult prawns.[54]

2. Visual Symptoms & Clinical Signs

External Lesions & Marks: The definitive sign is a distinct, milky-white, opaque appearance of the abdominal muscles (the tail).[50] This opaqueness typically begins at the telson (the very end of the tail) and spreads forward toward the head.[50] In severe cases, the tail and uropods may begin to rot or degenerate.[50]

Behavioral Changes: Infected post-larvae (PL) become extremely lethargic and stop feeding (anorexia).[50] Farmers often notice "mica-like flakes" floating in the tanks, which are actually abnormal, brittle moults from infected prawns.[50]

Internal Symptoms: In adults, which often act as carriers, the cephalothorax (head region) may appear swollen or double its normal size.[54] Dissection of the head reveals water-filled sacs (blisters) above the hepatopancreas.[54] Microscopically, there is severe Zenker's necrosis—a

dissolution of the muscle fibers—in the abdominal tissue.[50]

3. Susceptible Indian Species

Primary Targets: The Giant Freshwater Prawn (*Macrobrachium rosenbergii*), known as "Scampi," is the primary host.[51] Some marine shrimp species like *Penaeus indicus* and *P. monodon* can act as asymptomatic carriers.[50]

Life Stage Vulnerability: Larvae, post-larvae (PL), and early juveniles are highly susceptible, with mortality rates reaching 95% to 100% within five days of the first signs.[50] Adults are resistant to mortality but remain lifelong carriers.[50]

4. Environmental Triggers in Indian Aquaculture

Transmission Mechanisms: The virus is transmitted both vertically (from the mother to the eggs) and horizontally (through the water or by eating infected prawns).[50]

Vectors: Various aquatic insects (like backswimmers and water boatmen) and *Artemia* are known vectors that can bring the virus into a hatchery or pond system.[50]

Water Quality: High stocking densities and temperature fluctuations between $28^\circ\mathrm{C}$ and $32^\circ\mathrm{C}$ create the stress necessary for a viral breakout in hatcheries.[40]

Regional Context: The disease is a major threat in the Nellore district of Andhra Pradesh and the coastal regions near Chennai, where scampi hatcheries are concentrated.[54]

5. Prevention & Prophylaxis

Traditional/Local Indian Methods: There are no traditional herbal cures for this viral disease. Prevention relies entirely on biosecurity. Farmers are advised to avoid using wild-caught broodstock and to only buy seed from hatcheries that provide a "PCR Negative" certificate.[1]

Modern Aquaculture Practices: Mandatory RT-PCR or **Immunodot Assay** screening of all broodstock and PL batches.[54] Strict biosecurity, including the use of bird nets and insect screens to keep out vectors.[50] Heat treatment of water (to $65^\circ\mathrm{C}$ for 2 hours) or chlorination is used in hatcheries to kill the virus.[52]

6. Treatment & Remediation Strategy

Immediate Action Plan: There is no known treatment for White Tail Disease once it starts.[52] If an outbreak is confirmed, the entire stock in the affected tank or pond must be culled and buried

away from the water source to prevent the virus from spreading to neighboring farms.[54]

Chemical/Medical Treatments: No antibiotics or chemicals are effective against the MrNV virus. However, applying **CIFAX** can help reduce secondary bacterial infections in surviving carrier adults.[19]

Post-Treatment Care: All tanks, pipes, and nets must be thoroughly disinfected with high-strength Chlorine (100 ppm) and left to dry in the sun for several days before a new batch is started.[50]

7. Commonly Asked Questions

Q: Why are my baby prawns turning white like milk and dying? A: This is White Tail Disease (WTD), a very deadly virus. It can kill your whole stock in just 4-5 days.[50]

Q: Can I use medicine to cure the white tail? A: No. There is no medicine for this virus. You must focus on prevention by buying "PCR-Tested" seed and keeping the pond very clean.[52]

Q: How did the virus get into my pond? A: It could have come from the mother prawn (through the eggs), from water insects, or from contaminated water. This is why testing the parents is so important.[50]

Q: My adult prawns have "water blisters" on their heads. Is this related? A: Yes, that is often a sign that adult prawns are carrying the White Tail virus. They might not die, but they will infect all the babies.[54]

Q: Is Scampi farming in Andhra Pradesh safe from this? A: It is a major risk in Nellore and Chennai. Always ask for a "Disease-Free" certificate from the hatchery before you buy your seed.[54]

Works cited

1. Disease Management | PDF - Scribd, accessed March 8, 2026, https://www.scribd.com/document/951239550/Disease-Management
2. Present Status of Fish Disease Management in Freshwater ..., accessed March 8, 2026, https://www.heraldopenaccess.us/openaccess/present-status-of-fish-disease-management-in-freshwater-aquaculture-in-india-state-of-the-art-review
3. An Overview of Fish Disease Diagnosis and Treatment in Aquaculture in Bangladesh - MDPI, accessed March 8, 2026, https://www.mdpi.com/2673-9496/5/4/18
4. How to prevent red disease in fish farming | fish farming telugu - YouTube, accessed March 8, 2026, https://www.youtube.com/watch?v=IruHkFPP6aM
5. First report of natural co-infections and protein profile of Indian Major Carp,

Labeo rohita, accessed March 8, 2026, https://pubmed.ncbi.nlm.nih.gov/40675503/
6. Tataku tegulu - Fish diseases In telugu - YouTube, accessed March 8, 2026, https://www.youtube.com/watch?v=qoSdPW1KLEs
7. Disease management in fish farming | Red disease | - YouTube, accessed March 8, 2026, https://www.youtube.com/watch?v=71tZQBaPI5A
8. Common Diseases of Ornamental Fishes: Diagnosis, Treatment, and Preventive Management - CIFE, accessed March 8, 2026, https://www.cife.edu.in/sites/default/files/Common%20Diseases%20of%20Ornamental%20Fishes%20Diagnosis%2C%20Treatment%2C%20and%20Preventive%20Management.pdf
9. Common Fish Diseases and Their Symptoms: A Comprehensive Guide for Fish Farmers, accessed March 8, 2026, https://fishvigyan.com/common-fish-diseases-and-their-symptoms-a-comprehensive-guide-for-fish-farmers
10. Common Fish Disease Symptoms and Treatments - Aquarium Industries, accessed March 8, 2026, https://www.aquariumindustries.com.au/wp-content/uploads/2015/04/Common-Fish-Disease-Symptoms-Poster.pdf
11. Optimizing Shrimp Culture Through Environmental Monitoring: Effects of Water Quality and Metal Ion Profile on Whiteleg Shrimp (Litopenaeus vannamei) Performance in a Semi-Intensive Culture Pond - MDPI, accessed March 8, 2026, https://www.mdpi.com/2073-4441/17/19/2818
12. Health Management of Rohu, Catla, Mrigal in Traditional & Modern Indian Fish Farming, accessed March 8, 2026, https://fishingsagaonlinein.com/health-management-of-rohu-catla-mrigal-in-traditional-modern-indian-fish-farming/
13. ARGULUS OF INDIAN MAJOR CARPS IN SELECTED FISH FARMS OF MYMENSINGH - CABI Digital Library, accessed March 8, 2026, https://www.cabidigitallibrary.org/doi/pdf/10.5555/20173099447
14. (PDF) ICAR-CIFA Training Manual No. 93 Farmers' Meet and Awareness Programme on Scientific Freshwater Aquaculture Development in Manipur (Under NEH Programme of ICAR-CIFA) Training Manual Quality Fish Production and Health Management in Aquaculture - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/330881188_ICAR-CIFA_Training_Manual_No_93_Farmers'_Meet_and_Awareness_Programme_on_Scientific_Freshwater_Aquaculture_Development_in_Manipur_Under_NEH_Programme_of_ICAR-CIFA_Training_Manual_Quality_Fish_Producti
15. Rohu | Fish Information - Apni Kheti, accessed March 8, 2026, https://www.apnikheti.com/en/pn/livestock/freshwater-aquaculture/rohu
16. The Potential of Herbal Plants in the Prevention and Treatment of Fish Diseases: A Review - Journal UNRAM, accessed March 8, 2026, https://journal.unram.ac.id/index.php/jfh/article/download/5932/3288
17. Efficacy of medicinal plants on microbial fish pathogens - This document is

discoverable and free to researchers across the globe due to the work of AgEcon Search. Help ensure our sustain, accessed March 8, 2026, https://ageconsearch.umn.edu/record/276672/files/bangladesh-ag-uni-0282.pdf
18. CIFAX 500 ml Fish Medicine - Fish Health Management (Pack of 2) for Pond Water Quality Management - Toolsvilla, accessed March 8, 2026, https://www.toolsvilla.com/cifax-fish-medicine-fish-health-and-growth
19. Atc Cifax Fish Medicine, Packaging Type: Jar, Pack Size: 100 Ml To 5000 Ml - IndiaMART, accessed March 8, 2026, https://www.indiamart.com/proddetail/cifax-fish-medicine-9403641473.html
20. cifax - ICAR-CIFA – Central Institute of Freshwater Aquaculture, accessed March 8, 2026, https://cifa.nic.in/main-content/test-1/
21. Fish Health Management - ICAR-CIFA – Central Institute of Freshwater Aquaculture, accessed March 8, 2026, https://cifa.nic.in/fish-health-management/
22. FAO-Catla catla - FAO.org, accessed March 8, 2026, https://www.fao.org/fishery/docs/DOCUMENT/aquaculture/CulturedSpecies/file/en/en_catla.htm
23. Evaluation of Pacific White Shrimp (Litopenaeus vannamei) Health during a Superintensive Aquaculture Growout Using NMR-Based Metabolomics | PLOS One - Research journals, accessed March 8, 2026, https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0059521
24. Utilizing Medicinal Plants for Disease Treatment in Aquaculture An Approach to Improve Fish Health.cdr - MARKHOR (The Journal of Zoology), accessed March 8, 2026, https://markhorjournal.com/index.php/mjz/article/download/119/98/606
25. Therapeutic effects of Neem (Azadirachta indica) leaf extract on fin and tail rot diseased fish Anabas testudineus (Bloch, accessed March 8, 2026, https://ijam.co.in/index.php/ijam/article/download/5237/1473/11390
26. Use of aquadrugs, chemicals and biological products in freshwater aquaculture systems of West Bengal, India - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/358797522_Use_of_aquadrugs_chemicals_and_biological_products_in_freshwater_aquaculture_systems_of_West_Bengal_India
27. Removal of Turbidity of Water by Banana Peel Using Adsorption Technology - JOCPR, accessed March 8, 2026, https://www.jocpr.com/articles/removal-of-turbidity-of-water-by-banana-peel-using-adsorption-technology.pdf
28. Potential Application of Banana Shoot System as Coagulant for Turbidity Removal - IOE Graduate Conference, accessed March 8, 2026, https://conference.ioe.edu.np/publications/ioegc14/IOEGC-14-037-C1-3-368.pdf
29. Evaluation of plant-based natural coagulants for municipal wastewater treatment - PMC, accessed March 8, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5767163/
30. (PDF) RESPONSE SURFACE OPTIMIZATION OF TREATED TURBID WATER USING BANANA STEM JUICE. - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/307168086_RESPONSE_SURFACE_OPTIMIZATION_OF_TREATED_TURBID_WATER_USING_BANANA_STEM_JUICE

31. 15. fish disease prevention and treatment - FAO.org, accessed March 8, 2026, https://www.fao.org/fishery/docs/CDrom/FAO_Training/FAO_Training/General/x6709e/x6709e15.htm
32. CIFAX with Vigour & Strength, accessed March 8, 2026, https://4.imimg.com/data4/XV/EV/MY-4946313/aquaculture-cifax.pdf
33. A Guide to Preventing and Treating Common Fish Diseases - Bahtera Adi Jaya, accessed March 8, 2026, https://bahteraadijaya.com/en/blogs/a-guide-to-preventing-and-treating-common-fish-diseases
34. Saprolegniasis - Aquaculture, Fisheries, & Pond Management, accessed March 8, 2026, https://fisheries.tamu.edu/2020/01/24/saprolegniasis/
35. Saprolegniasis in Fish | Tree of Life Exotic Pet Medical Center, accessed March 8, 2026, https://treeoflifeexotics.vet/education-resource-center/for-clients/fish/saprolegniasis-in-fish
36. VM77/VM061: Use of Formalin to Control Fish Parasites - Ask IFAS, accessed March 8, 2026, https://ask.ifas.ufl.edu/publication/VM061
37. Effects of formalin on water quality and parasitic monogeneans on silver perch (Bidyanus bidyanus Mitchell) in earthen ponds | Request PDF - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/230381919_Effects_of_formalin_on_water_quality_and_parasitic_monogeneans_on_silver_perch_Bidyanus_bidyanus_Mitchell_in_earthen_ponds
38. Appearance of gut and hepatopancreas of healthy shrimp and those... - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/figure/Appearance-of-gut-and-hepatopancreas-of-healthy-shrimp-and-those-indicated-for-WFS-after_fig1_390952580
39. Behavioural indicators of welfare in farmed fish - PMC - NIH, accessed March 8, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3276765/
40. Hepatopancreas colors predict survival of shrimp to EMS - Responsible Seafood Advocate, accessed March 8, 2026, https://www.globalseafood.org/advocate/hepatopancreas-colors-predict-survival-of-shrimp-to-ems/
41. Bacterial communities and signatures in the stomach and intestine of juvenile Penaeus (litopenaeus) vannamei shrimp affected by acute hepatopancreatic necrosis disease - PMC, accessed March 8, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC11239581/
42. Serum Biochemical Parameters of Four Fresh Water Indian Carps from a Local Aquatic Body - ARC Journals, accessed March 8, 2026, https://www.arcjournals.org/pdfs/ijisabf/v2-i3/3.pdf
43. Bacterial Community Associated with Healthy and Diseased Pacific White Shrimp (Litopenaeus vannamei) Larvae and Rearing Water across Different Growth Stages - PMC, accessed March 8, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC5513922/
44. (PDF) Analysing farmers' perceptions of genetically improved G3 Rohu (Labeo

rohita) fish culture in Mymensingh District - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/publication/399182660_Analysing_farmers'_perceptions_of_genetically_improved_G3_Rohu_Labeo_rohita_fish_culture_in_Mymensingh_District
45. Health indicators for Penaeus vannamei breeders. - ResearchGate, accessed March 8, 2026, https://www.researchgate.net/figure/Health-indicators-for-Penaeus-vannamei-breeders_tbl3_368802723
46. Opinion: Benefits of animal welfare in Indian aquaculture, accessed March 8, 2026, https://enaca.org/enclosure/?id=1177
47. Comparative efficacy of individual and combined anti-parasitic treatments for Argulosis control - Indian Agricultural Research Journals, accessed March 8, 2026, https://epubs.icar.org.in/index.php/IJF/article/view/136120
48. argulus treatment | fish farming in telugu - YouTube, accessed March 8, 2026, https://www.youtube.com/watch?v=nBwEAQSidHw
49. Vision-2030 - ICAR-CIFA, accessed March 8, 2026, https://cifa.nic.in/cifa/wp-content/uploads/2023/06/Vision_2030.pdf
50. Infection with Macrobrachium rosenbergii nodavirus (MrNV), accessed March 8, 2026, https://www.agriculture.gov.au/sites/default/files/documents/infection-macrobrachium-rosenbergii-nodavirus.pdf
51. White Tail Disease of Freshwater Prawn, Macrobrachium rosenbergii - PMC, accessed March 8, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC3550746/
52. INFECTION WITH MACROBRACHIUM ROSENBERGII NODAVIRUS (WHITE TAIL DISEASE) - WOAH, accessed March 8, 2026, https://www.woah.org/fileadmin/Home/fr/Health_standards/aahm/current/2.2.07._MrNv_WTD.pdf
53. WHITE TAIL DISEASE - WOAH, accessed March 8, 2026, https://www.woah.org/fileadmin/Home/eng/Health_standards/aahm/2009/2.2.06_WTD.pdf
54. White tail disease of Macrobrachium rosenbergii, accessed March 8, 2026, https://enaca.org/enclosure/?id=706
55. Immunodiagnostic kit for white tail disease of Giant freshwater prawn | Official website of ICAR-Central Institute of Fisheries Education, Mumbai, accessed March 8, 2026, https://www.cife.edu.in/immunodiagnostic-kit-white-tail-disease-giant-freshwater-prawn