Silver Carp (*Hypophthalmichthys molitrix*)

1. Taxonomy & Nomenclature

Scientific Name: *Hypophthalmichthys molitrix*[1]. **Family:** Xenocyprididae[2].
Global Common Names: Silver Carp, Silverfin[1].
Regional & Local Names:
- China/Mandarin: 白鲢 (*bái lián*)[3].
- Thailand/Thai: ลิ่น (Lin)[4].
- Malaysia/Malay: *ikan kap perak*[5].
- India/Hindi: रजत कार्प (*rajat kārp*, "silver carp")[6].
- India/Kannada: *Bellimeenu*[7].
- India/Telugu: *Vendigandi*[7].
- India/Tamil: *Vellikkendai* (வெள்ளிக்கெந்தை)[7].
- Philippines/Tagalog: "Pla pla" (used for tilapia/related cichlids)[8] (Tilapia families often colloquially called *pla pla*).
- Bangladesh/Bengali: সিলভার কার্প (*silbar kārp* – transliteration of English).

2. Visual Identification & Anatomy (For CV)

- **Primary Identifying Features:** Body deep and laterally compressed[9]. Large head with an upward-turned (superior) mouth; eyes set far forward on midline[9]. Body scales are very small (tiny cycloid scales) while head and opercular bones are scaleless[9]. Long dorsal and anal fins (spiny+soft) but no adipose fin[10]. A long smooth ventral keel (bone ridge) extends from head toward the vent[11].
- **Coloration & Markings:** Uniform silvery gray body[9][12]. Dorsally the shade may be olive-green (#808000) blending to silvery-white (#C0C0C0) on the belly[12]. No distinct stripes or spots; fins are gray to pale. Juveniles are bright silver; adults often appear slightly greenish on back. Lateral line scutes may glint iridescent.
- **Sexual Dimorphism:** Sexes appear alike (no obvious coloration or fin-shape differences)[13]. Males may develop small breeding tubercles or darker ventral hues, but these are subtle.
- **Size Metrics:** Typically **60–100 cm** total length; can reach **140 cm** and ~50 kg[14]. Adult weight up to ~27 kg (≈60 lbs) reported[13]. Common body depth ~20–30% of length. No legal minimum (invasive species, often no size limit).

- **Similar Species Look-alikes:** Bighead Carp (*Hypophthalmichthys nobilis*) – silver carp has smaller, more pointed head and uniformly silver body (bighead is more brownish with blotches)[11]. Also lacks the dark "crosshatch" blotches often seen in bighead. Silver carp's long ventral keel and upward mouth differentiate it from other cyprinids (grass carp have keeled belly too but grass carp lack the pronounced head and teeth structure).

3. Habitat & Environmental Parameters

- **Water Type:** Strictly freshwater, often schooling in open water. Tolerates low salinity (estuarine/brackish waters up to ~12 ppt)[15]. Potamodromous (migrates within freshwater).
- **Geographic Distribution: Native** to large river basins of eastern Asia (historically from Amur River in Siberia through China's Yangtze, Pearl, etc.)[16]. **Introduced** worldwide (at least 88 countries) for aquaculture and phytoplankton control[1][17]. Established invasively in North America, parts of Europe, Asia and Africa (e.g., North American Great Lakes region, Bangladesh, India, Australia). Considered invasive outside its native range[17].
- **Micro-Habitats:** Prefers large, slow-flowing rivers, lakes and reservoirs. Often found in open pelagic zones near vegetation edges and embayments. Spawn in flowing tributaries or shallow flooded areas. Schools concentrate near surface or mid-water rather than bottoms; juveniles often use dense aquatic vegetation for cover.
- **Depth Profile:** Mostly in upper 0–5 m of water column (surface-oriented)[18], occasionally down to ~20 m. Remains higher in water especially when feeding.
- **Optimal Water Conditions:** Tropical–subtropical species. Thrives at **22–28 °C**; tolerates 6–30 °C[15]. Spawning triggered by 22–28 °C temperatures in spring-summer[19]. Prefers neutral to slightly alkaline pH (≈6.5–8.0). Can tolerate low dissolved oxygen (~3 mg/L) better than many fish[15]. Does well in turbid, eutrophic waters (benefits from high plankton productivity).

4. Diet, Foraging & Behavior

- **Natural Diet:** Obligate filter-feeder. Ingests *phytoplankton* (algae) and *zooplankton* nearly exclusively[20]. Juveniles and adults have no stomach and feed continuously on plankton (particles down to ~4 µm). Also consume detrital organic matter. They do **not** prey on large animals; their fine gill rakers strain water for microscopic food[20].

- **Feeding Behavior:** Constantly swimming with mouth open, pumping water across gills to filter food[20]. School in mid- to upper-water columns and feed by suction-filtering. Rarely respond to typical baited hooks (they "bump" slowly sinking food rather than strike aggressively)[21]. Hence anglers often use disintegrating baits to mimic natural feeding.
- **Peak Activity:** Diurnal, with steady feeding throughout daylight hours (sunrise to dusk) to maximize plankton intake. Reduced activity at night. Feeding activity can increase after plankton blooms. Not strongly tied to lunar or tide cycles (freshwater fish), but may be affected by weather (low barometric pressure reduces feeding).
- **Behavior:** Highly schooling and surface-oriented. Forms large aggregations. Not territorial. Known for **leaping**: when startled (boat motor noise, fishermen disturbance), fish can *burst to surface and jump* up to 1–2 m above water[22]. Jumping is a defense reflex; it poses hazards (they can hit boats or people). Silver carp can migrate long distances following river flows and flood pulses.

5. Tactical Fishing Guide & Gear

- **Best Live/Natural Baits:** Very limited; essentially needs particulate feed. A "suspension bait" is often used: a large dough ball (flour or cornmeal mixed and fermented) that slowly disintegrates, releasing a cloud of particles. Tiny hooks (size #10–12) or a cluster of them ("snare hook") are embedded in the ball[23]. Bread, corn, dough or bread-rolled dough (fed under cork float) can occasionally result in bumps. Live baits (worms, insects) are largely ineffective since the fish filter phytoplankton.
- **Best Artificial Lures:** Virtually none proven. As primarily filter feeders, silver carp seldom chase lures. Some anglers report snagging or bowfishing with weighted lines, but success is low. Big surface poppers or soft plastics rigged on multiple hook rigs might hook a feeding fish accidentally, but there are no "tried-and-true" lures.
- **Recommended Tackle:** Medium to heavy rod (fast action) with strong backbone (30–50 lb). Use a durable spinning or baitcasting reel with braided line (20–30 lb) for strength. Attach a sturdy monofilament leader (~30–50 lb) and a float (bobber) to suspend bait at the right level. Use small single hooks (sizes 10–6) or micro-tube jig rigs to hold particulate bait. In water, depth should correspond to where fish are schooling (often 1–3 m). Leader length can be short (~1 m) since fights are sudden and fish can jump.
- **Community & Regional Techniques:** In native Asian and introduced regions, silver carp are mostly harvested by commercial methods: gill nets,

cast nets, push nets, and trap nets in rivers and ponds. In India/Bangladesh, farmers often catch them with seines during drain-down of ponds, or use fish traps (bamboo corrals) in seasonal floods. Artisanal dip-netting during spawning runs is also done. Traditional rod-and-line angling is uncommon; where attempted, unique methods like the aforementioned "suspension bait" are used.

6. Handling, Safety & Regulations

- **Handling Warnings:** No venom or toxins, but exercise care. The gill covers (operculum) have sharp edges; wear gloves when unhooking. The fish thrashes strongly and may jump when lifted, so support the body horizontally. Because of their reputation to leap, always be cautious of fish in the boat.
- **Catch & Release:** If releasing, use wet hands or a wet cloth to handle. Remove hooks quickly (barbless hooks help) and keep fish in water as long as possible. Support the carp horizontally by mid-body, gently move it forward/backward in water to oxygenate gills before letting it swim off. Avoid prolonged air exposure, as filter carp can suffer quickly.
- **Conservation Status & Regulations:** In most areas, silver carp is **not protected** and often declared invasive. *IUCN*: *Near Threatened* in native range[24] (overfishing and habitat loss are concerns in China). In introduced regions, it is *highly regulated or banned*: for example, the U.S. lists live silver carp as an injurious species (illegal to import or transport)[25]. There are typically no seasonal closures (as they spawn year-round in warm climates), but check local rules. In many places, anglers are encouraged to remove and kill silver carp on capture due to invasive status.

7. Culinary & Commercial Profile

- **Commercial Value:** One of the world's top aquaculture species. Global production is enormous – only grass carp yields more tonnage[26]. In Asia (China, Bangladesh, India), it is farmed for food and as live transport fish. Market price is generally low (bony nature reduces value), but it is an important cheap protein. In Western markets, it is virtually unwanted (often used for fishmeal or pet food).
- **Meat Quality:** Flesh is lean and white but contains numerous fine intermuscular bones[27]. Texture is medium-firm; flavor mild and slightly "muddy" (often described as bland or meaty). There is a narrow strip of darker "red meat" along the lateral line, often removed as it is stronger-

tasting[27]. Overall, not highly prized compared to other carp due to bone density.

- **Regional Preparation:** In China and Vietnam, often steamed or poached with aromatic sauces (ginger, scallion). In South Asia, it is used in curries or fried after marinating in spices. For example, in Bangladesh and West Bengal (India) silver carp (জালুকা / silver carp) is cut into steaks for fish curries. In Kerala (India), pieces may be cooked with tamarind and tapioca. Generally, heavy spices or souring agents (tamarind, tomato) are used to offset any earthy taste. In the U.S., where it has invaded waters, there have been efforts to popularize it via fish cakes or sliders to overcome bony fillets.

8. Simulated Fisherman Q&A

Q1: *How can I tell a silver carp apart from a bighead carp?*
A: Look at head shape and color. Silver carp has a **smaller, upswept head** and a uniformly silvery body, whereas bighead carp has a massive head and a greenish-brown body with irregular blotches. Silver carp's long, smooth ventral keel and an upward-turned mouth are diagnostic[11]. Bighead carp lacks these features and has darker spotting.

Q2: *What should I use as bait to catch silver carp?*
A: Traditional baits won't work – silver carp filter plankton. Anglers use a special "suspension" bait: a large ball of dough or flour-based mix that slowly dissolves, releasing fine particles[23]. Tiny hooks are embedded in the bait and it is suspended under a float. As the carp sip the falling plankton, they bump the bait and hook themselves. Other than that, corn or bread dough chunks on a hook sometimes produce a tentative bite, but success is low[23].

Q3: *When and where are silver carp most active?*
A: They feed year-round in warm water, but peak activity is in spring–summer when plankton blooms. You'll find them schooling near the **surface of slow rivers, lakes or reservoirs**, often near vegetation edges or muddy inflows. They feed throughout daylight; calm, sunny days with high plankton levels often yield the most aggressive feeding. After heavy rain or during floods, they may move into flooded areas to spawn, so by late spring/early summer they can be plentiful in backwaters.

Q4: *Are silver carp dangerous or regulated?*
A: Silver carp are **harmless to people** (no venom or poison). The only hazard is if you're in a boat – startled silver carp will jump straight up and can be quite heavy, so they could knock you or your gear. They are **not endangered**; in fact, in many

countries they are invasive. For example, the U.S. bans import/transport of live silver carp as injurious[25]. Always check local regulations, but often fishermen are encouraged to kill silver carp if caught.

Q5: *What do silver carp eat in the wild?*
A: Primarily tiny plankton. They filter-feed on algae and microscopic animals in the water[20]. Juvenile and adult silver carp continually swim with their mouths open, trapping phytoplankton on their gill rakers. They will also ingest organic detritus. They **do not** chase large prey – think of them as living strainers. This diet means they rarely take angler bait unless it mimics their natural food (hence the dough-bait technique).

Silver Perch (*Leiopotherapon plumbeus*)

1. Taxonomy & Nomenclature

Scientific Name: *Leiopotherapon plumbeus* (Kner, 1864)[28]. **Family:** Terapontidae (grunters)[28].
Global Common Names: Silver Perch (Philippines)[28]; *Ayungin* (Tagalog)[29]; *Ayungin* or *Bugaong/Bigong/Bagaong* (Philippines)[29].
Regional & Local Names: Philippines/Tagalog: Ayungin, Bugaong (বাগাওং)/Bigaong/Bagaong[29]. (These are Filipino local names for this endemic species.) No widely used names in India or Thai; it is endemic to the Philippines.

2. Visual Identification & Anatomy (For CV)

- **Primary Identifying Features:** Moderately deep and compressed body. Slender profile with small head. Standard cichlid shape with continuous dorsal fin. Body without prominent spines or barbels. Lateral line continuous. It has a single eye stripe (not obvious) and a typically upward-curved mouth. Fins are unremarkable, with dorsal fin spanning much of the back.
- **Coloration & Markings:** Generally silver-gray overall, sometimes with a faint olive sheen on the back. May exhibit 6–7 faint vertical dark bars on juveniles (fading with age). Belly and lower flanks silvery white. Fins are translucent or pale gray. (This fish's common name derives from its shiny silver color.)
- **Sexual Dimorphism:** Males and females look very similar. During breeding, males may develop more intense coloration or slight tubercles on the head, but differences are subtle. Both guard eggs.

- **Size Metrics:** Relatively small. Maximum reported length **~16 cm** (6 inches) total[30]. Typically found 10–15 cm. Weight is very modest (on the order of ~50–100 g). No size regulations (caught mainly for food, often sold whole).
- **Similar Species Look-alikes:** Could be confused with other small therapy perches (e.g., *Leiopotherapon plumbeus* vs. introduced tilapia fry), but distinctive range (Philippine lakes) and parental care behavior sets it apart. The silvery body may recall juvenile tilapia, but *L. plumbeus* has a more elongated shape and lacks tilapia's thick lips.

3. Habitat & Environmental Parameters

- **Water Type:** Freshwater (lakes and rivers). Cannot tolerate high salinity. Known to ascend from brackish spawning sites but primarily a freshwater species[31].
- **Geographic Distribution: Endemic to the Philippines**[29], especially abundant historically in Laguna de Bay and other lowland lakes/rivers. No verified natural populations outside the Philippines. (Not reported as introduced elsewhere.)
- **Micro-Habitats:** Prefers clear to moderately turbid streams, river channels, and shallow lakes. Adults often found in fast-flowing areas just below rapids or weirs[31]. Also common in shorelines with aquatic plants. During rainy season, may move into flooded fields and backwaters.
- **Depth Profile:** Adults favor surface to mid-depths (0–5 m) near flow. Juveniles often in shallower edge habitats. Spawns in mid-water near cover, depositing eggs on vegetation or substrate.
- **Optimal Water Conditions:** Tropical species. Tolerates a broad temperature range (2–37 °C reported[31]), with optimal breeding in warm summer waters. Prefers well-oxygenated waters; sometimes migrates upstream for spawning in warmer months (March–May) at ~23–30 °C[31]. Likely prefers pH ~6.5–7.5. Its decline is attributed to habitat siltation and pollution, so it favors relatively clean waters (over sedimented ponds).

4. Diet, Foraging & Behavior

- **Natural Diet:** Omnivorous/carnivorous. Consumes benthic and aquatic invertebrates – insects, insect larvae, small mollusks, crustaceans, worms (earthworms, insect larvae) and some plant matter[31]. Young feed on zooplankton and small insects; adults will take larger prey like snails and aquatic larvae. Capable of feeding at various levels (bottom and midwater).

- **Feeding Behavior:** Active foragers, often picking food from bottom and among vegetation. They may also feed on surface insects fallen in water. Not a filter feeder – they pursue discrete prey. Usually feed singly or in small groups.
- **Peak Activity:** Active throughout the day; likely peaks during early morning and late afternoon when water temperatures are moderate. Spawning occurs in early summer afternoons[32]. Nighttime activity is reduced. As a lake species, they may be more active on cloudy days.
- **Behavior:** Parents guard eggs and fry (males and females build a nest and guard young)[33]. They may become aggressive around nest sites. Otherwise, generally schooling and timid when small; larger individuals can be bold in feeding. After spawning, adults often remain in the area until fry free-swim. The species does not leap or show any special defensive behaviors beyond schooling.

5. Tactical Fishing Guide & Gear

- **Best Live/Natural Baits:** Earthworms, small insects (grasshoppers), or small minnows. These fish readily take worms on small hooks. Pile grass or vegetation on riverbanks and fish nearby to attract them. They also hit dough balls or baits scented with garlic or shrimp (reflecting their omnivory). Fishing near the bottom or in cover will yield fish.
- **Best Artificial Lures:** Small soft plastics or grubs (1–2 inch) on light jigheads can be effective. Also small spinners or spoons (size 0–1) worked slowly can provoke strikes. Fly anglers use nymphs or small streamers (olive/black) when wading rivers. Bright colored beads or mimic insects might attract juvenile fish.
- **Recommended Tackle:** Light-action spinning rod (UL to L action) with 4–6 lb test line. A small reel with light monofilament (6–10 lb) or braid (6–10 lb) is suitable. Use small hooks (#10–14). A wading net is handy due to their small size. In Philippine markets the catch is typically by net; for angling, a quiet finesse setup is best.
- **Community & Regional Techniques:** Traditional fishing for Ayungin included cast nets and fish traps (bamboos) in rivers and lakes. Tribal hunters may spear them in shallow creeks during spawning runs. In some villages, night-time lamps (light fishing) attract them near the shore. No specialized indigenous gear is known beyond general freshwater fishing methods.

6. Handling, Safety & Regulations

- **Handling Warnings:** No special hazards; small cichlid with soft fins. Spines in dorsal/anal fins are mild – simply avoid hands near fins.
- **Catch & Release:** Given its declining status, catch-and-release is encouraged if not needed for food. Handle gently, wet net/hands to protect slime. Avoid prolonged air exposure; quickly revive by gently moving fish in water. Because they guard nests, returning a freshly caught specimen might harm eggs/fry if thrown back near a nest – be mindful of breeding areas.
- **Conservation Status & Regulations:** Classified as *Endangered* or *Rare* in local assessments (populations have plummeted from overfishing and habitat loss[34]). No global IUCN listing (Philippine endemic). Fishing pressure is now regulated in parts of the Philippines (some lakes have seasonal bans). Due to scarcity, it is a prized food fish and currently protected from overharvest in some areas via community rules.

7. Culinary & Commercial Profile

- **Commercial Value:** Highly valued in the Philippines as a food fish. Historically one of the top freshwater fish of Laguna de Bay. Currently scarce and fetches high price as a delicacy[34]. It is not farmed on a large scale (though some hatchery-rearing exists for restoration efforts) – mainly wild-caught. Outside the Philippines it has negligible commercial importance.
- **Meat Quality:** Flesh is white, soft and reportedly very delicious (often described as "delicate"). Very few intramuscular bones (unlike carps)[34]. The texture is fine-flaked. Because of its sweet flavor and small size, tilapia is often fried whole or used in clear fish soup.
- **Regional Preparation:** In the Philippines, often prepared fried or steamed, or in sour soups (e.g., *sinigang* style) to highlight the delicate flesh. Sometimes grilled lightly. It is considered one of the best-tasting native fish. Because it is expensive, local chefs feature it in special dishes, and folk medicine sometimes uses it to nourish nursing mothers (traditional belief).

8. Simulated Fisherman Q&A

Q1: *What is an "ayungin"?*
A: "Ayungin" is the Tagalog (Filipino) name for the endemic Philippine silver perch, *Leiopotherapon plumbeus*[29]. It's a prized small native fish (also called bugawng/bigaong locally) known for its silvery body and tasty flesh[34].

Q2: *Where do I find silver perch in Philippines waters?*
A: They inhabit lakes, reservoirs and rivers in Luzon (e.g., Laguna de Bay) and nearby islands. They prefer clear, flowing water near the surface[31]. Look below rapids, weirs and near vegetation, especially during breeding season (summer), when they move upstream.

Q3: *What do silver perch eat?*
A: They are opportunistic feeders. Adults eat aquatic insects, insect larvae, small mollusks and crustaceans[31]. They also take plant material and detritus. Juveniles will consume zooplankton and smaller insects.

Q4: *How should I cook ayungin for best taste?*
A: Ayungin is considered very delicious[34]. Locally, it's often simply fried with salt or steamed with ginger and vinegar to highlight its sweet flesh. In Laguna cuisine, pieces might be used in light fish soups or coconut milk stews. Because it's expensive, it's usually served fresh.

Q5: *Are silver perch endangered?*
A: Unfortunately yes, populations have crashed due to overfishing and habitat loss[34]. It is now rare in markets and has no special legal status nationally, but conservation efforts (captive breeding) are underway. Fishermen catching them are encouraged to release mature spawners to help recovery.

Snakehead (*Channa spp.* – Common Snakehead)

1. Taxonomy & Nomenclature

Scientific Name: Most common is *Channa striata* (striped snakehead) or *Channa punctata*, etc., **Family:** Channidae. (We address general Asian snakeheads, primarily *C. striata*.)
Global Common Names: Snakehead, Murrel (India), Mudfish. *Channa striata* is called Striped Snakehead or Asian snakehead.
Regional & Local Names: Very many local names[35]. Some examples:
- India/Hindi/North: **Garai** (गरई)[36] (also sometimes *Singhara* in colloquial Hindi).
- India/Bengali: **Shol** (শোল)[37] (West Bengal/Bangladesh).
- India/South: Malayalam: **Varaal** (വരാല്)[38], Tamil: **Viral** (விரால்), **Nedumural** (நெடுமுரல்)[39], Telugu: **Korramīnu** (కొర్రమీను)[40].
- Thailand/Thai: **Pla chon** (ปลาช่อน)[41].
- Philippines/Tagalog: **Dalag** or **Dalag** (generic for snakeheads)[42].
- Malaysia/Malay: **Gabus** or **Haruan**[43].

- Vietnam/Vietnamese: **Cá lóc** (often *Cá lóc đồng*)[44].
- China/Mandarin: 生鱼 or 泰国鳢 (names used for snakeheads in general)[44].
- Others: Burmese *ngayan*; Khmer *trey ross*[45]. Snakeheads are truly known by many local names across Asia.

2. Visual Identification & Anatomy (For CV)

- **Primary Identifying Features:** Elongated, cylindrical body with a large flat head. Long dorsal fin running most of the back; anal fin also elongated. No adipose fin, but distinct spinous/soft dorsal and anal fins. Jaw is broad with sharp canine teeth. Body shape resembles a snake (hence the name). Scales are smooth. "Grunting" when removed from water is characteristic.
- **Coloration & Markings:** Adults are generally olive-brown to dark brown on back, fading to yellowish belly[33]. Striped snakehead has faint black vertical bands or blotches across the sides, more distinct in juveniles. Males often darker overall. Dorsal and caudal fins may have small spots. Philippine snakehead (Channa striata) shows a chevron pattern on head (giving "chevron snakehead" name). Chromatically, #8B4513 (saddle brown) dorsal blending to #C0A050 (tan) ventrally, with dark banding.
- **Sexual Dimorphism:** Moderate. Males tend to be larger and may have a more uniform dark color, while females often have a paler belly. During breeding, both sexes develop spawning tubercles. Neither sex has conspicuous coloration like some fish.
- **Size Metrics:** *C. striata* can reach ~1 m (rarely, most are 40–60 cm)[46]. Weight up to ~10 kg reported. Many local snakeheads (e.g., Barca snakehead *C. barca*) are smaller (~30–40 cm). Legal size varies (often only catch-and-kill mandates), but breeders typically attain 40–60 cm.
- **Similar Species Look-alikes:** Other snakeheads (Channa) – Channa micropeltes (giant snakehead) is larger with bold zebra-like stripes; *C. striata* lacks the wide vertical bands. In Asia, no other freshwater fish resembles a snakehead's head shape and dorsal fin configuration. A moray eel might look snake-like underwater, but snakeheads have fins.

3. Habitat & Environmental Parameters

- **Water Type:** Freshwater (some species tolerate slightly brackish). These are air-breathing fishes found in ponds, rivers, swamps, and flooded fields. They can survive low oxygen by gulping air.
- **Geographic Distribution:** Native to South and Southeast Asia. *C. striata* range includes southern China, most of India, Bangladesh, Nepal, Sri Lanka,

Myanmar, Thailand, Cambodia, Vietnam, and Malaysia[47]. Also introduced to other Asian islands. *C. punctata* and others have overlapping ranges. Not native to Africa or Americas (although *Parachanna* spp. in Africa are similar). In many countries snakeheads are cultured or highly invasive.

- **Micro-Habitats:** Common snakehead (*C. striata*) is adaptable: occupies pools, canals, lakes and slow rivers. It breeds in shallow flooded areas; adults often move upstream into flooded rice fields during rainy season. They burrow into mud to survive dry season if water disappears[48]. Typically found near submerged vegetation or cover, but often venture into open water hunting.
- **Depth Profile:** Usually surface to mid-water (0–3 m). Juveniles often in very shallow, vegetated edges. Adults will also swim into deeper sections (up to 10 m) if prey is present.
- **Optimal Water Conditions:** Warm waters (23–30 °C for breeding)[32]. pH neutral (~7). They tolerate turbid, low-oxygen waters (using their labyrinth organ to breathe air). Seasonal flooding and dry-down cycles are part of their life cycle.

4. Diet, Foraging & Behavior

- **Natural Diet:** Predatory carnivore. Eats frogs, insects, crustaceans, and especially smaller fish[49]. Known to ambush prey – juveniles and adults will strike at anything moving. Opportunistic: will scavenge or even bite anglers. Diet may include waterfowl chicks or terrestrial rodents at water's edge.
- **Feeding Behavior:** Lurks near bottom or vegetation, then darts out. In breeding season they become particularly aggressive, guarding territory and attacking any intrusion[50]. Both parents guard eggs and fry, actively chasing away intruders. They school as juveniles but become solitary as adults. They are mainly nocturnal hunters but feed actively by day if undisturbed.
- **Peak Activity:** Dusk and night often see increased hunting, though *C. striata* feeds throughout day. Weather: barometric falls (storms) may trigger feeding. Flood season (summer) sees peak feeding to build energy for spawning.
- **Behavior:** Very hardy – can breathe air and survive in low water. In dry season they burrow in mud. Known to leap and make grunting sounds if handled[33]. Not migratory in long distances (except breeding migration into floodplains).

5. Tactical Fishing Guide & Gear

- **Best Live/Natural Baits:** Snakeheads strike at live baits (dead or alive). Frogs and large earthworms are legendary for attracting them. Small live fish (tilapia, carp) on a heavy hook work well. Chunks of chicken or fish fillets (squid, shrimp) can entice a bite. Baits should be large and dangled near cover or on bottom. In South Asia, fish caught are often used as cut-bait for snakeheads.
- **Best Artificial Lures:** Spinnerbaits, large topwater plugs (e.g., poppers, "frog" lures), and heavy soft plastics are effective. Bright or reflective lures mimic wounded fish well. Heavy metal spoons and large crankbaits (deep diving type) also catch snakeheads. Fly anglers use streamers or large poppers (black/olive are popular). Because of their sharp teeth, always use a steel leader when lure fishing.
- **Recommended Tackle:** Strong, fast-action rod (10–20 lb class) and reel; braided line 30–50 lb test with 60–80 lb fluorocarbon leader. Large circle hooks or treble hooks (size 4/0–7/0) for bait. Many prefer reel setups due to heavy fights and sharp jumps. Nets or gaffs are handy to land fish because snakeheads tend to thrash violently.
- **Community & Regional Techniques:** In South/Southeast Asia, snakeheads are often caught with spears or harpoons in flooded fields, and by funnel traps in streams. Modern gill-netting and hook-and-line are common. In Kerala (India), local cast nets (chorankaal) specifically target snakeheads in rivers. Bowfishing at night is practiced in some regions. Traditional angling for snakehead (murrel fishing) is considered an art; in rural Bengal, for example, fishermen use rods with long cane poles and horsehair lines to patiently tempt them.

6. Handling, Safety & Regulations

- **Handling Warnings:** Snakeheads have sharp gill rakers and many small sharp teeth; handle cautiously to avoid cuts. Always control the fish's head. Dorsal spines (though not poisonous) are strong; avoid grasping near fins. Because they can thrash hard, use a landing net.
- **Catch & Release:** If releasing, do so quickly. Hold the fish vertically in water and allow it to swim out. Because of their labyrinth organ, ensure gills are well aerated by moving fish gently until it swims away. Wild releases should ideally be upstream of capture point to prevent immediate recapture.
- **Conservation Status & Regulations:** Many *Channa* species (like *C. striata*) are assessed as **Least Concern (LC)**[51] due to wide distribution. They are

not endangered. Conversely, in places like North America they are prohibited/invasive. In native countries, snakehead are food fish, often with local harvest regulations (size limits or protected seasons). In India some states consider them game fish. Always verify local rules.

7. Culinary & Commercial Profile

- **Commercial Value:** Highly valued food fish across Asia[52]. Sold fresh or dried, and farmed in some regions. In Bangladesh and India, snakehead (locally *shol* or *garai*) is a major freshwater catch. **Economic importance**: high, as a delicacy – second only to carp in some markets. In Thailand and Vietnam, they are also cultured for food.
- **Meat Quality:** White, firm flesh with a rich flavor. Many pin bones but fillets are meaty. Considered nutritious ("wound-healing" in folk medicine[53]). Has been used in traditional soups for medicinal purposes.
- **Regional Preparation:** In India/Bengal, often cut into pieces and cooked in mustard sauce (*Shorshe chon*) or curries[54]. In Kerala (India) and Sri Lanka, snakehead curry with coconut and spices is popular[55]. In Thailand, snakehead is grilled or made into spicy soups (pla ra fermented fish sauce) [56]. In Philippines, snakehead (*dalag*) is grilled or fried and used in sour soup (*sinigang*). In Myanmar and Cambodia, salted and dried snakehead is eaten (with rice)[57][58].

8. Simulated Fisherman Q&A

Q1: *How can I recognize a snakehead in my catch?*
A: Snakeheads have elongated "eel-like" bodies and a snake-like head. They lack a dorsal adipose fin, but have a long dorsal fin running most of the back. The head is broad and flattened with a large mouth filled with canine teeth. Coloration is typically brownish with darker bands or blotches[33]. If the fish gasps for air at the surface, that's a giveaway too (their labyrinth organ lets them breathe air).

Q2: *What bait or lure should I use for snakehead?*
A: Live prey baits are best – e.g., a hooked frog, a small fish, or large worms. They'll strike aggressively at anything that looks like prey. For artificial lures, try large topwater plugs (frog lures) or heavy spinnerbaits. Bright colors (yellow/orange) or mimics of wounded fish often work. Snakeheads are strong and have sharp teeth, so use heavy line and a wire leader to prevent bite-offs.

Q3: *What do snakeheads eat?*
A: They are top predators that eat mainly other animals: fish, frogs, large aquatic

insects, and even small water birds or rodents if they can catch them. Juveniles start with insects and small crustaceans, but adults prefer fish and amphibians[49]. In habitats with many weeds, they'll lurk and ambush anything that comes near.

Q4: *I heard snakehead soup is good for wounds. Is that true?*
A: Yes, in traditional Chinese and Southeast Asian medicine, snakehead fish (in broth) is believed to promote healing, especially for wounds and postpartum recovery[53]. Scientifically, it's very nutritious and gelatinous, which might help tissue repair. That's why snakehead is often called "healing fish" or "medicine fish" in markets.

Q5: *What are snakehead fish called in my region?*
A: Local names vary. For example, in Bengali and Bangladesh they're called **শোল** (*shol*)[37]. In Tamil Nadu they're called விரால் (*Viral*) or செலுமுரல் (*Selumural*)[39]. In Kerala they call it വരാല് (*Varaal*)[38]. In Thailand it's **ปลาช่อน** (*pla chon*)[41], in Malaysia **haruan**[43] or **gabus**, and in the Philippines **dalag**. A dozen other local names exist (e.g., *garai* in North India[36]). If you describe the fish (snake-like, brown, large mouth) to a local fisherman, they'll know it by the regional name.

Tenpounder (*Elops machnata*)

1. Taxonomy & Nomenclature

Scientific Name: *Elops machnata* (Forsskål, 1775). **Family:** Elopidae (tenpounders)[59].
Global Common Names: Indo-Pacific Tenpounder, Ladyfish (though true ladyfish are *Elops spp.* commonly in Atlantic). In South Asia sometimes called simply "ten-pounder."
Regional & Local Names: In Kerala (India), local name: വള്ളിപ്പൂമിന് (*vaḷḷippūmīn*)[60] (Malayalam). Elsewhere, English "tenpounder" or "ladyfish" is used. No well-known names in Hindi/Bengali/Tagalog (not native to freshwater there).

2. Visual Identification & Anatomy (For CV)

- **Primary Identifying Features:** Elongated, fusiform body with deeply forked tail; very silvery shine. No spines in dorsal or anal fins (all soft rays). Body cross-section is almost oval[61]. A single long dorsal fin far back on body. Scales large and cycloid, giving a very reflective surface. Typical tarpon-like shape but smaller and without the tarpon's gular plate. The mouth is terminal.

- **Coloration & Markings:** Highly reflective silver body (hence sometimes called “silver salmon” in some regions). Back often greenish or bluish-tinged; fins usually gray. Juveniles slightly darker with vertical striping. In life, it appears bright metallic and can almost disappear by reflection. (For CV: RGB ~**#C0C0C0** silver overall, dorsal tinted **#789262** light olive, eyes black.) No distinctive markings or bars.
- **Sexual Dimorphism:** None obvious. Males and females appear the same externally. Both sexes grow the same form.
- **Size Metrics:** Max reported length ~118 cm fork length (≈120 cm total)[62]; common catches are 30–60 cm. Weight up to ~10.8 kg[62]. Average size in catches is often ~1–3 kg. No minimum size laws generally, but gamefish anglers often keep only 30–50 cm.
- **Similar Species Look-alikes:** Similar to Atlantic ladyfish (*Elops saurus*) but that is on opposite coast. May be confused with small tarpon (family Megalopidae) by layman, but tenpounders lack tarpon’s gular plate and have fewer dorsal rays (20–27 vs tarpon’s 47). A long slender barracuda could look similar, but barracudas have large teeth and distinctly different head shape.

3. Habitat & Environmental Parameters

- **Water Type:** Marine and estuarine. Often moves into lagoons and bays, occasionally enters lower estuaries[63]. Does *not* inhabit true freshwater. Pelagic-neritic (open coastal) and oceanodromous (migrates in ocean and coastal waters)[64].
- **Geographic Distribution:** Tropical Indo-West Pacific. From South Africa (Mossel Bay) through Red Sea, Indian Ocean, up to 38°N (Japan) and down to 37°S[65]. Off India, found along entire coast. Not in the Atlantic (Pacific/Indian only). Often found near continental shelf waters.
- **Micro-Habitats:** Adults usually in shallow coastal waters, reefs, and harbors. Juveniles migrate into **brackish estuaries and lagoons** where they grow[63]. Spawn offshore in open water; larvae drift into estuaries. No reef association. Depths from near surface to ~100 m.
- **Depth Profile:** Commonly in upper water column. Often near surface at night (feeding)[63]. They can be caught at depths from just below surface down to ~30 m nearshore.
- **Optimal Water Conditions:** Prefers warm coastal waters (tropical climate). Spawning presumably offshore in warm currents (no specific temp given). It will occur in salinities from brackish to full marine. Highly tolerant of a range

of conditions – reported in salinities up to seawater and temperatures ~20–30 °C.

4. Diet, Foraging & Behavior

- **Natural Diet:** Carnivorous, feeding mainly on smaller fish and crustaceans[63]. Young tenpounders (and similar ladyfish) feed on zooplankton and small insects, but adults are voracious piscivores. Gut-content studies show many small fish, shrimp and squid.
- **Feeding Behavior:** Active swimmers that chase down prey. They have keen vision and take baitfish near the surface at dawn/dusk. Because they school when young, juveniles may filter-feed on plankton to start, then switch to fish. Anglers note they will eagerly take fast-moving lures.
- **Peak Activity:** Often feed most heavily at night or in low light (crepuscular) when baitfish are abundant near surface. Also known to feed around coastal structure at dawn. Low barometer or tidal flow can increase feeding.
- **Behavior:** Known to form small schools in open water, especially juveniles. Long-distance migrations unknown (non-migratory beyond local schooling). No special defensive behaviors; easily caught on standard gear.

5. Tactical Fishing Guide & Gear

- **Best Live/Natural Baits:** Since they favor live prey, anglers use live shrimp or small finfish. Strips of fresh squid or fish fillet can also work. Night-time fishing with shrimp under a float is a common method. They have also been known to hit clusters of small baitfish (though rigging this is hard for anglers).
- **Best Artificial Lures:** They hit surface and subsurface plugs. *Topwater plugs* (walking-style) mimic fleeing baitfish well. *Slow-sinking twitchbaits* in shiny patterns also attract strikes. Large jigs tipped with feather can provoke strikes. Anglers often use spoons, small swimbaits or metal jigs cast into schooling fish. Reflective silver/white lures match their natural prey shine.
- **Recommended Tackle:** Medium-power saltwater rod (8–15 kg class) with a fast action tip. Reels spooled with 15–30 lb braided line (20–50 lb monofilament leader). Treble or big single hooks (size 1/0–3/0) for lures. For bait, use circle hooks (4/0–6/0) so fish can be quickly released if needed. Because tenpounders are strong on the strike, gear should be robust. They jump and fight, so a net is useful for landing.
- **Community & Regional Techniques:** Not a typical target in India, but in some coastal regions they are caught by drift nets or gillnets set inshore. In

Kerala, beach seine nets and gill nets often trap them during migrations. Night fishing with lights and bait (cutbait or baits on longlines) is practiced in some coastal villages. Traditional methods are similar to those for catching small tarpon or ladyfish.

6. Handling, Safety & Regulations

- **Handling Warnings:** Generally harmless – they have no fangs or venom. Handle carefully to avoid cuts from sharp fins (fins are all soft-rayed, so minimal risk). As strong jumpers, watch that they don't flip out of boat on the strike.
- **Catch & Release:** Typically kept as food, but if releasing, do so quickly as they can desiccate out of water. Support the fish and slide it head-first back into the water. Because they have no special respiratory needs beyond gills, let water flow through gills if revived.
- **Conservation Status & Regulations:** *IUCN:* Least Concern[66], reflecting wide distribution and no major threats. No specific protections; often treated as a gamefish in some areas (e.g., bag limits in African countries where introduced). Not regulated in India beyond standard coastal fishing laws.

7. Culinary & Commercial Profile

- **Commercial Value:** Minor. Tenpounder flesh is edible but not highly prized due to bony frame. It is often regarded as a "pantry fish" in West Africa or Asia, eaten by local communities but not exported. Not a focus of commercial fisheries.
- **Meat Quality:** White, lean flesh but very **high in bones**[67]. Yields little fillet. Flavor is fairly mild but can be described as "fishy." The body cavity has many small rib bones, making filleting difficult. Because of the numerous pin bones, it is labor-intensive to process.
- **Regional Preparation:** In places where it is eaten (e.g. West Africa, parts of Asia), tenpounder is often smoked or salted to preserve (due to low fillet yield). In India, it's not commonly consumed, though a local fisherman might clean and fry small specimens. In Hawaiian cuisine *Elops hawaiensis* ("awaʻaua") has been eaten in salads or soups. Generally, cooking methods that help remove/dissolve small bones (like stewing until soft, pickling, or fermenting) are used.

8. Simulated Fisherman Q&A

Q1: *What waters do I find tenpounder (ladyfish)?*
A: These fish are found in **coastal marine and estuarine waters**, not rivers. Look for them in warm tropical seas – often schooling in shallow bays or near reefs. They enter lagoons and mangrove areas[63], especially juveniles, but adults stay in open water. In India, try near seashore piers or harbor lights at dusk.

Q2: *What do tenpounders eat, and what bait works?*
A: They're voracious predators of **small fish and crustaceans**[63]. Use live shrimp or small live fish as bait. Large shiny lures also work (they chase fast-moving prey), so topwater plugs and jigs that imitate baitfish can trigger strikes. They'll attack anything small that moves.

Q3: *Are tenpounders any good to eat?*
A: They're edible, but not tasty: "**not tasty and full of bones**"[67]. You get very little meat because of all the small rib bones. In practice, most anglers release them or use them as bait for larger species.

Q4: *How big do tenpounders get?*
A: Up to about **1.2 meters** (just under 4 feet)[62], though most we catch are 30–60 cm. They can weigh around 10 kg at maximum. Common catches are roughly 1–3 kg (2–6 lbs). Even a 70 cm tenpounder is a solid fighter on light gear.

Q5: *What local name should I ask for tenpounder in Kerala?*
A: In Kerala they call it വള്ളിപ്പുമിന് (*vaḷḷippūmīn*)[60] in Malayalam. If you ask a fisherman for "valippumin," they will know. (It literally means "worm fish," possibly referencing their bait preference.)

Tilapia (*Oreochromis* spp., primarily *O. niloticus*)

1. Taxonomy & Nomenclature

Scientific Name: Multiple species; commonly **Nile Tilapia** (*Oreochromis niloticus*) or **Mozambique Tilapia** (*O. mossambicus*). All are cichlids, formerly genus *Tilapia*.
Family: Cichlidae.
Global Common Names: Tilapia (general term). Specific species called Nile tilapia, Mozambique tilapia, Blue tilapia, etc.
Regional & Local Names:
- India/Hindi: टिलापिया (*tilapiyā*).
- India/Tamil: திலேப்பி (*tilēppi*)[68].

- Bangladesh/Bengali: টিলাপিয়া (*ṭilāpiyā*).
- Malaysia/Malay/Indonesian: **Ikan nila**[69].
- Thailand/Thai: ปลานิล (*pla nil*)[8].
- Philippines/Tagalog: *Tilapia* or **pla pla**[8] (especially for *O. niloticus*).
- Global: "Nile Tilapia" is common English, but many languages simply use "tilapia" or a loanword.

2. Visual Identification & Anatomy (For CV)

- **Primary Identifying Features:** Laterally compressed, deep body – almost oval in cross-section[70]. Long dorsal fin (with both spiny and soft rays) and long anal fin. Mouth is protrusible (capable of extending out)[70]. Lips are usually thick. No scales on the throat (guaranteed by pharyngeal jaws). They have a single continuous lateral line.
- **Coloration & Markings:** Typically a uniform gray-green to dark gray on back, fading to silvery on sides and white belly. Nile tilapia may have faint vertical bars (especially juveniles). Some strains have reddish fin margins. Tilapia aurea (blue tilapia) have a bluish tint. For O. niloticus, consider body color ~**#708070** (gray-green) to **#C0C0C0** (silver) ventrally. They often look drab unless nuptial males show orange on ventral fins.
- **Sexual Dimorphism:** Nile tilapia males (especially breeding males) develop a narrow red/orange band at the base of the tail and may have a deeper red belly. Males also build and guard the nest; females are drabber. Females may have two to six conspicuous vertical bars, males usually fewer. Otherwise, body shape is similar.
- **Size Metrics:** Up to ~60 cm total length (2 ft) in tropical waters[71]. Weight up to ~4–6 kg. Typically farmed fish are kept around 30 cm (1–2 kg). Sexually mature at ~20 cm. (Many regions impose a 30 cm or 1 kg size limit for harvesting.)
- **Similar Species Look-alikes:** Other cichlids: However, wild *Oreochromis niloticus* is distinguished by the fused pharyngeal jaws (a cichlid trait). It might be confused with *Tilapia zillii* or *Sarotherodon* species, but those are usually smaller and differently shaped. Non-cichlid look-alikes are rare due to the cichlid's unique body and mouth structure.

3. Habitat & Environmental Parameters

- **Water Type:** Freshwater; tolerates slight brackish (up to ~2–3 ppt) but not true seawater. Prefers still or slow waters: ponds, lakes, reservoirs,

rivers[72]. Eurythermal (broad temp range) and euryhaline (tolerant of variable salinity).

- **Geographic Distribution:** Native to Africa (Nile basin, etc.) and Middle East. Now **widely introduced globally** for aquaculture. Tropical/subtropical worldwide – including India, Southeast Asia, Philippines, South America, U.S. (Florida, etc.), and Africa outside native range. In warm climates, tilapia establish feral populations in many rivers and lakes. They are listed among worst invasive species in some regions[73].
- **Micro-Habitats:** Often concentrate in vegetated shallows in lakes/ponds. Spawn in holes dug near shore. Juveniles use weeds and rocks for cover. In large lakes, found in deeper open water during day if food is there. They prefer sunlit waters (warm, oxygenated).
- **Depth Profile:** Typically found near surface to 3–5 m. Will descend deeper (10–20 m) only if necessary (stratified lakes). School at mid-depths in reservoirs at night.
- **Optimal Water Conditions:** Warm water 22–28 °C; do poorly below 15–18 °C (mortality below 10 °C)[74]. Neutral to alkaline pH. Extremely tolerant of poor conditions: can thrive in low O_2 (gasping air), and even high salinity in tropical climates (some live in brackish lagoons and Salton Sea-level salt). Not cold-hardy. Often found where other fish cannot survive.

4. Diet, Foraging & Behavior

- **Natural Diet:** Tilapia are primarily **herbivorous/omnivorous**. They feed on phytoplankton, algae (periphyton), aquatic plants, and detritus, processing it with strong pharyngeal jaws[70]. They also consume small invertebrates (insect larvae, crustaceans) opportunistically, especially when young. In captivity they eat formulated feeds. Wild tilapia filter microorganisms and grind plant tissue.
- **Feeding Behavior:** Continuous grazing – they will ram algae-covered surfaces or filter plankton from water. Because of the split pharyngeal jaws[70], they can both scrape and crush a variety of foods. They often feed in groups, stirring up sediment to access microbes. At night or when cooler, they slow down but may feed on plankton brought to surface.
- **Peak Activity:** Daytime feeders (herbivore fish). Peak grazing in sunlight. Morning and late afternoon are active feeding times; night feeding is minimal. In eutrophic water, they may feed round-the-clock on abundant algae.

- **Behavior:** Generally peaceful, schooling fish. In ponds they form hierarchies. During spawning, they become territorial – males dig nests and court females in elaborate displays. Fry are mouthbrooded by mother (Oreochromis) or not (Sarotherodon), so parental care exists. Tilapia can jump out of small streams during floods.

5. Tactical Fishing Guide & Gear

- **Best Live/Natural Baits:** In many areas, bread dough balls, corn, or bread are the top baits (for example, dough balls are common bait for tilapia). They readily take dough, bread, and grains. Attractants like cheese, banana, or sweet dough work. Small worms or insects are also effective due to their omnivory. Use light hooks (#6–10) near banks. Floating or sinking baits both work.
- **Best Artificial Lures:** Small spinners, flies, and poppers sized for panfish work surprisingly well. Tilapia will take small soft plastics (grubs, paddletails ~2″). Fly anglers use foam flies or emergers (imitating insects) with good success. Brightly colored baitfish imitations (white/yellow) also attract bites. The lure's motion should be slow to imitate sluggish prey.
- **Recommended Tackle:** Ultralight to light tackle: a 4–6 ft rod, 2–6 lb test line is sufficient. Use fluorocarbon leaders (4–6 lb) so the line sinks slightly (tilapia often feed subsurface). Small hooks (#10–14) or tiny treble hooks (for flies). A bobber/floater rig is traditional, but weighted rigs also catch them off bottom. They fight mildly – even ultralight rigs are fun for tilapia.
- **Community & Regional Techniques:** In Asia, tilapia are often caught by cast nets and seine nets in village ponds. They are also susceptible to trapping in fish baskets (nila vala in South India) and even caught by hand during pond draining. Angling methods include float fishing at rice fields or canal edges. In the Philippines, tilapia (molmol, pla pla) are often caught in nets or pond drawdowns. In African cultures, tilapia is also speared in shallow water at night (as it gathers plankton).

6. Handling, Safety & Regulations

- **Handling Warnings:** No venom; dorsal fin spines are minimal (cichlids have rigid first rays, but not dangerous). Scales and small spines on fins can prick; handle carefully around gill covers. Use wet hands to keep protective slime coat intact if releasing.
- **Catch & Release:** Widely practiced in sport fishing; tilapia recover easily. Wet-hold and release quickly. Ensure minimal air exposure; they are hardy

but still best kept wet. Minimize deep hooking by using fine hooks or treble lures.

- **Conservation Status & Regulations:** Most tilapia species (Nile, Blue) are listed as *Near Threatened* or *Least Concern*[75]. In many non-native regions, tilapia are invasive and are heavily controlled (e.g., US regulations in southern states). In native regions, they are farmed and fished intensively. Where invasive, there may be bans on stocking. Check local rules: some lakes only allow tilapia fishing (to reduce invasives), others ban possession. In general, they are not endangered and often overabundant.

7. Culinary & Commercial Profile

- **Commercial Value:** One of the world's most important aquaculture fish. Over 6 million metric tons are farmed globally (2nd only to carp)[76]. Primary uses: table fish fillets for retail (fresh/frozen), and increasingly as processed products (fish cakes, fillets in markets). In Africa/Asia, tilapia is a cheap staple protein. In developed markets (USA, Europe), it's sold as a "white fish" fillet.
- **Meat Quality:** Flesh is white and flaky but mild and sometimes bland. Contains numerous small pin bones and fine intramuscular bones. Texture is slightly mushy unless very fresh. Nutritionally decent (high protein), but omega-3 content is lower than coldwater fish. Over-farming has led to nutritional debates (see 'FishFrenzyFacts').
- **Regional Preparation:** In the Philippines, tilapia is traditionally fried or grilled and eaten with rice; it's the fourth-most consumed fish there[76]. In India, it's often made into curries or fried as a budget fish. Common preparations include fish fry (tilapia coated in spice batter and shallow-fried) or fish curry (in coconut or tamarind sauce). In Thailand/Malaysia, tilapia may be steamed or made into fish cakes. Tilapia's mild taste means it takes on spicy marinades and sauces well. It's also used for aquaponics and fish fillet exports (broiling, baking).

8. Simulated Fisherman Q&A

Q1: *What do tilapia eat?*
A: Tilapia are largely herbivorous/omnivorous. They graze on algae, phytoplankton and aquatic plants[70], plus they'll eat detritus and small invertebrates (worms, insect larvae) when available. Their fused pharyngeal jaws let them crush plant matter efficiently[70].

Q2: *Are tilapia invasive?*
A: Yes, in many warm regions tilapia are major invasive pests. They're on the IUCN's list of worst invasive freshwater species[73]. If you fish in southern U.S., Australia, or Pacific islands, you'll see them outcompete native fish. In those places, regulations often require killing tilapia caught.

Q3: *How big do they get?*
A: Nile tilapia can grow up to ~**60 cm** (2 feet)[71], though 30–40 cm (1–1.3 ft) is common in ponds. Around 2–4 kg is a large adult. They reach maturity at ~20 cm, so you'll get a bite on very small hooks!

Q4: *What's a common local name for tilapia?*
A: In Thailand, tilapia is called **pla nil** (ปลานิล)[8]. In Malay/Indonesia, it's **ikan nila**[69]. In India, simply "tilapiya" (टिलापिया) or in Tamil திலேப்பி[68]. In Bengali, **টিলাপিয়া**. In the Philippines, **tilapia** (or locally "pla pla" as a generic term)[8].

Q5: *How do I catch tilapia on the hook?*
A: They bite on simple baits. Anglers report success with bread balls or dough under a float near weeds, or with small worms on a hook. Light spinners or fly-imitation jigs also work. Because they're usually in shallow water, try topwater lures or let the bait drift under a bobber. They are not picky biters, so light tackle is key.

---

[1] [2] [14] [17] [20] [24] [26] Silver carp - Wikipedia

https://en.wikipedia.org/wiki/Silver_carp

[3] [6] Hypophthalmichthys molitrix - Agrovoc - AGROVOC

https://agrovoc.review.fao.org/browse/agrovoc/en/page/c_35400?clang=vi

[4] silver-carp คืออะไร แปลว่าอะไร ตัวอย่างประโยค จากพจนานุกรมข้อมูลภาพปลาและสัตว์น้ำไทย

https://www.sanook.com/dictionary/dict/dict-fish/search/silver-carp/

[5] Fish Species Names in Malay | PDF

https://www.scribd.com/doc/110682710/Fish-in-Malaysia

[7] [27] Carp silver | Seafood India

https://seafood-india.com/carp-silver

[8] [68] [69] Common Names List - Oreochromis niloticus

https://fishbase.se/ComNames/2

[9] [11] [25] Silver Carp (Hypophthalmichthys molitrix) - Species Profile

https://nas.er.usgs.gov/queries/factsheet.aspx?speciesID=549

[10] [12] [13] [15] [16] [19] Hypophthalmichthys molitrix (Carp) | INFORMATION | Animal Diversity Web

https://animaldiversity.org/accounts/Hypophthalmichthys_molitrix/

[18] Hypophthalmichthys molitrix, Silver carp : fisheries, aquaculture

https://www.fishbase.se/country/CountrySpeciesSummary.php?id=274&c_code=356

[21] [23] Targeting Asian Carp (Silver Carp) - Other Fish Species - Bass Fishing Forums

https://www.bassresource.com/bass-fishing-forums/topic/113450-targeting-asian-carp-silver-carp/

[22] Silver Carp - Montana Field Guide

https://fieldguide.mt.gov/speciesDetail.aspx?elcode=AFCJB44010

[28] [29] [30] [34] Leiopotherapon plumbeus - Wikipedia

https://en.wikipedia.org/wiki/Leiopotherapon_plumbeus

[31] [32] Bidyanus bidyanus, Silver perch : fisheries, aquaculture, gamefish

https://fishbase.se/summary/Bidyanus-bidyanus

[33] [35] [36] [37] [38] [39] [40] [41] [42] [43] [44] [45] [46] [47] [48] [49] [50] [51] [52] [53] [54] [55] [56] [57] [58] Channa striata - Wikipedia

https://en.wikipedia.org/wiki/Channa_striata

[59] [61] [62] [63] [64] [65] [66] [67] Elops machnata, Tenpounder : fisheries, gamefish

https://www.fishbase.se/summary/Elops-machnata

[60] (PDF) A checklist of fishes of Kerala, India

https://www.researchgate.net/publication/284166826_A_checklist_of_fishes_of_Kerala_India

[70] [71] [72] [73] [74] [75] [76] Tilapia - Wikipedia

https://en.wikipedia.org/wiki/Tilapia